

The Volta River Basin

Comprehensive Assessment of Water Management in Agriculture
Comparative study of river basin development and management

Boubacar BARRY
Emmanuel OBUOBIE
Marc ANDREINI
Winston ANDAH
Mathilde PLUQUET

January, 2005

Draft

TABLE OF CONTENT

List of Tables.....	7
List of Figures.....	9
I. INTRODUCTION.....	11
I.1 The Volta River Basin.....	11
I.2 Methodology and presentation of the study.....	14
II.THE VOLTA RIVER BASIN: HUMAN AND PHYSICAL SETTING.....	16
II.1 Climatic features.....	16
II.1.1 Distribution of Rainfall.....	16
II.1.2 Distribution of Evaporation and Evapotranspiration (ETP) compared to rainfall.....	22
II.1.3 Temperature and humidity.....	26
II.2 Human Context.....	29
II.3 Economic Sector and Policies.....	37
II.4 Agro-ecological Zoning.....	41
II.4.1 Drainage network and sub-basins.....	41
II.4.2 Relief.....	43
II.4.4 Geology of the Basin.....	47
II.4.5 Land use, land ownership and land degradation.....	49
i. BENIN.....	52
ii. BURKINA FASO.....	53
iii. CÔTE D’IVOIRE.....	55
iv. GHANA.....	56
White Volta Basin.....	57
Black Volta Basin.....	58
Lower Volta Basin.....	58
Oti Basin.....	59
Daka Basin.....	59
v. MALI.....	59
vi. TOGO.....	59
II.4.6 Agriculture.....	61
i. Crop production.....	61
ii. Livestock.....	65
iii. Fisheries.....	66
iv. Forestry.....	67
II.5 Water resources.....	67
II.5.1 Surface Water Resources in the Volta River System.....	67
i. GHANA.....	67
• The White Volta Basin.....	69
• The Black Volta Basin.....	69
• The Lower Volta.....	70
• Oti Basin.....	72
• Daka basin.....	72
ii. BURKINA FASO.....	73
iii. MALI.....	77
iv. TOGO.....	78
v. CÔTE D’IVOIRE.....	78
vi. BENIN.....	78
II.5.2 Groundwater Resources, Recharge, and Quality.....	78

i.	GHANA	79
(b)	Aquifers found in Ghana.....	84
(c)	Borehole Yields	84
(d)	Groundwater abstraction and distribution.....	87
(e)	Aquifer recharge	88
(f)	Groundwater quality	89
ii.	Burkina Faso	90
(a)	Aquifers in Burkina Faso.....	90
(b)	Groundwater availability	91
(c)	Aquifer yield.....	92
(d)	Recharge Mechanisms	94
(e)	Groundwater Abstraction and Distribution.....	94
(f)	Quality of aquifers in Burkina Faso.....	95
•	Nitrogen species.....	96
•	Salinity	96
•	Fluoride.....	96
•	Arsenic	97
•	Iodine	97
II.4.2	A geographical triptych.....	97
A.	Agro-Ecological Zones in the Volta Basin of Ghana	97
•	The High Rainforest.....	100
•	The Semi-Deciduous Forest (SDF).....	100
•	The forest-Savanna Transitional Zone (Derived savanna)	100
•	Guinea Savanna Zone	101
•	Sudan Savanna Zone.....	101
•	Coastal Savanna Zone.....	102
B.	Major Farming Systems in the Volta Basin of Ghana	102
•	The bush fallow system	103
•	The HUZA farming system	104
•	The permanent systems.....	104
•	The compound farming system.....	104
•	The Mixed Farming System.....	104
C.	Land tenure characteristics in several agro-ecological zones of the Volta basin of Ghana	105
•	Sudan and Guinean Savanna Zones.....	105
•	Forest Savanna Transitional Zone	106
•	Deciduous Forest Zone	106
D.	Major farming activities in the various agro-ecological zones of the Volta Basin in Ghana	106
i.	Cropping systems in the Volta basin of Ghana.....	106
•	Sudan Savanna Zone.....	106
•	Guinean Savanna Zone (GS)	107
•	Forest Savanna Transitional Zone	107
•	Deciduous Forest Zone	107
ii.	Crop husbandry processes and activities	107
•	Sudan Savanna Zone.....	107
•	Guinea Savanna Zone	108
•	<i>Forest Savanna Transitional Zone</i>	109
•	Deciduous Forest Zone	110
iii.	Livestock production	110
•	Sudan Savanna Zone and Guinea Savanna Zone.....	110

• Forest Transitional Savanna Zone	110
• Deciduous Zone	111
• Savanna Zone.....	111
• Savanna Transitional Zone	111
• Deciduous Forest Zone	111
i. The interior savanna zone	112
• Farming systems	112
• Interface between land evaluation and farming systems	112
• Conclusion	113
ii. The Forest-Savanna Transitional Zone.....	113
• Farming systems	113
• Interface between land evaluation and farming systems in the Transitional zone	115
• Conclusions.....	116
iii. The Deciduous Forest Zone.....	116
• Farming systems	116
• Interface between land evaluation and farming systems in the Deciduous Forest	117
Zone	117
• Conclusion	118
B. The Agro-ecological zones in the Volta basin of Burkina Faso	118
• The Eastern zone.....	118
• The Sahelian zone	118
• The North-western zone.....	118
• The Central zone	119
• The Western zone	119
C. Agricultural regions and production systems in Burkina Faso	119
• The Northern Region: The SAHEL	119
• The Central Region	120
• The North-Western Region.....	120
• The Eastern Region.....	121
• The Western Region	121
III. HISTORICAL DEVELOPMENT OF VOLTA BASIN.....	122
III.1 Changes in water supply and use	122
III.1.1 Water Institutions and Legislation	122
Benin.....	122
Burkina Faso	123
Côte d'Ivoire	124
Ghana	124
Policy framework.....	124
Water vision.....	125
Policy Objectives:.....	125
Guiding Principles for Ghana Water Policy	125
Mali	127
4.6 Togo	128
Overview of national institutional and legal framework for integrated management.....	129
• Regional Coordination	129
• International Cooperation	131
Surface water development and use.....	132
Water use:	137
Surface Water and health issues:	139

Groundwater development and use.....	140
Mali	140
Burkina Faso	141
Ghana	142
III.2 Agrarian and environmental transformations	142
A. Technical change (agriculture, water mobilization.....	142
1. AGRICULTURE	142
1.1 Agricultural changes in the pre-colonial era of Ghana	142
1. Agricultural changes in the Volta Basin of Ghana	144
Irrigation opportunity.....	146
II. WATER MOBILIZATION.....	147
*Animal production in the Volta Basin of Ghana	147
1. Fisheries	147
E. Health and Environmental Changes in the Volta Basin of Ghana.....	148
I. HEALTH.....	148
f Prevailing diseases before the Akosombo Dam Construction.....	149
f Post construction period.....	149
II. ENVIRONMENTAL CHANGES	151
(i) Deforestation	152
(ii) Aquatic weeds	152
(iii) Sandbars	153
(iv) Seismicity	154
(v) Sediment load changes	154
(vi) Morphological changes	155
(vii) Microclimatic changes (weather changes).....	155
III. LAND TENURE SYSTEM AND AGRICULTURAL POLICY CHANGES IN BURKINA FASO	156
1. Changes in the land tenure system in Burkina Faso	156
(i) The customary land system.....	156
(ii) The colonial land system.....	156
(iii) The post-colonial land system.....	157
2. Changes in Agricultural Policy in Burkina Faso	157
III.3 Changes in population, social and cultural patterns of water management.....	160
Population Distribution in Ghana	160
Urban/Rural Disparities	161
Annual population growth rate in the Volta Basin of Ghana.....	162
Agricultural land use in the sub-basins.....	163
Migrations and mobility in the Volta basin of Ghana.....	164
IV.3.4 Legal Framework: Presentation of the Water Laws	166
A. Statutory Legal Framework for Water	166
A.1 National Water Policy.....	166
A.2 National Water Legislation	167
A.3 The Ghana Water Company Limited.....	168
A.4 The Community Water and Sanitation Agency	169
A.5 District Assemblies	169
A.6 Regulatory Institutions	170
A.6.1 The Water Resources Commission	172
A.6.2 The Environmental Protection Agency.....	173
A.6.3 The National Environmental Action Plan.....	173

A.6.4 The Public Utilities Regulatory Commission	173
B. Nature and Status of Customary Water Rights	174
B.1 Description of Customary Water Laws.....	174
IV. 7 Urban Water Management.....	175
A. Access to Clean Water in Urban Ghana	175
A.1 Problems of GWCL	177
A.2 Reform of Urban Water Sector	177
B. Access to Clean Water in Rural Ghana.....	178
B.1 Community Water & Sanitation Programme (CWSP)-objectives.....	178
B.2 Access to Water under the CWSP	178
B.3 Assessment of the Various Regimes.....	179
B.3.1 Assessing GWSC'S Performance	179
B.3.2 Assessing the CWSP.....	180
B.3.3 Assessing the Customary Regime.....	181
B.4 Impact of Water Law & Water Rights on Water Supply and Access.....	183
B.4.1 Enabling Conditions.....	183
C. CUSTOMARY LAW AND TRADITIONAL WATER MANAGEMENT IN BURKINA FASO:	
.....	184
EXAMPLE OF THE YATENGA PROVINCE	184
1 - The governing system in the Yatenga kingdom	185
1.1- Traditional governance in the Yatenga kingdom	185
1.2- Division of powers within the kingdom	185
2.- Water in the Traditional Moagha Society of the Yatenga kingdom.....	186
2.1- Water as a universal myth	186
2.2- Water and the animist religion.....	186
2.3- Water in the peoples' settlement	186
2.4- Water as salvation for the migrant.....	186
2.5- Village names with inspiration from water.	187
2.6- A system of water property	187
2.7- Traditional water management: customary laws	188
2.7.1- Prohibitions.....	188
2.7.2- Totems	188
2.8- Gender and traditional water management	188
2.9- The causes of failure of water conveyance projects	188
Bibliography	190

List of Tables

Pages

- Table 1: Distribution of the Basin in the Six Riparian Countries.*
- Table 2: Average annual rainfall and evapotranspiration in the riparian countries of the Volta Basin*
- Table 3: Rainfall and length of growing seasons in the Volta Basin of Ghana*
- Table 4: Hydrometeorology Mean Annual (1961 – 1990) in the Volta Basin of Ghana*
- Table 5: Temperature and Relative Humidity in the Volta Basin of Ghana*
- Table 6: Demographic characteristics of the Volta Basin countries (2001)*
- Table 7: Population in the Volta River basin*
- Table 8: Population statistics in riparian countries (2004)*
- Table 9: Education and literacy in the Volta Basin countries*
- Table 10: Agricultural labour force in the riparian countries of the Volta River Basin*
- Table 11: Structure of Economic Output at the National Level*
- Table 12: Macroeconomic indicators in the Volta Basin countries (2001)*
- Table 13: Human development indicators in the Volta basin countries (1998)*
- Table 14: Poverty and income distribution in the Volta Basin countries (latest year available)*
- Table 15: External Debt of the Riparian Countries (1998)*
- Table 16: Area and length of main tributaries in the Volta River Basin in Ghana*
- Table 17: Some Important Relief Characteristics*
- Table 18: Identified Soil Groups in the Volta River Basin of Ghana*
- Table 19: Characteristics of the Zones*
- Table 20: Major landuse types in the Volta basin of Burkina (hectares)*
- Table 21: Erosion Hazards of the Volta Basin in Ghana*
- Table 22: Rate of Occupation of Cultivable Land in the Basin in Togo (1,708,800 ha)*
- Table 23: Evolution of Various Vegetation Formations in Togo (1979-1991)*
- Table 24: Production Levels of Selected Crops by Regions in the Volta Basin in Ghana (Tones)*
- Table 25: Cereal Production in the Mopti Region of Mali (Tones)*
- Table 26 : Crop Production in Côte d'Ivoire (1996)*
- Table 27: Production trends in Burkina Faso (x1000 tons)*
- Table 28: Population of Major Livestock in the Volta Basin in Ghana*
- Table 29: Population of Major Livestock in Togo*
- Table 30: Annual Fish Production in Ghana*
- Table 31: Surface Water Resources of the Volta River in Ghana*
- Table 32: Surface water flows of the Black Volta in Ghana*
- Table 33: Surface Water flows of the Lower Volta of Ghana*
- Table 34: Sub-basins of the Main Volta River in Ghana*
- Table 35: Surface water resources from within the Lower Volta Basin in Ghana*
- Table 36: Potential Surface Water Resources of the Volta Basin in Burkina Faso*
- Table 37: Potential of Surface Water Resources of the Nakambe Basin*
- Table 38: Potential of surface Water Resources of the Mouhoun Basin in Burkina Faso*
- Table 39: Hydrological Characteristics of the Groundwater aquifers in the Volta Basin of Ghana*
- Table 40: Summary of borehole yields of hydrologic provinces and sub provinces*
- Table 41: Minimum Recharge and Replenishable Groundwater Capacities*

Table 42: Chemical analysis of water samples in the geologic formations of Ghana (all values except pH are in mg/l)

Table 43: Groundwater potentiality of Burkina Faso

Table 44: Average Rate of well-sinking success and discharge by region

Table 45: Proportion of Volta Basin in Agro-ecological zones of Ghana

Table 46: Access to land

Table 47: Proportion of area allocated to major crops species and yield per hectare in the GS

Table 48: Cropping patterns in the Guinea Savannah zone

Table 49: Percentage of use of each technology

Table 50: Percentage of farmers practicing cropping system in Wenchi

Table 51: Percentage of farmers keeping livestock, 1992

Table 52: Estimated Average National wholesale prices of crops/tonne (1998)

Table 53: Sequence of planting in the Techiman District of Ghana

Table 54: Types of farming systems in the Techiman District of Ghana

Table 55: Estimated average national wholesale prices of crops/tonne (1998)

Table 56: Types of farming systems in the Atwima District of Ghana

Table 57: Estimated Average National wholesale prices of crops/tonne (1998)

Table 58: Ministries and Departments for managing water and Land resources in Benin

Table 59: Departments for Managing Land and Water Resources in Burkina Faso

Table 60: Ministries and Their Responsibilities in Côte d'Ivoire

Table 61: Ministries for the Management and Use of Land Resources in Côte d'Ivoire

Table 62: Ministries and Departments Responsible for Water Resources Development and Utilization in Ghana

Table 63: Ministries, Departments, and Institutions Responsible for the Management of Water in Togo

Table 64: Bilateral Cooperation among Riparian Countries

Table 65: Dates of Ratification of Major International Environmental Conventions

Table 66: Water demand during year 2000 (km³)

Table 67: Projected water demand in Burkina Faso (km³)

Table 68: Large dams in the Volta Basin of Burkina Faso

Table 69: Information on Dams in the Volta Basin of Cote d'Ivoire

Table 70: Information on Dams in the Volta Basin of Togo

Table 71: Domestic/Industrial Water Demand of the Volta River Basin ($x 10^6 m^3$)

Table 72: Irrigation Water Demand of the Volta River Basin ($x 10^6 m^3$)

Table 73: Water Demand for Livestock of the Volta River Basin ($x 10^6 m^3$)

Table 74: Total Consumptive Water Demand of the Volta River Basin ($x 10^6 m^3$)

Table 75: Dependency by region on groundwater supply for domestic use (1984)

Table 76: Categories of infected people according to Farooq, 1963

Table 77: Annual population growth rate by sub-basins, 1960-200

Figure 1 : NOAA-AVHRR satellite image composition LANDSAT 1999-2000 image composition

Figure 2a: Declining patterns in rainfall in Burkina Faso.

Figure 2b: Changes in Climatic zones of the Volta Basin of Burkina

Figure 2c: Annual rainfall at several locations in the Volta Basin of Burkina Faso

Figure 3: Distribution of rainfall in the Volta Basin in 1990

Figure 4: Evaporation in several locations within the Volta Basin in Burkina Faso

Figure 5: Spatial distribution of potential evapotranspiration in the Volta Basin

Figure 6: Spatial distribution of temperature in the Volta Basin

Figure 7: Spatial distribution of population density in the Volta River Basin

Figure 8: Distribution of population density in Ghana

Figure 9: Pyramidal ages in Ghana

Figure 10: Distribution of soil types in the Volta basin of Ghana and Burkina

Figure 11: Map of Geology of the Volta River Basin.

Figure 12: Land cover Map of the Volta River Basin.

Figure 13: Hydrographic network of Burkina Faso

Figure 14: Location of reservoirs in the Volta Basin

Figure 15a: Water balance of the Mohoun River Basin

Figure 15b: Water balance of the Nankambe River Basin

Figure 16: Hydrogeological provinces and river systems of Ghana

Figure 17: Hydrogeological subprovinces of the Basement Complex

Figure 18: Hydrogeological subprovinces of the Voltaian System

Figure 19: Distribution of borehole yield in Ghana (Water Resources Research Institute 1994)

Figure 20: Regional distribution of borehole and annual abstraction

Figure 21: Map showing discharge of wells across Burkina Faso

Figure 22: Distribution of Piezometric heads of different wells in Burkina Faso

Figure 23: Spatial distribution of boreholes in Burkina Faso

Figure 24: Map showing the agro-ecological zones of Ghana and Burkina Faso

Figure 26: Location of Dams in the Volta Basin

Figure 27: Regional distribution of groundwater use among sectors

Figure 28: Water Weed Encroachment

Figure 29: Sandbar on the Volta River mouth

Figure 30: Annual percent population growth rates by sub-basins, 1960-2000

Figure 31: Average cropped area by sub-basins, 1992 & 2000

I. INTRODUCTION

I.1 The Volta River Basin

The Volta River Basin is located in West Africa and covers an estimated area of 400,000 km². The Volta basin stretches from approximately latitude 5° 30' N in Ghana to 14° 30' N in Mali. The widest stretch is from approximately longitude 5° 30' W to 2° 00' E but the basin becomes more narrow towards the coast of the Gulf of Guinea. The Volta basin is spread over six West African countries (43% in Burkina Faso, 42% in Ghana, and 15% in Togo, Benin, Cote d'Ivoire and Mali).

The Akosombo dam is by far the most significant structure built in the basin and the *Volta Lake is the largest man-made lake in the World*. The Volta Lake reservoir itself has a surface area of about 8,500 km², an average depth of about 18.8m and a shoreline of about 5,500 km. The deepest portions of the lake are about 90m. The total volume of water in the reservoir at full supply level (about 84.73m³) is approximately 150 billion m³. The seasonal rise and fall is about 2.0-6.0m and the areas covered by seasonal fluctuations are about 100,000ha. The Kpong Head pond, i.e. the area between the Akosombo dam and the Kpong Dam, has a surface area of 12 km² with a total volume of about 190 km³. The average head pond elevation is about 14.7m. The lower Volta area, i.e. the area between the Kpong Dam at Akuse and the estuary at Ada, is about 68,600km². Water available from the reservoirs is primarily used for hydro-energy production, other significant uses being transportation, fishery, water supply (commercial and domestic purposes), tourism and irrigation

The Akosombo dam is of strategic importance to the economy of Ghana. It generates 80% of the power produced in the country. The primary purpose of the project was to supply cheap electricity to smelt aluminum and the secondary one is the development of the country. Contrary to many hydropower dams that are generally built upstream, the Akosombo dam is built closed to the ocean because of the flat relief of the basin and the difficulty to find potential locations. Construction of the Volta Lake led to the resettlement of about 80,000 people from several hundred of villages to fifty newly built townships (more than 1% of Ghana's population at that time). In addition to the resettlement of the river communities, damming affected local health, agriculture, fishing, and navigation.

Population in the basin lives generally away from the main water courses because of water borne diseases such as Onchocerciasis and Schistosomiasis which are prevalent in the region. Public health related problems were those of a switch between two disease episodes – i.e. the one that thrives well in riverine system and the other which thrives well in (benthic) riverine systems. It was a switch between onchocerciasis (river blindness) and urinary schistosomiasis. Onchocerciasis is transmitted by the black-fly (*Simulium damnosum*) which breeds solely in fast flowing waters. Urinary schistosomiasis, commonly called bilharzias is a chronic snail-borne infection, which frequently occurs in water development schemes in tropical countries. The disease was endemic in Ghana long before the creation of the Volta Lake; but endemicity was low along the Volta River. Prevalence in school children was 5% according to an epidemiological survey made in 1960-61 before the lake was formed.

The creation of the lake and the consequent biological explosion of aquatic weeds associated with the

aquatic snail, the “intermediate host,” together with mass migration into the fishing communities above

The head pond area from regions in which the disease was endemic led to a great increase in the prevalence of the disease in many localities around the lake.

In the other riparian countries of the basin, small and large dams have been built by governments, NGOs and local people to secure food production after the severe droughts that occurred in the 1970s and 1980s. In the Nakambe sub-basin (Burkina Faso) alone more than 600 small dams have been built most of them during that period. More recently power generating dams have also been built in some of the Volta main tributaries Bagre and Kompienga (Burkina) with generating capacities of 41.5 GWH and 31.0 GWH respectively and on the Oti River, at the border between Togo and Benin within a power generating capacity of 35GWH.




Figure 1 : NOAA-AVHRR satellite image composition ANSAT 1999-2000 images composition

The distribution of the area of the basin among the six riparian countries is shown in the following Table 1.

Country	Area of Volta River Basin (km ²)	% of Basin	% of Country in the basin
Benin	17,098	4.10	15.2
Burkina Faso	178,000	42.65	63.0
Côte d'Ivoire	12,500	2.99	3.9

Mali	15,392	3.69	1.2
Togo	26,700	6.40	47.3
Ghana	167,692	40.18	70.0
Total	417,382*	100%	

Figures for area of Volta River Basin are from respective National Reports. The areas recorded from the country reports are quite similar to that quoted from Moniod, et al (1977).

As indicated in Table 1, the relative proportion of the basin area found within a country does not necessarily reflect the relative importance of that part of the basin in that country. While a country may have only a small percentage of the total basin within its borders, as in the case of Togo, this area might comprise a significant proportion of the entire country. Additionally, the area of the country within the basin might hold abundant natural resources with respect to the entire country, such as in the case of Mali, Burkina Faso, Ghana and Togo.

I.2 Methodology and presentation of the study

This report is a compilation of various documents collected in Burkina Faso and Ghana over the past year. Institutions and many people involved in research and development projects in the Volta Basin in both countries have also been visited during the same period. Some of the information especially socio-economic data presented in this report are country specific. However, because the basin covers 63% of Burkina, and almost 50% of Ghana, it can be estimated that such information are also valid for the basin. A GIS database of the Volta Basin has been created and is now available along with a collection of reports.

II. THE VOLTA RIVER BASIN: HUMAN AND PHYSICAL SETTING

II.1 Climatic features

The climate of the region is controlled by two air masses: the North-East Trade Winds and the South-West Trade Winds.

The North-East Trade winds, or the harmattan, blowing from the interior of the continent, are dry. In contrast, the South-West Trade winds, or the monsoons, are moist since they blow over the seas. The inter-phase of these two air masses is called the Inter-tropical Convergence Zone (ITCZ). There is a lot of convective activity in the region of the ITCZ; hence the region is associated with a considerable amount of rainfall. The ITCZ moves northwards and southwards across the basin from about March to October when rainfall is received in the region.

II.1.1 Distribution of Rainfall

Three types of climatic zones can be identified in the region: the humid south with two distinct rainy seasons; the tropical transition zone with two seasons of rainfall very close to each other; and, the tropical climate, north of lat 9° N, with one rainfall season that peaks in August. Average annual rainfall varies across the basin from approximately 1600 mm in the southeastern section of the basin in Ghana, to about 360 mm in the northern part of Burkina Faso. Figure 3 shows the spatial distribution of rainfall in the Volta Basin and Table 4 shows the average annual rainfall and evapotranspiration in the riparian countries.

There have been a number of changes in the precipitation patterns in some sub-catchments in the basin, as rainfall and run-off reductions have been evident since the 1970s (Opoku-Ankomah, 2000). Some areas that used to have bi-modal type of rainfall have only one mode as the second minor season has become very weak or non-existent. This situation means that rainfed agriculture can only be carried out once instead of twice a year.

In Burkina Faso, the Volta basin stretches into three climatic zones:

1. the Sudan zone with an annual rainfall between 900 and 1,200mm distributed on average over 74 rainy days. It is located below the 11° 30'N parallel
2. the Sudano-Sahelian zone with an annual rainfall between 600 and 900mm on average over 43 rainy days, located between the 11° 30'N and 14°N parallels
3. the Sahelian zone located above the 14°N parallel with a mean annual rainfall between 300 and 600mm over 38 rainy days.

In the Sahelian zone, the rainy season lasts for about 3 months. It lasts 4 to 5 months in the Sudano-Sahelian zone and 6 to 7 months in the southern part of the Sudan zone. Dominant winds blow east-west from January to March which is the harmattan season. Over the past 40 years, the rainfall amount has been decreasing leading to severe droughts during the 1970s and 1980s and only a slight recovery in 1985-1995. As a consequence of this instability in the rainfall pattern most of the rivers have dried up, most of the land cover have been degraded and the water table has been drawing down.

This has led to a shift of the climatic zones in a southerly direction. Consequently most of the Volta basin in Burkina Faso is now located in the Sahelian and Sudano-Sahelian zones. The following figures show the declining pattern in rainfall in Burkina Faso, the shift of the climatic zones, as well as

the rainfall variability and intensity over the past 40 years in several locations within the Volta basin in Burkina.




Figure 2a: Mean Annual rainfall in Burkina Faso.





Figure 2b: Changes in Climatic zones of the Volta Basin of Burkina Faso









Figure 2c: Annual rainfalls at several location in the Volta Basin of Burkina Faso




Figure 3: Distribution of rainfall in the Volta Basin in 1990

Table 2: Average annual rainfall and evapotranspiration in the riparian countries of the Volta Basin

Riparian Country	River	Area of the Volta Basin km ²	Upstream Riparian Countries	Average Annual Rainfall (mm)	Average Annual Evaporation (mm)
Ghana	Volta	165,830	Burkina Faso, Mali, Togo, Cote d'Ivoire, Benin	1320	1415
Cote D'Ivoire	Volta	9,890	Burkin Faso, Mali	1358	1486
Togo	Volta	25,545	Burkina Faso, Benin	1305	1697
Burkina Faso	Volta	171,105	Mali	950	2130
Benin	Volta	13,590	Burkina Faso, Togo	1294	1400
Mali	Volta	12,430	None	685	3015

It has been estimated that 340 km³ of rain must fall on the catchment before run-off occurs at significant levels. Once this threshold has been reached, approximately half of the precipitation becomes run-off. This indicates that only small changes in rainfall could have dramatic effects on run-off rates. Although rainfall decreased by only 5% from 1936 to 1998, run-off decreased by 14% (Andreini, 2000).

Simulations of run-off using GCM-based climate scenarios developed by Minia (1998) showed 15.8% and 37% reduction in run-off of the White Volta Basin for years 2020 and 2050, respectively (Opoku-Ankomah, 2000). These projections showed that projects whose design was based on historical records without considering climate change, such as the hydropower dam at Akosombo, could be vulnerable

Table 3: Rainfall and length of growing seasons in the Volta Basin of Ghana

Agro-ecological Zone	Mean annual Rain (mm)	Growing Period (Days)	
		Major season	Minor season
Rain Forest	2,200	150 - 160	100
Deciduous Forest	1,500	150 - 160	90
Transitional	1,300	200 - 220	60
Coastal	800	100 - 110	50
Guinea Savanna	1,100	180 - 200	-
Sudan Savanna	1,000	150 - 160	-

II.1.2 Distribution of Evaporation and Evapotranspiration (ETP) compared to rainfall

Evaporation in the basin is relatively high especially in the Sahelian zone. In Burkina (43% of the basin), the lowest record is about 1,900mm/yr.

In the Volta Basin of Ghana, the Potential evapotranspiration varies from a minimum of 1,450 mm per annum in the Black Volta sub-basin to a maximum of 1,968 mm per annum in the White Volta sub-

basin. Table 5 shows the range of potential evapotranspiration as well as other hydrometeorological parameters in each sub-basin of the Volta Basin in Ghana. Pan evaporation is the same (2540 mm/an) in all the sub-basins except the Lower Basin.

Table 4: Hydrometeorology Mean Annual (1961 – 1990) in the Volta Basin of Ghana

Volta Basin System	Rainfall (mm)	Coeff. Of Var.	Pan Evaporation (mm)	Potential Evapotranspiration (mm)
Black	1023.3 – 1348.0	0.17 – 0.23	2540	1450.0 – 1800.0
White	929.7 – 1054.2	0.16 – 0.20	2540	1650.0 – 1968.0
Oti	1050.0 – 1500.0	0.18 – 0.20	2540	1550.0 – 1850.0
Lower	876.3 – 1565.0	0.17 – 0.35	1778	1450.0 – 1800.0

In Burkina Faso, average annual pan evaporation values are generally very high and at the same time progressively increase from south to north. The minimum annual value in the Volta basin is about 1900mm. Average levels recorded in the basin are 2334 mm in the north (Ouahigouya), 2120 mm in the centre (Ouagadougou) and 1932 mm in the south (Gaoua) (Monido et al.,1977). Figure 4 shows average pan evaporation values at several locations within the Volta River basin.




Figure 4 : Evaporation in several locations within the Volta Basin in Burkina Faso

Figure 4 shows the distribution of potential evapotranspiration in Ghana and Burkina Faso (about 85 % of the Volta Basin). Generally, ETo decreases downstream of the basin.




Figure 5: Spatial distribution of potential evapotranspiration in the Volta Basin

II.1.3 Temperature and humidity

The annual mean temperatures in the Basin vary from about 27° C to 30° C. Daily temperatures can be as high as 32° C - 44° C, whereas night temperatures can be as low as 15° C. The humidity varies between 6% and 83% depending on the season and the location. Figure 5 shows the spatial distribution of temperatures in the Volta Basin. Generally, temperatures are higher in the upstream of the basin and decreases in the downstream

Generally, in Ghana (42 % of the Basin) the mean temperature never falls below 24°C in the country. This is explained by the fact that no part of the country is really far from the equator. The hottest month of the year is March-April and the coolest is August. The southern section of the country is more humid than the north. In the coastal area of Ghana the relative humidity are 95-100% in the morning and about 75% in the afternoon. In the north values can be as low as 20-30% during the harmatan period and 70-80% during the rainy period.

In Burkina Faso, the mean temperature in the Sahel zone is always higher than 29°C while in the Sudano-Sahelian zone it lies between 28°C and 29 °C and in the Sudan zone below 28°C. The seasonal variation in temperatures is characterized by four periods: two extremely hot periods and two relatively cool periods. The first hot period is in March-April with average maximum temperatures of 37°C and 41°C in the south, centre and north of the basin respectively, while the average minimum temperatures are 24°C (south), 25°C (centre) and 26°C (north). The second hot period occurs immediately after the rainy season. It is not as hot as the first, with average maximum temperature of 34°C, 36°C and 38°C in the south, centre and north respectively. Minimum temperatures vary between 21°C and 22°C.

The first cool periods occur in December-February, with average maximum temperatures varying between 33°C and 35°C in January and average minimum temperatures in the order of 14°C (north), 17°C (centre) and 19°C (south). This is the period of harmattan, a dry dusty wind that makes it cold at night and hot during the day. The second cool period coincides with the rainy season. During this period, the moisture content of the air is at the highest. The average maximum and minimum temperatures during this cool period are subject to extreme variations due to the irregularity of the rains. These extreme regional and temporal variations, combined with the shortness of the rainy season, are one of the ecological factors that play a key role in limiting crop production and making a major contribution to the degradation of the vegetation cover.

Relative humidity can reach a maximum of 80% in August which is the wettest month of the rainy season. There are sometimes heavy falls of dew, especially in the south where levels of between 0.8mm and 1mm have been recorded. Total evaporation in August is generally lower than 100mm. From November to April relative humidity in Burkina is about 50%. Maximum pan evaporation (>400mm) is observed during March-April.




Figure 6: Spatial distribution of temperature in the Volta Basin

Conditions of temperature and humidity in the Volta Basin in Ghana do not differ too much from what prevails generally over the portions of the country within the basin. Table 5 below gives figures of

temperature and relative humidity of the different sub-basins of the Volta Basin in Ghana.


Table 5: Temperature and Relative Humidity in the Volta Basin of Ghana

Volta System	Basin	Temperature (°C)	Relative Humidity (%)
Black		25.0 – 27.8	59 – 77
White		28.0 – 28.6	54 – 68
Oti		27.0 – 27.8	62 – 72
Lower		23.5 – 28.1	6 – 83

II.2 Human Context

A. Human resources

The geographic distribution of the population within the basin is highly variable with a density ranging from 8 to 104 persons/km². In areas with high density population one can notice a real pressure on land and water resources (Ghana's Upper East regions: 104 persons/km²). Figures 7, 8, and 9 show the spatial distribution of population density in the Volta River Basin, Ghana and Burkina Faso respectively. In general, areas with low population density are either national park (Comoe one of the largest West African national parks, in Cote d'Ivoire) or regions where *onchocerciasis* or *river blindness* is prevalent (valleys of the Black Volta). Three cities in the basin (Ouagadougou, Bobo Dioulasso and Tamale) can be described as large, each having a population of over one hundred thousand (100,000) people.



Basin Area (sq. km.):	407,093
Average Population Density (people per sq. km.):	43
Number of Large Cities (>100,000 people):	3
Water Supply per Person (1995) (m ³ /person/year):	2,054

Figure 7: Spatial distribution of population density in the Volta River Basin




Figure 8: Distribution of population density in Ghana

In Ghana the most populated areas are located outside the Volta Basin, except Tamale and the upper regions (Fig 8). The two largest cities of Burkina (Ouagadougou and Bobo-Dioulasso) are located within the basin. The average population density in the Volta Basin of Burkina Faso is 29 inhabitants/km² with strong regional variations. While the high population density (79 inhabitants/km²) on the Central Plateau has led to the over exploitation of natural resources and a marked degradation of the environment, the sparsely populated eastern and western regions (10 inhabitants/km²) are attracting a number of immigrants. This is causing problems in terms of rational land management and the degradation of natural resources.

Table 6: Demographic characteristics of the Volta Basin countries (2001)

Setting	Total Population (million)	Estimated Population 2005 (million)	Population growth Rate (annual %)	Population density (people per sq km)	Population density, Rural 2000 (people per sq km)	Population (% Rural)	Rural population growth rate (annual %)	Population ages 0-14 (% of total)	Population ages 15-64 (% of total)	Population ages 65 and above
Benin	6.44	7.12	2.56	58.19	185.70	57.00	1.32	45.85	51.36	2.71
Burkina Faso	11.55	12.66	2.38	42.22	247.68	83.13	2.04	47.08	50.14	2.66
Côte d'Ivoire	16.41	17.65	2.20	51.60	306.08	55.96	1.70	41.90	55.05	2.54
Ghana	19.71	20.91	1.83	86.61	341.99	63.60	1.55	43.06	51.87	4.53
Mali	11.09	12.05	2.25	9.09	163.43	69.14	1.37	46.97	49.81	3.00
Togo	4.65	5.02	2.40	85.56	120.19	66.08	1.94	43.57	52.58	3.15

Data source: WDI (2003); and “..” denotes data not available.

Burkina Faso and Mali have the highest percentage of young population, 0-14 years (47%), while Ghana has the lowest (43%). A major fraction of population falls in working ages 15-64 years, for all countries. The highest percentage of the working age groups occurs for Côte d'Ivoire (55%) and for the rest of the riparian countries, it varies between 50% and 53%. Figure 9 shows a population pyramid of Ghana after the 1999 Population Census




Figure 9: Pyramidal ages in Ghana

Population in the Volta Basin is generally rural (64-88 %, Table 7) and the people depend to a large extent, on the exploitation of the natural resources for their livelihood, which may not be environmentally sustainable in the future. In Burkina Faso which represents 43 % of the total area of the basin, about 78 % of the total population live in the Volta Basin which has a vital socio-economic importance for the country. Out of the 45 provinces in the country, 37 are located within the

Volta Basin within two sub-basins Mouhoun and Nakambe with population density of respectively 41 and 53 persons/ km². In Togo, the Volta Basin covers 48 % of the country's land and is home to 35 % of the total population in 2000.

Table 7: Population in the Volta River basin

Country	1990	2000	2010	2020	2025	Growth Rate (%) (2000)	P/km ² Density (2000)	Urban %	Rural %
Benin	382,328	476,775	596,000	746,000	820,000	2.27	43.4	36	64
Burkina Faso	7,014,156	8,874,148	11,227,366	14,204,605	15,997,351	2.38	41.53	22.6	77.4
Côte d'Ivoire	-	397,853	497,469	632,313	717,672	2.53	8 - 22	23	77
Ghana	5,198,000	6,674,376	8,570,068	11,004,185	11,696,054	2.5	26 - 104	16	84
Mali	380,000	625,000	880,000	1,140,000	1,260,000	2.78	45 - 75	12.2	87.8
Togo	1,189,900	1,594,446	2,153,719	2,891,457	3,385,266	2.80	66	30	70
Total	14,474,276	18,642,598	23,924,622	30,618,560	33,876,343				
Average						2.54	48.49	23.30	76.70

The total basin population is expected to grow significantly from an estimated 18,600,000 hbts in 2000 to approximately 33,900,000 hbts in 2025. This is an expected increase of 80 % within a twenty-five year period. This high growth is due to the high average population growth rate (2.54 %) in the basin.

About half of the population in each of the riparian countries is in the age group of (15-64) years (Table 8), which forms the majority of the labor force. This age pattern might also be true for the areas within the Volta river basin, meaning that the basin is significantly rich in human resources. Other important population statistics in the riparian countries of the basin are presented in Table 8.

Table 8: Population statistics in riparian countries (2004)

Country	Life Expectancy (Yrs)	Death Rate (per 1000)	Literacy Rate (%)	Sex Ratio (male(s)/female)	% Age Structure (Yrs)		
					0-14	15-64	64 and over
Benin	51	13.69	40.90	0.98	46.8	51.0	2.3
Burkina Faso	42	18.79	26.60	0.97	46.0	51.1	2.9
Côte d'Ivoire	42	18.48	50.90	1.01	45.1	52.6	2.2
Ghana	56	10.67	75.00	1.00	38.0	58.3	3.7
Mali	45	19.12	46.40	0.96	47.1	49.9	3.0
Togo	53	11.64	60.90	0.97	43.9	53.6	2.6
Average	48	15.40	50.12	0.98	44.48	52.8	2.8

Source: CIA, 2004

Population pressures in countries with a weak economic base, as in the basin, induce unsustainable use of forest and land resources. This constitutes a major factor in the degradation of natural resources especially in Burkina Faso. Between 1985 and 1995, the population of Burkina Faso increased by 2.7%. Since then, the increase has been in the order of 2.6%. In the near future the population is expected to reach almost 16 million. This rapidly expanding population automatically requires a great area of cultivated land, which in turn means more land clearance. This inevitably contributes to the destruction of small amount of available vegetation cover. According to the Ministry of Agriculture, the high population density on the central plateau which lies entirely within the Volta basin and the high levels of intensive farming (between 50% and 70%) mean that the agro-demographic threshold has been reached, if not exceeded, and has led to a process of land degradation

The easy movement of people across national boundaries in the sub-region under the ECOWAS protocol makes population pressure a transboundary cause of the above-mentioned environmental problems in the basin. There have been some population migrations in the basin. In Ghana, the decline in the fishing industry in the Lower Volta following the establishment of the Volta Lake upstream has attracted people to move upstream to live near the lake for their livelihood. It is unfortunate however, that these recent settlements are often close to the banks of the lake.

In Togo, some people in the basin (Savannas and Kara regions) who migrated to the southern regions of the country before 1990 are now returning due to socio-political unrest.

The population of Burkina Faso experiences an important migration at both internal and external levels. The average ratio between internal and external migrations is as follows: out of 100 migrants people, 71.2% migrate inside the country and 28.8% abroad. In Burkina Faso during the severe Sahelian droughts of the 70s and 80s numerous populations have moved from the northern and central regions (Plateau Mossi) of the country to the south in search of a better life. Many Burkinabes are currently migrating from one region to another, or from rural to urban areas. There are three types of migration:

- The exodus towards urban centres
The migration of young people towards urban areas in search of paid work. This uncontrolled migration led to the over-exploitation of land and deforestation around the large towns, and creates problems in terms of urban development. Between 1975 and 1985, the urban growth of Ouagadougou and Bobo-Dioulasso was 156% and 101% respectively
- Migration to other countries
This type of migration is characteristic of rural areas, usually in regions with poor financial resources. The migrants move mainly towards Cote d'Ivoire and, to a lesser extent to Ghana. The departure of these members of the working population depopulates the villages and deprives them of the workforce required to implement Soil and Water Conservation and other labour intensive initiatives.
- Rural migration
Migration within the country involves the individual or a group and it is spontaneous or organized migration from regions that are usually infertile and/or over-populated to regions that are under-populated and under-exploited, with the intention of settling temporarily (during the rainy season) or permanently.

Rural migrations within Burkina Faso have increased since the drought of 1970-72. They are regarded as a purely temporally solution to the problem of over-population and the absence of cultivated land. The problems experienced in the central region will soon be repeated in the newly colonized region since the settlement of the migrant population and the cultivation of these areas happen in a haphazard way that results in the exploitation of the land and vegetation cover. Rural migrations occur as follows:

- individual or collective departures occurring from usually infertile and overpopulated regions (e. g. : the central plateau or Mossi land) towards more fertile and hardly exploited areas; the main host provinces are those of Houet, Mouhoun, Gourma, Tapoa, Kossi and Sissili;
- spontaneous departures towards places where there are real possibilities of earning more substantial incomes (e. g. : gold sites), noticed in the provinces of Sanmatenga, Passoré, Séno and Soum;
- State organized migrations towards areas developed for agricultural exploitation; these concern the developed plain of the Sourou province and the plain of the Kompienga and Bagré dam and the Kou valley.

The settlement of populations in these areas is anarchistic. However, as far as organised migrations are concerned, the National Office in charge of Land Development (ONAT) is trying to organize the settlement of migrants.

The growth rates of migrations in host provinces between 1975 and 1985 were as follows : 88% for the Houet province, 44% for Mouhoun, 73% for the Tapoa, 64% for the Kossi and 106% for the Sissili (Kessler et Geerling, 1994).

Mali has also seen migration into the “forest” of Samori, a sub-basin of the Volta. This movement is caused by the quest for new land for farming activities. Others moved into the basin after the drought of 1985. Another sub-basin in Mali, the Seno, has seen such a surge in population that there is no longer sufficient farmland to allow land to lie fallow, resulting in the impoverishment of the land. Additionally, there has been some migration out of the basin and into the urban areas where jobs are sought.

As noted in **Table 8**, the literacy rate is in the range of 26 to 75% with a mean of approximately 50% for the entire basin. This average level can impede environmentally sustainable development. Also, there are significant disparities in the schooling and literacy rates for men and women. In Togo, for example, between 43 and 83% of women are illiterate, while the illiteracy rates for men is between 25 and 50%.

Table 9: Education and literacy in the Volta Basin countries

Country	Education efficiency coefficient (%)*	Net intake rate in grade 1 (%) 2000		Adult illiteracy rate (%) ≥15 yrs, 2001		Youth illiteracy rate (%) 15-24 yrs, 2001	
		Male	Female	Male	Female	Male	Female
Benin	49.8*	46.54	75.44	28.37	62.75
Burkina Faso	67.7	24.57	17.22	65.09	85.08	53.14	75.46
Côte d'Ivoire	59.0	29.84	24.08	39.66	61.57	28.73	46.42
Ghana	87.5	29.29	29.34	18.94	35.46	6.12	10.62
Mali	66.4	63.33	83.35	51.81	74.00

Togo	44.9	50.11	43.63	26.62	56.03	12.24	34.79
------	------	-------	-------	-------	-------	-------	-------

Data source: WDI (2001; 2003); *=primary school only;

Life expectancy in the basin is fairly low, varying between 42 and 56 years with an average of 48 (Table 8). The infant mortality rate in the basin is high, estimated in 2004 to be between 49.25 and 55.1 per 1000 births in Ghana and between 90.39 and 106.7 in Burkina (CIA, 2004). A major factor in the short life expectancy of the population in the Volta basin is inadequate access to health care. Access to public health care in the Volta Basin in Ghana is poor and significantly below the national average of 37.2% .

In Burkina Faso, only 28% of children between 12 and 23 months old are fully vaccinated against the six main diseases affecting children. About 29% of children between 0 and 5 years old are malnourished. In Togo, within the Volta basin about 42% of children under age three are concerned. Although health care is being continually improved, it is marked by a high level of infant mortality. Only 31.7% of children are in full-time education and only 12.3% of the population are literate. Although health care is being continually improved, it is marked by a high level of infant mortality.

Agricultural labor force

Ghana has the largest human labor force of 9.5 million, followed by Côte d'Ivoire with 6.6 million, Burkina Faso and Mali have 5.7 and 5.4 million respectively. Benin and Togo have the lowest labor force of 2.9 million and 1.9 million respectively. During 1980-1999, labor force growth rates have varied substantially in Côte d'Ivoire as it was 3.4%, in Ghana 2.9% and Togo 2.7%. It has remained around 2% in Mali and Burkina Faso. Benin had during the same period the lowest labor force growth rate of all the riparian countries, 0.4%. Agriculture accounts for 86% of total employment in Mali and 92% in Burkina Faso; for all other countries it accounts for not less than 60%.

Employment rates in the agriculture sector of the riparian countries vary according to gender. In the agriculture sector of Côte d'Ivoire, the employment rates accounts for 54% of total male employment, and 72% of total female employment. Employment patterns are more or less the same in almost all the other riparian countries except Ghana where, the agriculture sector accounts for a higher proportion of female employment than male employment. Women in the Volta Basin are heavily depend on the agriculture sector for securing their livelihood.

Table 10: Agricultural labour force in the riparian countries of the Volta River Basin

	Total population (million) 2001	Labor force		Employees in agriculture (latest year available)**		Employment in agriculture (% of total employment)**
		Total 2001 (millions)	Average annual growth rate (%) 1980-99*	Male (% of male employment)	Female (% of female employment)	
Benin	6.44	2.91	0.4	61.90	65.20	63.50
Burkina Faso	11.55	5.68	1.9	91.40	93.50	92.40
Côte d'Ivoire	16.41	6.60	3.4	54.20	72.00	60.00
Ghana	19.71	9.40	2.9	62.20	58.70	62.20
Mali	11.09	5.42	2.2	83.00	89.00	85.80
Togo	4.65	1.93	2.7	66.20	64.50	65.50

Data source: WDR (2000); WDI (2003); * (WDI, 2001); **=latest year available (1990s).

II.3 Economic Sector and Policies

The riparian countries of the Volta River Basin are some of the poorest in the world having underdeveloped economies. Mali has the highest proportion of the population below the poverty line (64 %-2001 est.), 44.5% in Burkina Faso in 1997 and Ghana recorded the least in 1992 (34.1 %). See Table 9. According to the World Development Report 2000/2001, all of the Volta River Basin countries are considered to be in the low income category (GNP per capita of \$755 or less). Per capita GNP and economic growth rates for the riparian countries are shown in the following tables.

Côte d'Ivoire has the highest GNP in the region with \$710 per capita, while Mali is ranked lowest with only \$190. The average GNP/capita is \$372, making the region one of the poorest in the world. Although figures quoted in the table are national values, the condition in the basin is not better than in the remainder of the countries. The average annual growth rates in the range of -0.3 to 2.7% of GNP/capita also shows low performance of the economies of the region. During 2001, Benin, Burkina Faso Ghana registered a robust GDP growth rate of 4 to 5 percent, while Côte d'Ivoire experienced a negative growth rate. Likewise, Côte d'Ivoire, Mali, and Togo showed a negative per capita GDP growth rate, while all others have a positive figure. Over all, per capita GDP growth rates are lower in all countries than GDP growth rates, which are picking up the effect of fast growing population in all countries, the difference being widest in the case of Togo with the fastest growing population.

Economic activities in the basin are quite similar in all of the countries such as crop production, livestock breeding, fishing, lumber, agro-industry, transportation, and tourism. These activities can be grouped under agriculture, industry, manufacturing, and services.

Agriculture includes crop and livestock production, fisheries, and forestry, while industry involves mining and quarrying, electricity supply, and construction; services include transport, storage, communication, wholesale and retail trade, restaurants and hotels, government services, etc. The economic outputs of these activities in the riparian countries are shown in Table 11.

Table 11: Structure of Economic Output at the National Level

Country	Gross Domestic Product		Value added as % of GDP							
	Millions of \$		Agriculture		Industry		Manufacturing		Services	
	1990	1999	1990	1999	1990	1999	1990	1999	1990	1999
Benin	1,845	2,402	36	38	13	14	8	8	51	48
Burkina Faso	2,765	2,643	32	32	22	27	16	21	45	41
Côte d'Ivoire	10,796	11,223	32	24	23	24	21	20	44	52
Ghana	5,886	7,606	45	36	17	25	10	9	38	39
Mali	2,421	2,714	46	47	16	17	9	4	39	37
Togo	1,628	1,506	34	43	23	21	10	9	44	36
	-	-	37.8	36.7	19.0	21.3	12.3	11.8	43.5	42.2

(World Development Report 2000/2001)

Table 11 shows economic outputs for the countries as a percentage of GDP in 1990 and 1999. The activity output is shown as a percentage of the GDP. From the table, it can be observed that services and agriculture are most prominent in the sub-region, averaging 42% and 37% of GDP, respectively (1999). Industry follows in third place. The services sector averaged 19% and 21% of GDP

in 1990 and 1999, respectively. The services sector is dominant in the urban areas, whereas agriculture

dominates in the rural areas. It is worth noting that the type or intensity of activities did not change significantly over the 1990 to 1999 period.

The agriculture sector (Table 11) contributes between 35-40 percent of the value added to GDP in all countries, except Côte d'Ivoire, where it contributed only 1/4th. Côte d'Ivoire, and Mali, also experienced a negative agricultural value added growth. Also, agriculture value added per worker varied considerably: lowest contribution in Burkina Faso (\$185), followed by Mali (\$265), Togo (\$528), Ghana (\$574), Benin (\$627) and highest in Côte d'Ivoire (\$1085). As agriculture sector value added per worker is a measure of agriculture sector productivity/efficiency, it implies that with the minor exception of Côte d'Ivoire, agriculture sector is least efficient, which points to capacity constraints, underemployment, low productivity, market distortions, and poor infrastructure in these countries.

Table 12: Macroeconomic indicators in the Volta Basin countries (2001)

Country	Per capita GDP (constant 1995 \$US) 2001	GDP growth (annual %)	GDP per capita growth (annual %)	Agriculture value added (% of GDP)	Agriculture value added per worker (constant 1995 US\$)	Agriculture value added growth (annual %)	Employment in sector (% of total employment) *			Aid dependency ratio	
							Agriculture	Services	Industry	% of GNI	% of gross capital formation
Benin	423.75	5.00	2.31	35.53	626.78	3.10	63.50	28.40	8.10	11.62	60.05
Burkina Faso	250.22	5.64	3.09	38.22	185.21	7.05	92.40	5.80	1.80	15.68	61.68
Côte d'Ivoire	714.62	0.90	-3.30	24.25	1084.8	-1.60	60.00	30.50	9.60	1.91	18.17
Ghana	421.02	4.00	1.88	35.90	573.95	3.69	62.20	27.90	10.1	12.72	51.23
Mali	291.60	1.44	-0.88	37.77	264.57	-13.0	85.80	12.20	2.00	13.92	62.65
Togo	322.33	2.70	-0.09	39.44	527.77	4.78	65.50	24.40	10.10	3.77	17.86

Data source: (WDI, 2003); WDR (2003); WDR (2004); *=latest year available during 1990s; years may not be same for various sectors, so figure may not add to 100.

The last column in (Table 12) shows aid dependency ratio, defined in terms of foreign aid as a percentage of gross national income and gross capital formation. The number shows that relative aid dependency is lower for Côte d'Ivoire and Togo, while for all other countries, over 50-60 percent of gross capital formation depends on foreign aid. In general, the basin countries have high debt burden, with average debt burden being about 72 percent of GNP (WDR, 2003).

Also, the region is saddled with a heavy burden of external debt as indicated in (Table 11). The debt burden ranges from 32% to as high as 122% of the GNP. The average, as well as the median, debt burden for the sub-region is about 70% of the GNP. This poor economic situation can potentially inhibit any meaningful sound environmental development about the exploitation of natural resources for socio-economic development. The other important measurement of external debt is debt servicing as a percentage of exports of GDP and the value of debt in relation to the value of exports. In Burkina Faso for example, debt servicing in 1996 was equivalent to 56% of GDP, and 19% of export revenues from goods and services. The present value of Burkina Faso's external debt in 1996 compared to the value of its exports was 254%.

Following the United Nations typology, Human Development Index (HDI), Human Poverty Index (HPI), and Gender Development Index (GDI) can be used to construct relative poverty profiles and discern gender differentials, if any, for the Basin countries. Of the two poverty indices, the HDI is a combination of *longevity* (life expectancy), *knowledge* (literacy and education) and *standard of living* (purchasing power and cost of living), where the lower the figure, the worse is the state of human development in a country. The HPI converts this into a poverty index, which becomes a measure of economic development and human welfare. The HPI is a measure of *human deprivation* defined in terms of *survival* (percent dying under 40 years), *knowledge* (illiterate adult percent) and *economic provisioning* (health, sanitation, infant mortality). Therefore, both HDI and HPI are composite indices and the underlying conception of poverty is that of multiple human deprivations.

The GDI measures women's achievement using the same indicators as the HDI. The divergence between HDI and GDI for each setting measures the relative status of women's opportunities. The closer the values of the HDI and GDI, the greater the degree of gender equality is. For all countries, the GDI is consistently lower than HDI, with the divergence being highest in Mali, followed by Burkina Faso and Benin, but lowest in Togo and Ghana (Table 13). This implies that gender disparities do exist in all the countries, and remain widely pervasive except in the last two

The human development index (HDI) of Burkina Faso is among the lowest in the world 0.320. Out of the six riparian countries of the basin, only Ghana is ranked medium human development country.

Table13: Human development indicators in the Volta basin countries (1998)

Country	HDI	HDI rank	HPI (%)	HPI rank	GDI	GDI rank
Benin	53.5	157	48.8	74	0.391	132
Burkina Faso	44.7	172	58.4	84	0.290	142
Côte d'Ivoire	46.9	154	45.8	72	0.401	129
Ghana	60.4	129	35.4	59	0.552	105
Mali	53.7	165	51.4	81	0.371	137
Togo	49.0	145	37.8	63	0.448	120

Data source: HDR (2000)

In terms of population below international poverty lines, Mali, Burkina Faso, and Ghana have about half to two-thirds of their population living below \$1 a day poverty line, with mean income shortfall varying between 1/4th to 1/3rd of reference poverty line. For \$2 a day, poverty rates are extremely high.

Table14: Poverty and income distribution in the Volta Basin countries (latest year available)

Country	Rural population below national poverty line (%)	Urban population below national poverty line (%)	Population below \$1 a day (%)	Poverty gap at \$1 a day (%)	Population below \$2 a day (%)	Poverty gap at \$2 a day (%)	Gini Index (%)	Income share held by lowest 10%	Income share held by highest 10%
Benin	33.00*	33.00*
Burkina Faso	51.0	16.5	61.2	25.5	85.8	50.9	55.1	2.0	46.8
Côte d'Ivoire	36.8*	36.8*	12.3	2.4	49.4	16.8	36.7	3.1	28.8
Ghana	34.3	26.7	44.8	17.3	78.5	40.8	40.7	2.2	30.1
Mali	72.8	37.4	90.6	60.5	50.5	1.8	40.4
Togo	32.3*	32.3*	33.8**

Data source: WDR (2003); (WDI, 2003). *=National figure; **= Deininger and Squire (1997) database/WIID (2000).

Gini index is a measure of income inequality, with value ranging between zero and one. A zero value shows a perfectly equal distribution of income, while a value of one implies perfect inequality. The higher the Gini value, the higher is the degree of income/consumption inequality. Income/consumption share held by highest/lowest 10 percent is the share that accrues to respective sub-groups of population indicated by deciles. The numbers show that poorest countries, such as Burkina Faso and Mali, also have highest income inequality with Gini coefficient ranging between 50 to 55 percent.

During the 1980's, most of the governments of the riparian countries began negotiations with the World Bank and agreed on a Structural Adjustment Loan. This was followed in the 1990's by an IMF Enhanced Structural Adjustment Facility. There was a considerable degree of opposition to these relations with the IMF and World Bank from opposition parties and trade unions in most riparian countries.

Adjustment programmes in key sectors like education and health, agriculture, the environment and public enterprises accompanied these macro-economic programmes. Funds were loaned to improve the sectors condition of these sectors regarding certain changes in their operation, as prescribed by the Bank. This high degree of funding for sectoral adjustment from the start was unusual in sub-Saharan Africa.

Attempts to increase government revenue in most of the riparian countries through reforms of the tax system and improved collection largely failed. For example, in Burkina Faso, a large wage increase in the public sector in 1991 increased government expenditure and a large budget deficit ensued.

External debt continues to place a heavy burden on the Burkina economy. Total outstanding debt almost tripled from 41.3% of GDP in 1981 to 85.6% of GDP in 1994. Although the Paris Club of debtors rescheduled some debt in 1991 and 1993, this made a little difference. In a country such as Burkina Faso, government expenditure on debt servicing greatly depletes funds available for essential social development. The money that Burkina Faso desperately needs to spend to improve education and health provision is rather used to pay debt servicing. Under HIPC Burkina Faso will receive \$200 million in debt relief, which although is not a huge amount, could still be used to improve education and health-care provision.

Table 15: External Debt of the Riparian Countries (1998)

Country	Millions of Dollars	% of GNP
Benin	1,647	46
Burkina Faso	1,399	32
Côte d'Ivoire	14,852	122
Ghana	6,884	55
Mali	3,202	84
Togo	1,448	68
Average	4,905	70

(World Development Report 2000/2001, 2001)

II.4 Agro-ecological Zoning

II.4.1 Drainage network and sub-basins

The Volta Basin is drained by several major rivers: the Black Volta, the White Volta with the Red Volta as its tributary, the Oti River and the Lower Volta. The mean annual flows of the Black Volta, White Volta, and Oti River are $7,673 \times 10^6$, $9,565 \times 10^6$, and $11,215 \times 10^6$, respectively (MWH, 1998-Table 15). The Oti River with only about 18% of the total catchment area contributes between 30% and 40% of the annual flow of the Volta River System. This situation is due to the steep topography and the relatively high rainfall in the Oti sub-basin. A dominant feature in the basin is Lake Volta which covers about 4% of the total area in Ghana. It generates hydropower (1060MW) at Akosombo and Kpong about 100 km north of Accra. Some 95% of the generated power in the basin comes from these two sites. Table 12 presents the area of sub-basins and length of the main tributaries of the Volta River.

Table 16: Area and length of main tributaries in the Volta River Basin in Ghana

Sub-basins	Total Area (km ²)	Rainfall (mm)	Mean Annual Runoff (x10 ⁶ m ³)	Run-off Coefficient (%)	Length (km)
Black Volta	149,015	1023.3–1348.0	7 673	8.3	1,363.3
White Volta	104,749	929.7–1054.2	9 565	10.8	1,136.7
Oti	72,778	1150.0–1350.0	11 215	14.8	936.7
Lower/ Main Volta	62,651	1050.0–1500.0	9 842	17.0	
Total	400,710	876.3–1565.0			

The White Volta began as the Nakanbé River in Burkina Faso. The Red Volta, referred to as Nazinon in Burkina Faso, and Sissili are tributaries of the White Volta and they all have their source in Burkina Faso. The mean annual flow of the White Volta Basin is estimated to be about 300 m³/s where the percentage of flow from outside Ghana to the total flow is estimated to be 36.5% (UNEP, 2002).

The Sourou from Mali and the Mouhoun from Burkina Faso join in the latter country and flow downstream to Ghana as the Black Volta. In Burkina Faso, apart from the Mouhoun, all of the rivers, including the Nakanbé, Nazinon and Sissili, dry up for approximately two months in a given year. The mean annual flow of the Black Volta at Bamboi is about 200 m³/s, out of which about 42.6% originates from outside Ghana.

The Oti River begins in the Atakora hills of Benin at an altitude of about 600 m and flows through Togo and Ghana. In Benin, the Oti River is referred to as the Pendjari River. Tributaries include the Koumongou, Kéran, Kara, Mô, Kpanlé, Wawa, Ménou, and Danyi Rivers. Due to the regularization by the Kompienga Dam in Burkina Faso, the Oti River has a permanent flow with an annual average flow of 100 to 300 m³/s, and can reach more than 500 m³/s. Virtually all the tributaries stop flowing during the dry season, however, and their annual average flows are only in the range of 5 m³/s. In Ghana, the Black Volta, the White Volta and the Oti join the main Volta at Volta Lake, which was created by the Akosombo Dam.

II.4.2 Relief

The basin is flanked by a mountain chain on its westernmost section. From the sea and northeastwards rise the Akwapim ranges, followed by Togo Mountain, Fazao Mountain, and the Atakora ranges in Benin. The Kwahu plateau branches northwestwards after the Akosombo Gorge. The only other significant relief on the western part of the basin is the plateau of Banfora. The basin in general has a low relief with altitudes varying between 1 and 920 m. The average mean altitude of the basin is approximately 257 m, with more than half the basin in the range of 200 – 300 m.

The global slope index is between 25 and 50 cm/km. Some of the characteristics of the relief are shown in Table 17.

Table 17: Some Important Relief Characteristics

Elevations at MSL (m)	Black Volta	White Volta	Oti	Main Volta
Minimum altitude	60	60	40	1
Maximum altitude	762	530	920	972
Average altitude	287	270	245	257

Source: ORSTOM Hydro. Monographs edited by Moniod et al., 1977.

II.4.3 Soils in the Basin

The soils of the Main Volta Basin in the sub-humid Savannah Zones are Savannah Ochrosols, Groundwater Laterites, Savannah Ochrosols – Groundwater Laterite (GWL), Savannah Ochrosol–GWL Intergrades, Savannah Ochrosol – Rubrisol Intergrades, Tropical

Black Clays, Luvisols, Tropical Grey Earths, Sodium Vleisols, and Savannah Gleysols. The soils of the Oti Basin are Savannah Ochrosols, Groundwater Laterites, Savannah Ochrosol-GWL Intergrades, Savannah Lithosols, Savannah Gleysols, and Forest Lithosols. The major soil groups in the Black and White Volta are Savannah Ochrosols, Groundwater Laterites, Savannah Ochrosols–Groundwater Laterite Intergrades, Savannah Lithosol, Savannah Gleysols, Savannah Ochrosols – Rubrisol Intergrades, and Savannah Gleysol–Luvisol Intergrades.

Table 18 gives a detailed description of identified soil groups in the basin

Table 18: Identified Soil Groups in the Volta River Basin of Ghana

Soil Group	Predominant Relief	Predominant Texture	Erosion Hazard
Savannah Ochrosols	Upper and middle Slopes gently Undulating	moderately heavy to light	Moderate sheet and gully erosion
Groundwater Water Laterites (GWL)	Near level to level lower slopes to valley Bottoms	Light over concretions and Iron pan	Severe to very severe sheet erosion
Savannah Ochrosols GWL Intergrades	Gently undulating to level middle to lower Slopes	Medium to light	Moderate to severe sheet erosion
Savannah Lithosols	Summits with steep Slopes	Medium to light	Severe gully erosion
Savannah Gleysols (GLE)	Near-level to level lowlands	moderately heavy to very heavy	Slight sheet erosion
Savannah GLE-Alluviosol Intergrades	Lowland terraces	Light to very light	Moderate to slight sheet

Erosion

In the Volta Basin in Burkina Faso, the major soil types found are:

- Lithosols and Aridisols: They are characterized by little or no chemical and biological alteration, traces of organic matter to a depth of less than 20 cm and an almost non-existent biological activities
- Degraded Ferruginous soils or Ferralitic soils: they are slightly or moderately unsaturated and have developed on material that is deficient in humus and material resulting from schist and sandstone. They are characterized by the complete alteration of the primary mineral, an abundance of synthetic products (aluminosilicates, iron hydroxides and oxides, etc.) and, in some cases highly developed organic matter.
- Ferruginous soils: they occupy most of the basin (central plateau and south). They usually contain low levels of organic matter. These soils can be with low or high cultivation according to their location
- Halomorphic soils: they have a layered structure and are fairly rare in the Volta basin
- Hydromorphic soils: they occupy the low-lying areas and alluvial plains
- Vertisols: mainly found in the valleys and usually dark in colour. They are characterized by the

presence of highliexpansive clay which swells and contracts alternately under the effects of very wet and very dry periods.




Figure 10: Distribution of soil types in the Volta basin of Ghana and Burkina

II.4.4 Geology of the Basin

The geology of the main Volta is dominated by the Voltaian system. Other geological formations include the Buem formation, Togo series, Dahomeyan formation, and Tertiary-to-Recent formations. The Voltaian system consists of Precambrian to Paleozoic sandstones, shales and conglomerates. The Buem series lie between the Togo series in the east and the Voltaian system in the west. The Buem series comprise calcareous, argillaceous, sandy and ferruginous shales, sandstones, arkose, greywacke and agglomerates, tuffs, and jaspers. The Togo series lie toward the eastern and southern parts of the main Volta and consist of alternating arenaceous and argillaceous sediment. The Dahomeyan system occurs at the southern part of the main Volta Basin and consists of mainly metamorphic rocks, including hornblende and biotite, gneisses, migmatites, granulites, and schist.

The Oti Basin is underlain mainly by the Voltaian system, the Buem formation and the Togo series. The White Volta Basin comprises the Birimian system and its associated granitic intrusives and isolated patches of Tarkwaian formation. The other significant formation is the Voltaian system. The Birimian system consists of metamorphosed lavae, pyroclastic rocks, phyllites, schists, tuffs, and greywackes. The Black Volta Basin consists of granite, the Birimian and Voltaian systems, and, to a minor extent, the Tarkwaian system. The Tarkwaian formation consists of quartzites, phyllites, grits, conglomerates, and schists. The underlying rocks of the basin have no inherent porosity. Thus, groundwater storage occurs only in fractured zones of the rocks. Figure 11 shows the geology of the entire Volta River Basin.

Volta Basin - Geology




Figure 11: Geological Map of the Volta River Basin.

II.4.5 Land use, land ownership and land degradation




Figure 12: Land cover Map of the Volta River Basin.

As agriculture and animal husbandry are the primary economic activities in the basin, land resources are critical to the basin inhabitants. The resources currently meet these needs, but the growing population pressure that will require additional land, combined with the anthropogenic and climatic threats to land resources, suggest that this might not always be the case.

Information on demand for land resources is inadequate in the basin. Statistics of land use were given, for example, in Togo and out of a basin area of 2,670,000 ha only 428,000 (16%) were put under cultivation in 1995, while an area of 528,420 ha (19%) was under forest reserve. Fertility of the soils was not discussed, but this could be a limiting factor due to the expansion of agricultural lands. For the Volta Basin in Burkina Faso, 3,905,500 ha, representing 22.5% of the basin area were under cultivation. It was indicated that land availability in that area for farming is becoming limited but is not yet in a crisis situation. Throughout the basin, the loss of soil fertility due to erosion, over-use of manure, and uncontrolled bushfire were identified as problematic issues.

Farming practices could determine the size of land needed for future activities. With the increase in population, the available data and information suggest that demand for land for farming will increase with a view to achieve food self-sufficiency in the basin, as well as increasing food exports. Population pressure has not only expanded pasturage, but has reduced its quality due to reduced crop rotation and not allowing insufficient time for croplands to lie fallow to regenerate essential nutrients.

Farming and animal husbandry are significant contributors to land degradation in the basin. Agricultural practices in the basin have in the past included crop rotation and leaving fields fallow for a

period of time. With rising population, however, the fallow periods have been reduced and crop rotation declined, leading to the loss of soil fertility and less food production per unit area of cultivated land. Increasing livestock production has resulted in the loosening of soils and the degradation of vegetation, both of which exacerbate erosion. Increased exploitation of forested areas is also a significant contributor to soil degradation and erosion.

Forested areas are cut to provide additional land for agriculture and animal husbandry, and to provide fuel. Also, timber resources are over-exploited in many parts of the basin. This is done to meet rising demands for foreign exchange, as well as to meet increasing domestic needs. Unfortunately, the timber exploited is not processed for higher value and thus more volume of timber is required to be exported for adequate foreign exchange receipts for the countries' socio-economic development. In the long-term, these practices are not sustainable and have detrimental effects on both land and water resources in the region.

Land degradation in the basin encompasses soil degradation, intense erosion and desertification. As discussed above, the basin's population heavily depends on the land resources of the region for subsistence agriculture and livestock breeding. The increasing demographic pressures have resulted in the overuse and misuse of land resources. Soil degradation, erosion, and desertification processes manifest themselves in low agricultural productivity, destruction of the soil's natural productive capacity, compacting of the soil, degradation of water quality, and loss or reduction in vegetation cover. The increased mobility of sediments also affects reservoir capacities and their useful lives.

Land ownership in the region remains primarily traditional, meaning that lands are often owned or managed by local elders or leaders. As a result, the major institutions involved in land administration are the traditional leaders. Thus, a significant problem associated with land resources is the institutional and legal framework governing the release of land for use. Some lands are also preserved as natural habitats for flora and fauna and are unavailable for use however, illegal exploitation of the land resources has reduced their value.

Further, some lands are already degraded to the point of non-productivity. For example, in the Lower Volta Basin, the establishment of the Akosombo Dam has rendered some of the soils in the area more acidic. As a result, yields from farms in the region have been reduced considerably. The potential of such lands has been reduced and will require remedial measures.

Land degradation in the region has both transboundary causes and effects. Transhumance, defined as the movement of cattle, sheep, and people across national boundaries, is common within the basin. This phenomenon is usually accompanied by reckless destruction of vegetation, watering sources, etc. The situation also creates social tension and disruption of socio-economic activities, sometimes proving fatal.

Bushfire has no respect for national boundaries and can move from one country to another country in the basin. This phenomenon of bushfire across frontiers does happen in the basin and constitutes a transboundary cause. While controlled bushfires are used to enhance the fertility of agricultural lands, many of the bushfires occurrences in the region whether intentional or unintentional region can easily get out of control and burn large areas.

Deforestation occurs across frontiers, particularly where transhumance is a major problem as in the basin. The animals being moved are fed on leaves from trees illegally cut down by the

herdsmen. In addition, the trampling of the soils by the hooves of the animals renders the soil loose for erosion. Deforestation also occurs across borders when there are inadequate laws in neighboring countries. For example, people from Burkina Faso travel to Mali because the laws are less strict there.

Population pressures in countries with a weak economic base, as in the basin, induce unsustainable use of forest and land resources. The easy movement of people across national boundaries in the sub-region under the ECOWAS protocol makes population pressure a transboundary cause of the above-mentioned environmental problems in the basin.

The transboundary nature of the effects of soil degradation and erosion arise mainly in the sediment transport and degradation of water quality. Due to erosion occurring upstream, sediments fill river channels and reservoirs, and decrease water quality. Further, the transhumance of livestock occurs when new pastures must be found due to land degradation.

Socio-economic impacts related to land degradation is extensive, but difficult to assess. Land degradation, however, has serious consequences on the ability to produce food in the region, which in turn has serious consequences on human health and security.

i. BENIN

Only a small amount of land is suitable for agriculture, livestock, and for dwellings in the Volta Basin of Benin. As a result, competition exists over these finite resources and the region is experiencing significant demographic pressure. Land pressure also hinders economic development in the region.

The majority of land in the basin is owned and is passed down by families, rather than sold to outsiders. Additionally, communes remain under the control of managers who determine how lands are divided. Thus, the land tenure system remains somewhat traditional as local leaders have significant control.

Land resources in the basin are seriously threatened by anthropogenic activities, and this in turn threatens waterways. The overexploitation of the vegetation occurs as a result of overgrazing by livestock caused by the increasing density of the zone’s population. The abusive use of artificial fertilizers and pesticides, the reduction of the duration of fallow periods, and other poor agricultural practices scour the land and deplete the soil’s minerals. Bushfires accompanied by hard rains and strong winds further accentuate erosion and add to the siltation of waterways. Further, the destruction of forests and the deforestation of riverbanks exacerbate the degradation of the land and threaten the Oti River. Table 19 below gives the characteristics of the agro zones in Benin.

Table 19: Characteristics of the Zones

Agro Zones	Ecological Practices of Agricultural Systems	Problems	Causes
<ul style="list-style-type: none"> • West Atacora • Cotton • Oti National Park 	<ul style="list-style-type: none"> • Burning of biomass • Farming of hollows • No use of mineral manure • Food crops: sorghum and millet, niébé, groundnut, fonio, voandzou, yam, maize, and rice • Breeding: system of breeding based on transhumance and 	<ul style="list-style-type: none"> • Degradation of the soils and vegetation cover • Food security • No market for groundnut and rice • Conflicts between livestock breeders and farmers 	<ul style="list-style-type: none"> • Farming practices • Demographic pressure • Ferruginous soils with inadequate water and poor fertility except in the hollows • Lack of water sources and pasture

	changing pastures		
--	-------------------	--	--

ii. BURKINA FASO

Burkina is endowed with significant natural resources. Apart from arable lands (9,000,000 ha), forestry and pastoral lands (16,000,000 ha), there are nearly 1,200 water courses and dams covering a total area of 120,000 ha to 150,000 ha. These natural water bodies (Mouhoun, Nakambé, Nazinon, Comoe, etc) offer potential fishery resources of close to 12,000 tons per year (Kafando, 1995).

Five agro-ecological have been identified by the CNRST, based on agro-climatic, socio-demographic conditions and regional constraints and potentials. These zones have the following characteristics:

- The Eastern zone covers an area of 60,600 sq.km with a population density of less than 20 inhabitants per sq. km. It has an annual rainfall of between 500 mm and 1000 mm. It has great potential in animal husbandry, fishery and wildlife resources (5 reserves). The farming system is dominated by sorghum and millet.
- The Sahelian zone covers an area of 36,896 sq. km with a very low population density of 10 to 17 inhabitants per sq. km; it has a rainfall of between 300 mm and 600 mm. and a highly degraded vegetation cover due to drought. Its agriculture is subsistence and consists mainly of cereal. The main source of income is animal husbandry.
- The North-western zone has an area of 30,870 sq. km., a rainfall of between 500 mm. and 800 mm. It has an average population density of 41.1 inhabitants per sq. km. Although the farming system is based on sorghum and millet cultivation, the zone has great potential for irrigated crops like vegetables, rice and maize. Animal husbandry is also significant.
- The Central zone covers an area of 94,000 sq. km with a high population pressure resulting in severe degradation of the plant cover and soils. Rainfall varies between 600 mm and 900 mm. The agro-pastoral and forestry potentials are limited and the farming system which consists of agriculture combined with animal husbandry, is dominated by small ruminants and poultry.
- The Western zone has an area of 52,000sq.km, and a rainfall of between 700 mm and 1200 mm. This zone has a great potential for natural resources. It is the converging area for immigrants from the Central and Northern parts of the country. Its agriculture is much diversified with high output in cotton, fruits, vegetables and cereal dominance (sorghum, maize, millet, rice and fonio). It has a high potential for animal husbandry, and possesses a large stock of cattle.

Land in Burkina Faso is threatened by agricultural practices, deforestation, and, in some areas, by mining activities. The land tenure system is governed by the Land and Agricultural Reform Law. This statutory framework has been read through several times with a view to adopting the law in the socio-cultural context of the regions of the country, but there has not been a bold application of the law due to the existence of customary laws on land ownership. This situation should improve soon. An FAO

study in 1987 revealed four (4) types of land holding rights in force in the rural areas of Burkina:

- the right to permanent use for each member of a lineal group with the collective ownership right vested in the head of the family;
- the right to permanent use as a result of the clearing and development of vacant land;
- the right to use inherited from a farmer who has himself been granted that right by the holder of a collective ownership right; this is usually permanent;

The provisional right of use lent out by the holder of the right to permanent use or a collective ownership right (lending or hiring). Under the framework of the Agrarian and Land Reform Law (RAF), the National Lands Sector (DFN) which is the right to exclusive State ownership was established in 1984. Despite the existence of these basic land tenure regulations, there is also the existence almost throughout the whole country, of the customary law governing lands placed under the authority of "Land Chiefs" who assume religious and legal functions in the villages. Their duty is to ensure that land is used in accordance with the rules laid down by society. The underlying principle is to ensure that every man, whether he belongs to a village community or not has a right to adequate land, in order to support himself and his family. This redistribution of lands to those who do not possess any is in the form of land lending or renting, which is unpaid for.

However, the only requirement on the beneficiary is the compliance with certain socio-cultural rules of the lender or the receiving community (in the case of immigrants). Apart from this land redistribution system, the common method of land possession is through inheritance. These land acquisition methods enable everyone, no matter his status in life, either rich or poor, to have access to land. However, in most cases, the right to land ownership is the exclusive preserve of men. Women can have access to it for subsistence farming, through the head of the household or family. On redistribution of land, especially through lending, any action which tends to jeopardize the ownership right of the lender is forbidden, such as investments leading to permanent occupation of the land.

In the Mouhoun Basin, land tenure is based on social stratification. It is also based on the right of collective appropriation distributed between the founders of a village and the right of temporary or permanent use of the land allotted to an individual. Individual appropriation does not exist, however, and land rights can only remain within the social group.

In the Nakanbé Basin, land tenure is related to the existing structure. The area is occupied primarily by animal herders who require a large amount of land for pasture, but the land is not used intensively. The land is collectively owned by the group, and they do not have the right to refuse an outsider the use of the land if they have valid reason.

Forested areas in Burkina Faso are shrinking significantly as population pressure in the region increases. In the National Action Plan for the Fight Against Desertification, it is noted that forested areas decreased by 1.26 million hectares between 1980 and 1992. It was estimated in 1996 that 105,000 ha of forested areas were yearly lost. About 880,000 ha are classified forests and 8,790,000 ha are protected forests. In the Volta Basin about 788 433 ha of forest have been considered classified forests since 1954.

The Volta Basin in Burkina covers a total area of 172, 968 sq.km (17.3 million ha) and is primarily occupied by tropical ferruginous soils and degraded land less favorable to subsistence farming and cashcrops. It is estimated that about 9 million hectares representing 1/3rd of the country's

total area, is arable land. However only 3.5 million hectares are yearly cropped for subsistence farming based on staple cereals: millet, sorghum and maize. The three cereal crops cover 88% of the total cropped land in the country. The average yield is about 850 kg/ha.

In 2000, about 2,927, 737 ha were cropped in the basin. This is an increase of 12% compared to 1997 when about 2,612,245 ha were cropped. According to the Agricultural Strategic Plan in 2000, the cropped area in the Volta Basin represented 82.5% of the total cropped land in Burkina. About 12,000 ha of land were also planted with fruit trees. One of the objectives of the Strategic Plan is to increase agricultural production by 5 to 10% per year over the next ten years.

In short, the occupied surfaces in the basin are:

- agriculture: 3,000,000 hectares
- water bodies: 83,000 hectares
- human settlement: (not estimated)
- roads: 34,000 hectares (16300 ha representing 38 161 km rural dirty roads and 17530 ha of 9100 km of improved roads)
- classified forests: 788,500 hectares

Table 20: Major landuse types in the Volta basin of Burkina (hectares)

Agriculture	3 000 000
Water bodies	83 000
Human settlements	Not Estimated
Roads	34 000
Forests	788 500
Total	3 905 500

The total of **3,905,500 hectares** represents 22.5% of the total surface of the basin.

iii. CÔTE D'IVOIRE

The rural areas in the basin tend to follow the traditional system of land tenure. At the village level, each "great family" has a field on which the members cultivate. The appropriation of the ground is thus collective, but its exploitation is individual.

Today, land and water resources are subjected to the modern system of land tenure defined by the laws of the Rural Land Code and Water Code promulgated in December 1998. These laws aim to protect the resources from overexploitation.

As in the rest of the basin, agriculture is the dominant economic activity and a significant contributor to land degradation in the Volta Basin of Côte d'Ivoire. Cotton is a main crop in the area, as well as anacarde, corn, sorghum, rice, and groundnuts.

The area is also used as pastureland to a significant extent. Herders come from Mali and Burkina Faso to use lands in Côte d'Ivoire.

Bushfire is used extensively in the region for hunting, managing pastures, preparing agricultural lands, and for other purposes. Uncontrolled bushfires contribute to land degradation.

The lands in the basin are occupied approximately in the following manner:

- 3% Infrastructure, urban areas, water, and rocky zones
- 12% State-owned land: national parks and reserves
- 75% Rural areas (50% savannah and forest, 25% cultivated)

iv. GHANA

Land ownership within the basin is basically traditional except for areas demarcated for control by the government agencies, such as the Volta River Authority, as well as forest reserves, wildlife, and national parks. The details of traditional ownership vary from place to place. As the system puts all resources under the control of the traditional authorities families and individuals have both access and control of resources through birth into a particular community or after payment of certain amounts either in kind or cash (Nyankpala Agricultural Research Station). In the northern sector of the Volta Basin, usufruct rights to land might not necessarily include rights to economic trees like the dawadawa and shea trees. While individuals and families might own lands along riverbanks, the rivers always remain communal or public property and are used as such.

Agriculture is the dominant economic activity within the Volta River Basin. As in other areas of the basin, soil is being rapidly degraded as a result of shortened fallow periods. This is especially pronounced in the northern parts of the basin.

Environmental problems arising from livestock production are becoming sources of great concern. The maintenance of large herds of livestock has tended to exceed the carrying capacity of the ecosystem, particularly in the northern part of the basin where mean annual rainfall is about 1000-1200 mm. During the dry season of November to April, large herds of cattle cross from the neighboring countries to graze on the limited fodder available. As a result of trampling, soils are severely exposed to erosion, and watersheds to rapid evaporation. The prolonged exposure of the soil renders it susceptible to erosion and reduces its regenerative capacity. In the northern parts of the basin, large tracts of arable land have become infertile and crop yields have declined tremendously.

Table 21: Erosion Hazards of the Volta Basin in Ghana

Volta Basin System	Erosion Hazard
Black Volta	<ul style="list-style-type: none"> • Northern Section: slight to moderate sheet erosion. • South-western Section: A combination of moderate to severe sheet and gully erosion but more of the latter with areas of very severe sheet and gully erosion. • SE Section: A combination of moderate to severe sheet and gully erosion but more of the latter.
White Volta	<ul style="list-style-type: none"> • Same as in the Black Volta Basin
Daka	<ul style="list-style-type: none"> • Combination of sheet and gully erosion but more of the former.
Oti	<ul style="list-style-type: none"> • Combination of moderate to severe sheet and gully erosion but more of the latter, especially within the central and southern sections
Lower Volta	<ul style="list-style-type: none"> • NS – Combination of moderate to severe sheet and gully erosion, especially the southern parts. The extreme northern part is however subject to slight sheet erosion. • CS – Moderate to severe sheet and gully erosion but more of the latter.

Volta Basin System	Erosion Hazard
	<ul style="list-style-type: none"> • SS – Slight to moderate sheet erosion within the savannah areas and severe to very severe gully erosion within the forest and highland areas. • ES – Severe to very severe sheet and gully erosion but more of the latter.

As increasing populations look for additional land to farm, deforestation often occurs. Although figures are not known for the Volta basin, nationally deforestation occurs at a rate of 2.5-5% annually in areas that are not forest reserves. However, within the forest reserves, deforestation occurs annually at a rate of 1.3%.

Bush burning is used to clear land for agricultural purposes, hunting, creating fire belts at the onset of the dry season, and inducing rapid re-growth of rangeland during the dry season, which often results in enormous damage to vegetation, wildlife, and properties because it is typically not controlled. The risk of bushfire is highest on the grazing lands in the savannah zones of the basin where as many as 120 to 150 outbreaks can occur within a single year. Along the border areas of the savannah zone, particularly in the Oti, White Volta, and Daka Basin, the problem of bushfire is especially severe, probably as a result of the association with transhumance.

Urbanization in Ghana is another cause of land degradation in several areas within the basin that are becoming population nodes as people migrate from the rural areas to urban centres in search of better livelihood and to escape tribal conflicts. Settlement growth in areas of the basin considered to be potential biodiversity conservation priority areas, particularly in the White Volta and Lower Volta Basin, is of great concern as important habitat is lost. Although no population statistics is available, however, it is believed that the population in designated protected areas within the basin has not changed significantly over the past decade.

Another problem of rapid urbanization is that infrastructure development often lags behind population growth resulting in the development of poor sanitation situations that adversely affect surface water resources. A report by EPA (2001) shows that surface water resources close to urban centres have exceptionally high faecal coliform counts.

Mining is a final cause of land degradation. Several small-scale artisanal groups carry out gold mining in areas underlain by the Birimian formation with little regard for environmental protection. Thus, their operations have led to serious degradation of the land in portions of the Black Volta and White Volta Basin. Limestone mining in the Black Volta basin and in the Lower Volta is also causing damage to land.

White Volta Basin

The predominant land use is extensive land cultivation which is two-to-six miles from the village on upland areas (NAES, 1993), with widespread grazing of large numbers of cattle and other livestock up to 100 cattle/km² (FAO, 1991); and compound cropping (home gardening) around the house (Wills, 1962; Adu, 1967; USAID/ADB, 1979; FAO, 1963; NAES, 1993). Estimates of land use and land cover in 1989 showed that about 50% of the land in the northeast and northern parts of the basin was in the compound and bush fallow cultivation cycle (IFAD, 1990). Farm sizes are usually less than three acres. Grazing land including that obtainable under natural condition is generally poor. Annual bush burning further reduces the quality and quantity of fodder.

Extensive valley bottoms in many parts of the basin, particularly in the guinea savannah areas, have in recent years been cultivated for rice under rain-fed conditions. In the north and northeast, the best agricultural soils are derived from granites, sandstones, and greenstones. These areas remain the most densely populated.

A long period in the upland areas away from the valley bottoms, which had been infested with the *Onchocerciasis simulium* vector, and the intensive cultivation and grazing without proper management practices have led to widespread soil erosion and loss of fertility of the upland soils. Outcrops of rocks, iron pan soils, as well as the scarps are usually avoided by farmers and may be uninhabited or only sparsely inhabited. Fuelwood and other wild produce gathering is widespread.

Urban land use is small and most intensive in such centres as Bolgatanga, Bawku, Wa, Navrongo, Tamale, and Tumu. Due to the decentralisation of administration to the district level, urban type land use is becoming important in some of the district capitals, especially those along major trunk roads.

Black Volta Basin

The major land use is agriculture with food crop cultivation under extensive bush fallow. The major food crops include yam, cassava, maize, sorghum, millet, groundnuts, and beans. Animal grazing on the free range is a significant activity. In the northwest of the basin, particularly the Lawra district, lands are highly degraded both in terms of physical status and fertility levels and can hardly support meaningful crop cultivation. Vegetation has also been degraded due to the incidence of annual bushfire. This has led to seasonal human migration and great reduction in the number of livestock.

Lower Volta Basin

Current land use is short bush fallow cultivation along the immediate banks of the river, and less intensive bush fallow cultivation elsewhere. Animal grazing is common while the lakeshores are extensively settled by fishing families. Charcoal burning involving the cutting of wood has become an extensive economic activity in the southern dry forest and transitional environments (e.g., various parts of the Afram sub-basin.)

The Afram plains and other areas in the south have been the focus of increasing settlement and agricultural development since the 1960s, having been generally thinly populated in the past as part of the empty "middle belt". The forest and transitional areas are intensively farmed with cocoa, coffee, plantain, cocoyam, cassava, oil palm, and maize on small bush fallow plots. A large modern commercial farm at Ejura specializes in maize production. Some timber extraction takes place in these areas.

Recent developments, particularly below the Akosombo Dam, include irrigated rice, sugar, and vegetable cultivation in the areas immediately adjoining the Volta River. The construction of the Akosombo Dam has reduced the annual flooding in the Lower Volta areas.

The areas around the coastal lagoons, such as the Songhor, are used for salt mining. Urban land use is limited to a few towns like Kpandu, Kwamekrom, Akuse Sogakpe, and Ada-Foah.

Oti Basin

Current land use and land cover are extensive bush fallow cultivation and grazing with tree savannah regrowth and small patches of reserved forest areas on the hills in the southeast. The main crops that are grown in the basin include yams, guinea corn, maize, rice, millet, and groundnuts. Fishing is common along the river while grazing, as in other parts of the savannah, is commonly practiced. Settlements within the basin are small.

Daka Basin

The predominant land use is bush fallow cultivation of yam, maize, and guinea corn with free grazing animals. A recent land use problem within the greater part of the Volta basin especially in the Black Volta, White Volta, Afram, Daka, and Oti sub-basins is the activity of alien herders who graze their large herds of cattle indiscriminately, leading to widespread destruction of vegetation and even crop farms. In some cases bushfires is set to hasten the re-growth of fresh vegetation leading to high rates of soil erosion and loss of soil productivity.

v. MALI

Land in Mali officially belongs to the state. This ownership, however, does not preclude the traditional authority, which manages land ownership according to the following criteria:

- The water and land belong to the head of the land (first occupants and their descendants). This title can be passed down through the family.
- These grounds can then be yielded, lent, pawned or sold to a third party.
- Village leaders play an important role in resolving land disputes.

Approximately 80% of the land in the basin is used for agriculture, livestock, or dwellings. The high population density in the region places enormous pressure on the land. There is competition between the livestock breeders and the farmers for scarce land and water resources. This is a consequence in part from the increasing population pressure on the land, thereby agricultural practices are not sustainable. Land is no longer allowed to lie fallow for a sufficient amount of time before they is replanted.

The basin of the Sourou River is considered to be the granary of the country, but poor agricultural practices have steadily degraded the land. The lands are now no longer very fertile and are prone to wind erosion. The degradation of the soil in Mali has resulted in a decline in production from 4 to 20% in the sahel zone and 8 to 20% in the Sudan zone.

There is a great deal of competition for resources between those agriculture and those raising livestock. The droughts in the northern region of Mali have resulted in livestock herders migrating into the Sourou region to find water. This transhumance results in significant destruction of the forests of Samori.

vi. TOGO

Togo is divided into six major agro-ecological zones and three are entirely located within the Volta basin. These are: the Dry Savannah zone in the North, the Oti sub-basin and the Southern dry zone of the Atakora Mountains. The dry Savannah zone in the North is dominated by three major soil units: deep weathered ferruginous soils, lithosols, and hydromorphic soil with gley while in the Oti sub-basin zone the major unit soils are shallow weathered ferruginous soils and in the southern dry zone of the Atakora Mountains they are tropical ferruginous soils, ferralitic soils vertisols and lithosols. Soil erosion is highly present in this zone. In Togo, the land resources are governed by a combination of local and tribal leaders and the national government.

Land degradation of the land in Togo results from a variety of factors. First, trees are harvested at an unsustainable rate in some areas as the demand for wood has increased. This increases erosion and desertification as the land cover is removed. Second, poor agricultural practices, such as the misuse or overuse of pesticides and fertilizers, have damaged soil resources. Finally, overgrazing of the land further exacerbates the problems of erosion and desertification.

Forest resources in Togo have experienced extensive degradation in recent decades due to population increase, unsustainable cutting of trees for firewood and charcoal, unsustainable cutting of sawlogs, clearing for agricultural use, and bushfire. The forests of the Volta Basin provide more than half of the national production of sawlog, and during the political crisis of the 1990s, much illegal cutting of forests took place. At the national level, it is estimated that forest cover is degraded at the rate of 15,000 ha/year.

While there are significant protected areas in Togo, these have been threatened by encroachment from those populations living around the reserves. In 1992, a national commission was formed to examine the areas facing the greatest threats, which suggested turning towards participatory management of protected areas.

Since 1970 when coffee and cacao trees and cotton production were introduced into the region, vegetation cover in the region has changed. During that period, there was significant immigration into the region as people came to grow these products. Considerable amounts of land were cleared for agriculture and livestock production. The agricultural practices, including shortened fallow periods, used in the region often result in land degradation. Table 22 shows that the areas under cultivation will continue to increase over the next decade.

Table 22: Rate of Occupation of Cultivable Land in the Basin in Togo (1,708,800 ha)

Years	Cultivable Area (ha)	Cultivated Area (ha)	Rate in %
1990	1,456,188	264,030	15.45
2000	1,291,759	434,014	25.39
2010	1,087,310	646,784	37.85
2020	782,632	963,862	56.4
2025	578,179	1,176, 637	68.85

Source : Projection à partir des données des Recensements Nationaux de l'Agriculture (1972, 1982, 1996 – DESA)

Livestock are also taking a significant toll on soil productivity in the region. Although there are sparse data on the specific effects of livestock, it is clear that they are negatively influencing the area. The areas that have experienced the most severe degradation include the Savannah and Kara regions. The Plateau (Danyi), the Central region (Fazao), Kara (Kantè), and Savannah (Dapaong) have experienced strong degradation. The Plateau (Danyi), Power Station, and of Kara have also experienced average degradation. The zones with weak degradation extend around the Togo Mountains, in the Plateau area (Danyi and Wawa), in the Savannah area (Mandouri), and Kara (National Park of Kéran).

Table 23: Evolution of Various Vegetation Formations in Togo (1979-1991)

Vegetation	Area (km ²)		Variation in % of the area
	1979	1991	
- Dense forests	2931	1264	- 56 %
- Mountainous forests	863	525	- 39 %
- Dry dense forests	677	315	- 53 %
- Regrown forests	1159	615	- 47 %
- Savannahs with trees	12922	6048	- 53 %
- Shrub savannahs	5138	2720	- 47 %
- Agriculture zones and others	1840	1944	+ 5.6 %

About strongly degraded savannah, the soil erosion was evaluated in 1969 to be between 1,500 and 2,000 tons per km² annually. The prefecture of Oti has records from the same time period showing from 600 to 1,500 tons per km². These figures can be multiplied by as much as four or six times to account for the current level of degradation.

The zones with weaker degradation are currently threatened by the phenomenon of savannisation and from impoverishment of the soil due to the disappearance of forests. While the causes and effects of land degradation have been described well, supporting data have been provided only sporadically and would need to be augmented. For many countries, information has not been provided on the areas that are experiencing the most severe degradation, the amount of soil lost to erosion annually, and rates of deforestation and desertification. Neither has a quantification of the loss of productivity of lands been provided for all countries. In addition, information on demand for land resources in the future has not been given, except in Togo.

II.4.6 Agriculture

i. Crop production

Accurate and specific data are not easily available on the economic output of the basin as these data are embedded in the national figures. It may be claimed, however, that agricultural production in the basin, which has a higher rural population than the national averages will not be less than 40% of the entire economic output of the basin. To demonstrate the importance of agriculture in the basin, some information from national reports is presented.

In Ghana, Table 23 shows the production levels of selected staple crops by regions in the basin. From the Table, it can be observed that the basin in Ghana produces 78% of the total national output of yams, 31% of cassava, 40% of maize, and 69% of rice. Table 24 shows the statistics of cereal production in two districts of the Volta Basin of Mali. These regions are considered to be the granary of the Mopti region and 85% of the local population is engaged in agriculture.

Table 24: Production Levels of Selected Crops by Regions in the Volta Basin in Ghana (Tones)

Region	Yam	% of National Total	Cassava	% of National Total	Maize	% of National Total	Rice	% of National Total
Upper East	-		-		16,280	1.6	65,379	26.3
Upper West	263,416	7.8	-		56,725	5.6	9,281	3.7
Northern	518,000	15.4	68,500	0.8	81,800	8.1	71,360	28.7
Volta	112,265	3.3	424,350	5.2	48,980	4.8	14,530	5.8
Eastern*	529,014	15.7	767,460	9.5	97,014	9.6	2,250	0.9
Ashanti*	186,248	5.5	373,674	4.6	12,530	1.2	706	0.3
Brong Ahafo*	1,000,337	29.7	854,659	10.5	91,985	9.1	32	0.0
Greater* Accra	-		38,603	0.5	2,269	0.2	8,469	3.4
Total in Basin	2,609,280	77.6	2,527,246	31.2	407,583	40.2	172,007	69.1
National Total	3,363,000		8,107,000		1,013,000		249,000	

Source: SRID, MOFA, 2000

*Figures are totals for districts that fall within the Volta Basin, whether wholly or partially.

Table 25: Cereal Production in the Mopti Region of Mali (Tones)

KORO											
Cereals/Years	1990-1991	1991-1992	1992-1993	1993-1994	1994-1995	1995-1996	1996-1997	1997-1998	1998-1999	1999-2000	2000-2001
Millet	67930	64770	66440	67800	67800	66735	65500	68225	68400	68700	69000
Sorghum		3500	3950	3660	3760	4400	4400	4275	4300	4600	4650
Paddy rice	320	270	270	360	330	380	350				4150
Niébé	1545	3245	3640	980			1950	1700	2320	2370	2426
Fonio	5020	4190	4264	4990	5350	4530	3380	4030	1940	1940	2000
Groundnuts	1620	5470	5750	5520		5985	5670	6150	5975	6030	6324
BANKASS											
Cereals/Years	1990-1991	1991-1992	1992-1993	1993-1994	1994-1995	1995-1996	1996-1997	1997-1998	1998-1999	1999-2000	2000-2001
Millet	48700	42600	44300	45800	45900	45600	46900	46800	47500	48000	48945
Sorghum		6300	8100	8300	8500	9230	9900	9900	9800	9800	10500
Paddy rice	170	275	1050	1600	1900	3250	3800	3900	5600	5400	4150
Niébé	2490	4200	4000	1600	1450	1360	1500	1700	1800	1875	2050
Fonio	4900	4650	4850	4900	4825	4330	4100	3950	3500	3500	3335
Groundnuts	1760	4750	4800	5800	5800	4900	5650	5100	5850	5750	5880

Crop production in Côte d'Ivoire is presented in **Table 26**.

Table 26 : Crop Production in Côte d'Ivoire (1996)

Produce	Quantity (Tonnes)
Cocoa	816
Coffee	1,099
Cotton	1,511
Cashew nut	12,482
“Roucou”	1,951
Yams	225,703
Maize	14,500
Cassava	42,695
Rice	2,341
Groundnut	4,330

In Burkina, agriculture is the main activity and it involves mainly food crops, with a clear predominance of cereals. The 1990 MEE/CILSS study which included data for the period covering 1961 to 1988 indicated that the following are the main food crops: sorghum (which takes 51 % of the country's rain-fed cereal area), millet (40 %), maize (6.7 %), rain-fed rice (1.6 %) and fonio (10.7 %).

The main cash crop is cotton, followed by pulses and oil seeds (groundnuts, sesame, cowpea, Bambara bean, etc.), fruits and vegetables (citrus, mango and vegetable products). Table 27 below shows the production trend between 1992 and 1996 in Burkina Faso

Table 27: Production trends in Burkina Faso (x1000 tons)

Crop	Year				
	1992	1993	1994	1995	1996
Cereals	2091.6	2157.1	2099.5	1936.6	2054.6
Oil seeds and pulses	239	306.3	310.8	293	301.1
Cotton	163.3	116.6	143.1	151.3	170
Fruits and vegetables	319.1	331.9	345.2	360.7	376.9

Source : CC-PASA (1997)

In the Mouhoun sub-basin, cotton and maize are the major crops. During the 1997/1998 season the cotton production in the sub-basin accounted for about \$50 million in revenue leading to a substantial increase in cultivated land the following years (from 250,000 ha in 1997/1998 to 400,000 ha in 2001/2002). Cereal production (maize, fonio, millet, sorghum) covers 60% of the cultivated land while cash crops (cotton, groundnuts, etc.) cover only 30%.

Like in most of the riparian countries in the Volta basin, for Togo, productivity in the agriculture sector is low. Agriculture employs 59% of the active population and is widely dominated by a subsistence farming system. Most of the 3.5 millions arable land in Togo

are located in the Volta basin and are used for staple crops (maize, fonio, yam cassava, millet sorghum and rice, peanuts etc.) and for cash crop (cotton, coffee, cocoa etc.) Cash crop production is almost entirely intended for export (100,000T of cotton, 20,000T of coffee and 20,000T of cocoa etc.)

Irrigated land as a percentage of cropland for 1995–97 for Benin, Burkina Faso, Côte d’Ivoire, Ghana, Mali, and Togo are 0.8, 0.7, 1.0, 0.2, 2.1, and 0.3, respectively (World Development Report 2000/2001, 2001). Thus, crop production under irrigation is negligible in the sub-region as most arable farming is predominantly rainfed. With current climate change, rainfall is believed to be becoming more variable and unreliable. Extensive crop farming coupled with variable and unreliable rainfall patterns in a region where poverty is predominant has far-reaching implications for the environment and food security.

In the Nakanbe sub-basin alone (in Burkina) more than 400 dams and small reservoirs have been built over the past years to develop irrigation and generate electricity. Thus, about 47 irrigation schemes have been identified but only 2,620 ha have yet been developed and out of that 1,000 ha are for the Bagre dam alone. An estimated 2,175 ha of inland valleys are to be added to the potential of irrigable land in the basin. Irrigated land in the Nakanbe sub-basin is relatively small compared to the vast potential for irrigation that the large volumes of water stored behind the numerous dams can offer.

ii. Livestock

The rich savannah grassland provides good fodder for livestock production. In Ghana, data show that animal husbandry in four regions, the Upper-East, Upper-West, Northern and Volta, which fall exclusively in the basin, account for 83.5%, 57.7%, 64.1%, and 68.8% of cattle, sheep, goats, and pigs, respectively. **Table 28** gives the population of major livestock in the Ghana side of the Basin.

Table 28: Population of Major Livestock in the Volta Basin in Ghana

Livestock	Population	% of National Total
Cattle	1,111,707	89.1
Sheep	1,672,395	69.1
Goats	1,854,749	70.5
Pigs	231,760	65.3
Poultry	5,479,352	37.5

Livestock is also taking a significant toll on soil productivity in the region. Although there are little data on the specific effects of livestock is scanty, it is clear that livestock negatively influences the area.

In Togo livestock production is not as developed as in Burkina and Ghana because of water and pasture scarcity and the presence of trypanosomiasis vector. **Table 29** shows national figures of the 1995 agricultural survey. Most of the livestock production is located within the Volta basin

Table 29: Population of Major Livestock in Togo

Livestock	Population
Cattle	202,340
Sheep	501,139
Goats	813,532
Pigs	330,750
Poultry	5,738,400

In Burkina Faso, animal husbandry, just like agriculture, occupies an important place in the farming system in all the regions of Burkina. It plays an important role in the nation's economy approximately 16 % of exports of goods and nearly 12.69 billion CFA in absolute value en 1989 (MARA, 1991). About stocks, there are: 4.260.900 heads of cattle, 5.680.600 sheep, 7.242.100 goats, 552.300 pigs, 445.300 asses, 23.032 horses, 13.056 camels and 18.776.400 fowls (DSAP/MARA, 1995).

iii. Fisheries

Fish production is also an important para-agricultural activity in the basin. In the Oti River in Benin, fishes are abundant. Downstream in Togo, fishing is also found to be a secondary activity for most of the population.

During the drought of the 1970s, fishing gained importance as an economic activity and as a source of food in Burkina Faso. Fishing is done in rivers (Mouhoun, Oti, Kompienga, Comoé, Léraba, Béli, Faga, Garoual, Sirba, and Tapoa) and in reservoirs (Kompienga, Bagré, Sourou, Moussodougou, Zega, Loumbila, and Kanazoé). Families of fish that are exploited include Cichlidae, Centropomidae, Mochokidae, Clariidae, Bagridae, Clatoteidae, Characidae, Mormyridae, and Osteoglossidae. Although this area has not been studied adequately, it is estimated that between 8,000 and 8,500 tons of fish are caught annually at a national level, and that this amount constitutes roughly 60-70% of biological capacity. Fish resources are being modified and threatened by overexploitation in certain areas due to the degradation of waterways, however.

Fish farming and fishing in Côte d'Ivoire experienced a takeoff in 1978 with a fishery development project. A provisioning center of 8 ponds with a capacity of 60,000 alevins per year and 52 fish farming ponds were established. The fish farming activity has strongly regressed since 1993, however, largely due to inadequate water resources and the slowdown in farming.

In Ghana, the Volta Lake created by the Akosombo Dam produced about 87,500 metric tons of fish in 2000. It is stated that the Volta Lake produces about 98 % of the inland fresh water fish in Ghana (Brammah, 2001). The increases in fish landings from the Volta Lake in the last half-decade are the result of deployment of active gear, such as the winch net, with unapproved mesh sizes in the lake. **Table 30** gives the annual fish production in Ghana between 1996 and 2000.

Table 30: Annual Fish Production in Ghana

Source	1996		1997		1998		1999		2000	
	No.	%	No.	%	No.	%	No.	%	No.	%
Marine	378,000	84	377,600	80	336,700	74.3	384,700	83.7	421,320	82.8
Inland	74,000	16	94,400	20	116,200	25.7	75,000	16.3	87,500	17.2
Total	452,000		472,000		452,900		459,700		508,820	

Source: SRID, MOFA, 2001

Prior to the construction of the dam on the Mouhoun in Burkina Faso, fishing in Mali was done using rudimentary equipment and the activity did not provide a significant source of income. When the waters rose, however, several villages moved to the edge of the river and fishing became a more important activity. The industry has since moved from subsistence to commercial.

iv. Forestry

Forests in the basin are cut to provide firewood and charcoal for local populations. In addition, forests are developed to provide lumber. For example, the forests in the Togo section of the basin provide more than half of the country's production of sawlog (Togo National Report). Forests in the region have been severely overexploited, however, and are threatened. The forested areas in the Volta basin of Burkina Faso have been reduced by 1.26 million hectares from 1980 to 1992. In 1996 alone, over 105,000 hectares in the Burkinabe portion of the Volta basin were deforested.

II.5 Water resources

Major sources of water in the Volta River system and riparian countries are natural rainfall, rivers, streams, lakes, groundwater and artificial impounded water (dams, dug-outs and reservoirs). The estimation of direct recharge to the system is based on the assumption that recharge occurs when actual evapotranspiration and direct run-off are balanced by precipitation. This occurs when the soil is saturated to the field capacity, which is likely to occur when precipitation exceeds evapotranspiration. Analyses of rainfall data from various stations within the Volta River system indicate that the months in which precipitation exceeds the evapo-transpiration are usually June, July, August, and September. The annual recharge for the Volta River system ranges from 13.4 % to 16.2 % of the mean annual precipitation. On average, the mean annual recharge of the Volta River system is about 14.8 % of the mean annual precipitation

II.5.1 Surface Water Resources in the Volta River System

i. GHANA

Naturally, rainfall is the single source of water that feeds all other sources. Whenever

rainfall is insufficient, recharge of water from other sources is low. In the extreme south-west of Ghana, the mean annual rainfall is over 2000 mm. Rainfall reduces eastward and northwards to about 800 mm in Accra (the capital) and 1000 mm in the north of the country (see Figure 3). A comparison of the total annual rainfall between 1960 and 1991 shows that there is no apparent decrease in rainfall in the northern and coastal parts of the country. The Ashanti and Brong-Ahafo regions may be getting slightly wetter.

Rivers constitute another important source of surface water supply in Ghana. There are 3 main river systems that drain the country. These are the Volta River System, the south-Western River system and the Coastal River System. These cover 70, 22 and 8 percent respectively of the total area of Ghana. The Volta River system consists of the Black, White and Main/Lower Volta and Oti and Daka Rivers. The southwestern rivers comprise the Bia, Tano, Ankobra and Pra rivers. The coastal rivers are made up of Chi-Nakwa, Ochi Amissah, Ayensu, Densu and the Tordzie rivers (MWH, 1998).

The yields from runoff in the various river basins generally follow the rainfall pattern. In June, July and October when rainfall is heaviest in the south of the country, flows are high. In northern Ghana, where rainfall is unimodal, high flows occur in August through October. The dry season extends from November to March and low flows are observed during this period throughout the country (FAO-RAF, 2000). The total annual runoff for the country is 54.4 billion m³ of which about 70 % is accounted for by the Volta River system.

All the tributaries of the Volta River enter Ghana and converge in the Lower Volta Basin. Rainfall in the basin varies from approximately 1000 mm to 1600 mm. The surface water resources received annually from outside and within the country are shown in Table 31.

The flows into the Lower Volta were based on specific yield of the catchment and may not be very accurate. Approximately 54% of the flows of the transboundary tributary originate from outside the country. An earlier estimation by Nathan Consortium (1970) puts this figure around 70 %. This may be explained by the reduction of rainfall magnitudes in the Sahel in the high latitudes of West Africa since the 1970s (Nicholson, 1983). Further, for the Oti River, approximately 76 % of the water resources originate from outside the country. The total mean annual flow of the entire Volta River system is estimated to be 38.3 billion m³ (MWH, 1998). The surface water resources of the Sub Rivers in the entire Volta river basin are described below.

Table 31: Surface Water Resources of the Volta River in Ghana

	River	Mean Flows (m ³ /s)	Mean Annual Flow (10 ⁹ m ³)
Water resources that originate from outside the country	White Volta	110.7	3.49
	Black Volta	103.75	3.27
	Oti	276.4	8.72
	SUB-TOTAL	490.85	15.49
Water resources	White Volta	192.57	6.08

from within the country	Black Volta	139.55	4.40
	Oti	89.1	2.81
	Lower Volta	289	9.12
	SUB-TOTAL	710.22	22.41
	TOTAL FLOW	1,201.07	37.90

Source: (Opoku-Ankomah, 1998)

- **The White Volta Basin**

The White Volta sub-basin covers about 49210 km² in Ghana, representing 46 % of its total catchment area of 10741.67 km² distributed in Ghana, Burkina and Togo. Its main tributaries are Morago and Tamne. The Morago has a total area of 1608 km² with an area of 596 km² in Ghana and 912 Km² in Togo. The Tamne lies entirely in Ghana with an area of 855 km². The White Volta covers mainly the north-central Ghana and some parts of the Upper and Northern Regions. It is located within the Interior Savanna Ecological Zone and is underlain by the Voltaian and granite geologic formations.

Annual rainfall in the sub-basin ranges between 1000 in the north and 1200 mm in the south; pan evaporation is about 2550 mm per year and runoff from within the basin averages about 96.5 mm per year. The average annual runoff from the White Volta is about 272 m³/s and the mean monthly runoff from within the basin varies from a maximum annual flow of 1216 m³/s to a minimum of about 0.11 m³/s. Potential storage sites have been identified within the basin totaling nearly 8180 10⁶ m³ which could regulate the basin yield at a minimum flow of about 209 m³/s. It contributes about 20 % of the annual total flows to the Volta Lake. Specific suspended sediment yield in this basin is between 8.5 and 14.0 tonnes/yr/km². Current surface water uses in the basin are estimated at about 0.11m³/s for domestic water supply and about 2 m³/s at numerous small irrigation projects.

Development potentials have been identified in the White Volta Basin which include a total of 63 megawatts of installed hydroelectric generating capacity, 155,809 hectares of irrigation, flood control, domestic water supply, navigation and recreation.

- **The Black Volta Basin**

The Black Volta has a total catchment area of 142,056 km² including areas outside Ghana. Only 33,302 km² (23.5%) of the catchment area are located in Ghana. Its main tributaries are Kamba, Kuon, Bekpong, Kule Dagare, Aruba, Pale, San, Gbalon, Chridi, Oyoko, Benchi, Chuco and Laboni. The catchment areas are all within Ghana. The Black Volta basin is primarily located in northern-western Ghana. The basin includes portions of the Upper, Northern and Brong Ahafo Regions.

Annual rainfall in the basin varies between about 1150 mm in the north and 1380 mm in the south; pan evaporation is on the order of 2540 mm per year, and runoff is about 88.9 mm per year.

The annual runoff from the Black Volta Basin is about 243 m³/s. Mean annual flow is about 8300 106 and the mean monthly runoff from the basin within Ghana varies from a maximum of about 623 m³/s to a minimum of about 2m³/sec. It contributes for about 18 % of the annual total flows to the Volta Lake. The potential storage site at Bui has a volume in excess of 12.3 10⁹m³ and could regulate the basin yield at a minimum of about 200 m³/s. The specific suspended sediment yield in this basin ranges from 8.0-12.0 tonnes/yr/km². Current surface water use is estimated to be only about 0.03 m³/s for domestic water supply.

The surface water resources of the Black Volta Basin consist of runoff from outside and within the country. The inflow into the country can be estimated from the discharges measured at Lawra, which is near the border. Similarly, the total discharge in the basin can be estimated from Bamboi, the southern most gauging station (See Table 32).

Table 32: Surface water flows of the Black Volta in Ghana

Station	Catchment area (k m ²)	Annual discharge (m ³ /s)	Dry season discharge (m ³ /s)	Wet season discharge (m ³ /s)
Lawra (inflow)	90,658	103.75	34.75	172.13
Bamboi	128,759	218.97	62.83	373.79
Catchment outlet (outflow)		243.30	69.81	415.32
Flow from within the catchment in Ghana		139.55	35.06	243.19
% Inflow/total Outflow		42.64	49.7	41.45

Development potentials have been identified in the basin which include 230 megawatts of installed hydroelectric generating capacity, about 38,446 hectares of irrigation, flood control, and recreation.

- **The Lower Volta**

The Lower basin is located below the confluence of the Black Volta and the White Volta rivers, excluding the Oti river drainage area. The surface water resources in the basin consist of flows from outside the country and flows from within the country. Discharges of White Volta at Nawuni and Mole River at Lankatere were used to estimate the total basin discharge (Table 33).

Table 33: Surface Water flows of the Lower Volta of Ghana

Station	River	Catchment area	Annual discharge (m ³ /s)	Dry season discharge (m ³ /s)	Wet season discharge (m ³ /s)
Nangodi	Red Volta	10,974	30.72	0.34	61.12
Yarugu	White Volta	41,619	80.00	2.17	157.00
Total inflow			110.72	2.51	218.12
Nawuni	White Volta	96,957	229.98	18.95	440.05
Lankatere	Mole		73.31	15.78	131.33
Total outflow	White Volta		303.29	34.73	571.38
Total flow from within the catchment area in			192.57	32.22	353.26

Ghana					
% Total inflow/total outflow			36.5	7.2	38.0

The Lower Volta Basin covers a total area of about 68588 km² and most of that (50 432 km²) is located in east-central Ghana. The basin includes also portions of the Northern, Brong Ahafo, Volta, Ashanti, Eastern Regions and parts of Togo.

Annual rainfall in the basin varies from about 1100mm in the northern part of the basin to about 1,500mm in the central, and to about 900mm in the southern part. Pan evaporation is about 1,800mm per year and runoff from within the basin is estimated to be about 89 mm per year. The natural total mean runoff from the Basin is estimated to be about 1,160 m³/s; the Volta Lake behind Akosombo dam providing extensive regulation. Current river water withdrawals in the basin include about 1.86 m³/s domestic water supplies, about 0.71 m³/s for irrigation water supply, and about 566 m³/s for power. In the future, nearly all the regulated outflow from Akosombo will be used for power generation. **Table 34** gives the estimated annual flow of the tributaries of the Main Volta River basin

Table 34: Sub-basins of the Main Volta River in Ghana

Sub-basin	Area (km ²)	Specific yield (m ³ /s/km ²)* 1000	Estimated annual basin flow (m ³ /s)
Daka	8,283	7.996	66.2
Kularakum	5,931	8.000	47.4
Pru	8,728	2.176	19.0
Sene	5,366	2.176	11.7
Obosom	3,620	2.176	7.9
Dayi	1,828	8.289	15.7
Asukawkaw	2,233	8.081	18.0
Alabo	1,023	3.086	3.2
Afram	11,396	8.766	99.9
Total flow from the tributaries	48,478		289.0

The surface water resources from within the basin were estimated from the annual flows of these primary tributaries. Summary of flows from outside the country, within the country but outside the Lower Basin and within the Lower Basin are shown in **Table 35** below:

Table 35: Surface water resources from within the Lower Volta Basin in Ghana

	River	Study (m ³ /s)	Nathan Consortium (m ³ /s)
Flows that originate from outside the country	White Volta	10.7	152.06
	Black Volta	103.75	186.32
	Oti	276.40	389.34
	Sub-total	490.85	727.72
Flows from within the country but outside the Lower Volta	White Volta	192.57	118.65
	Black Volta	139.55	57.20
	Oti	89.10	111.85

	Sub-total	421.22	287.70
Total inflows into the Lower Volta Basin		912.07	1015.44
Flows from within the Lower Volta	Total	289.0	140.73

The total inflow into the Lower Volta Basin is 912 m³/s compared to 1015 m³/s estimated earlier by Nathan Consortium. Thus, the percentage of current flows into the Lower Volta basin is about 90 % of the amount estimated in the earlier report. The observation here is that the flows that originate from outside the country into the Volta Basin have considerably reduced while the flows in the country have increased by 46%.

Development potentials identified in the basin include a total of about 202,350 hectares, 140 megawatts of additional hydropower generating capacity exclusive of potential additional installations at Akosombo, flood control, navigation, recreation and water supplies for domestic, municipal and industrial uses.

- **Oti Basin**

The Oti River Basin has a surface area of 16801 km² in north eastern Ghana. The basin includes portions of the Northern and Volta Regions. It also expands to Togo where it covers more than 40 % of the land. The relief varies considerably from 150 m to 450 m in Ghana and much more in Togo. Annual rainfall in the basin varies from 1010 mm in the north to 1400 mm in the south; pan evaporation is about 2540 mm per year and runoff is about 254 mm per year.

The average annual runoff from the Oti basin is about 500 m³/s and the mean monthly runoff from the basin within Ghana is from 849 m³/s to 1.1 m³/s. The mean annual flow is estimated at 12, 606 x 10⁶ m³. The Oti River Basin contributes about 25 % of the annual total flows to the Volta Lake due to the steep topography and the relatively high rainfall within the basin. Potential storage sites have been identified within the basin, totaling about 406 million m³, which could regulate the basin yield at a minimum flow of about 37 m³/s. Specific suspended sediment yield in this basin is 27.7 tonnes/yr/km². It is the highest in the Volta River system of Ghana

The Oti River Basin is entirely within the Interior Savanna Ecological Zone and is underlain by the Voltaian and Buem geologic formations. Current surface water usage in the basin is negligible and there have been no irrigation development potentials identified in the basin.

- **Daka basin**

The Daka Basin covers an area of 7,424 km² located almost entirely in eastern part of northern Ghana. The annual rainfall on the basin varies between 1120 mm and 1340 mm and most of the biophysical characteristics found in the basin are similar to those of the Oti River Basin.

Another important source of surface water supply in Ghana is from surface impoundments and dugouts. The largest reservoir in the country is the man-made Volta Lake created in 1964 at Akosombo to generate hydro-electric power. The dam is 88.2 m high with a crest length of 638.4 m and a generating capacity of 833 MW. It has a mean annual inflow of 4.05 ha/m and a live storage of 6.03 ha/m (FAO-RAF, 2000/1)

ii. BURKINA FASO

Burkina Faso has a total mean annual precipitation volume of 165 km³, of which only 9 km³ is runoff. The country has eight large dams in operation, of about 2100 dams in total. The total storage volume of all reservoirs is approximately 4.6 km³

The hydrographic network of the country is characterised by three great river basins: the basins of the rivers Nazinon, Nazinga and Mouhoun, of the River Comoé and the River Niger.



Figure 13: Hydrographic network of Burkina Faso

The rivers Mouhoun (Black Volta) and Nakanbé (White Volta) constitute the two main sub-basins of the Volta Basin in Burkina Faso. The country has other smaller rivers – the Bougouriba, Comoé, Béli, Sirba and Tapoa. Many of these rivers dry up in the dry season (October-June), with the exception of the Mouhoun and Comoé in the south-west which are fed by springs (Bandre *et al.*, 1998).

Mean annual rainfall in the Mouhoun and Nakanbé sub-basins varies from approximately 900 mm to less than 600 mm. Over the past 40 years, the precipitation patterns have been of increased dryness, especially between 1970 and 1980. While rainfall increased between 1985 and 1995, the last decades are still marked by the following trends:

decrease in river flow, decreased availability of groundwater, the drying up of source waters, and degradation of vegetation cover.

The surface water resources of the basin are made up of stream flows and water in reservoirs. **Table 36** illustrates the distribution of the available surface water resources in the Volta River Basin of Burkina Faso.

Table 36: Potential Surface Water Resources of the Volta Basin in Burkina Faso

Sub-basin	Annual Flow Volume (x 10 ⁹ m ³)	Volume in Reservoirs (x 10 ⁹ m ³)	Potential in the Basin (x 10 ⁹ m ³)
Mouhoun (Black Volta)	2.64	0.29	2.75
Nakambé (White Volta)	2.44	2.20	3.32
Total	5.08	2.49	6.07

Source: Etat des lieux des ressources en eau et de leur cadre de gestion

The Nankambe River is one of the main tributaries of the Volta River. Its basin covers about 81,932 km² located in the following four sub-basins:

- Sissili basin 7,559 km²
- Nazinon basin 11,370 km²
- Pendjari basin 21,595 km²
- Nouhoa basin 4,050 km²

The first intermittent flows in the Nakambe occur in May. They become permanent flows in July and August at the Wayen station (20,800 km² watershed) and larger downstream when they reach Bagre (33,120 km²) where there are average values of 65.4 m³/s in July, 145 m³/s in August and 107 m³/s in September. During the three months, 88% of the annual flows at Bagre (29.7m³/s representing 946 Mm³) occur.

The Nazinon and the Sissili Rivers are two of the Nankambe tributaries that collect runoff water from the South-West part of the Mossi plateau. The Pendjari River which forms the border with Benin has three main tributaries: the Doudodo, the Singou and the Kompienga Rivers. These rivers flow during the rainy season (750 -900 mm) and dry up in November.

Annual surface water resources from the basin are shown in the following **Table 37**.

Table 37: Potential of Surface Water Resources of the Nakambe Basin

Sub-basin	Mean flow (m ³ /s)	Contribution of the basin (10 ⁶ m ³)
Nakambe	33.4	1,054
Nazinon	6.04	160
Sissili	2.11	67
Pendjari	28.5	899
Nouhao	7.59	239
Total Nakambe Basin in Burkina	77.64	2,444

Existing reservoirs in the Nakambe basin have a total storage capacity of 4.3 billion m³ including the Ziga reservoir. About 27 of the main reservoirs in the basin are frequently monitored and the total volume stored in these reservoirs is estimated at 2.04 billion m³ which represent half of their capacity. Bagre and Kompienga are the largest reservoirs in the Nakambe basin and are used for power generating, irrigation and environmental purposes.

The Mouhoun River is the main tributary of the Volta River. Its basin covers 91,036 km² and can be divided into three sub-basins:

- Upper Mouhoun basin 20,978 km²
- Sourou basin 15,256 km²
- Lower Mouhoun basin 54,802 km²

The upstream portion of the Mouhoun has perennial flows with base flows rarely below 2 m³/s at the Samendeni station or at Nasso on the Kou River. At the confluence with the Sourou River, the Mouhoun and its main tributaries (Plandi, Kou, Voun Hou) produce a mean flow of about 25 m³/s. However this flow is irregular. With the construction of a water abstraction structure on the Sourou River in 1984 and the recurrent precipitation deficits, the natural regime has been changed during the base flow and flood periods. Thus, the estimated average base flow of 5.9m³/sec at Boromo (ORSTOM, 1977) has since decreased to a dry up of the river in 1984. Similar observations have been made at the Dalopa and Noubel stations where average flows were estimated at 7.35 and 9.3m³/s respectively.

Table 38: Potential of surface Water Resources of the Mouhoun Basin in Burkina Faso

Sub-basin	Mean flow (m ³ /s)	Contribution of the basin (10 ⁶ m ³)
Mouhoun at Ouessa	43.0	1,356
Sub-basin Bougouriba	28.0	884
Su-basin Bambassou	12.9	405
Total Mouhoun Basin in Burkina	83.9	2,646

Existing reservoirs in the Mouhoun basin have an overall capacity of 438x10⁶ m³. A planned reservoir at Samandeni on the Mouhoun, 30 km north-west of Bobo-Dioulasso will alone hold about 500x10⁶ m³. Only the Sourou reservoir at Yaran with a capacity of 250x10⁶ m³ is currently monitored by hydrologists. This reservoir and those on the Nakambe River have never reached their capacity. It is estimated that existing reservoirs in the basin store about 300x10⁶ m³. Small reservoirs however, get full during normal years.




Figure 14: Location off reservoirs in the Volta Basin




Figure 15a: Water balance in the Mohoun River Basin




Figure 15b: Water balance in the Nankambe River Basin

iii. MALI

Though very little data are available on water resources in Mali, the country possesses substantial surface water resources made up of: (1) rainwater, ranging from 1,400 mm per year in the south to less than 150 mm per year in the north; (2) perennial surface water, with an annual flow of about 56.5 billion cubic meters, divided between the Niger River at Koulikoro (46 billion) and the Senegal River at Kayes (10.5 billion); and (3) non-perennial surface water like the Sourou River (the Volta Basin of Mali)

In the Volta Basin region of Mali, water has been in such short supply that the first priority has been to provide water for the inhabitants, while research was overlooked. The Sourou River is the main source of surface water in the region. Annual rainfall is approximately 400 mm and surface flows are only ephemeral as streams dry up after 3 to 5 months of the rainfall season. Discharge measurements are limited and available data could not allow for quantitative assessment of surface water resources. However, about 52% of the villages in the region depend on surface water (i.e., streams, lakes, ponds, etc.) for short periods.

The water level of the Sourou River in Mali increased significantly after a dam was constructed in Burkina Faso in 1989. Sourou valley now forms the northern end of the reservoir created by the dam.

The only other sources of surface water are small temporary ponds that appear in Seno during the winter. There are 12 ponds in the southern zone (Dioura) and 9 others in the Bankass District and these dry up after 3 to 5 months.

The chemical and bacteriological quality of the surface water is generally bad, largely due to faecal contamination. The polluted water is a source of water-borne diseases.

iv. TOGO

The northern section of the Oti basin of Togo receives between 1000 and 1200 mm of rainfall annually, while the southwest region receives from 1000 to 1500 mm per year. Surface water resources estimated for the basin are about $4.71 \times 10^9 \text{m}^3$ per year. Most streams dry up during the dry seasons due to high evapo-transpiration. In the northern part of the basin, the Oti, enlarged by its tributaries and Mò exceed $100 \text{m}^3/\text{s}$ in the Savannah region and 100 to $300 \text{m}^3/\text{s}$ in the Kara area. The extreme variability of the flows between the wet and dry seasons makes dependency on the surface water for irrigation difficult. In the southwestern section of the basin, the Menou, Wawa, and Danyi have much smaller flows of between 1 and $6 \text{m}^3/\text{s}$, but these are perennial flows as the climate is wetter.

Water shortage in Togais projected to be exacerbated by the effects of climate change. It is estimated that by 2025, average monthly temperatures will rise from South to North at 0.48 to 0.58% , which is 0.8 to 1°C over 1995 levels. Precipitation is expected to decrease 0.1 to 0.3% .

v. CÔTE D'IVOIRE

Lack of hydrometeorological, hydrogeologic, hydroclimatic, water quality, and sediment transport data for the Black Volta Basin of Côte D'ivoire makes it difficult to accurately evaluate the surface and subsurface waters in the Côte D'ivoire area. Average annual rainfall over the basin is approximately $1,000$ mm. The surface water resources derived from the Black Volta are about $0.788 \times 10^9 \text{m}^3/\text{yr}$. There are 43 dams established in the basin, with a storage capacity of 3 million m^3 .

vi. BENIN

The average annual rainfall in the Oti River Basin of Benin is approximately $1,100$ mm. In normal years, flows are about $58.6 \text{m}^3/\text{s}$ and the annual flow volume is estimated at approximately $1.85 \times 10^9 \text{m}^3$. Estimation of groundwater resources in the Oti Basin is difficult since these resources are almost inseparable from the groundwater resources of the Niger basin in Benin.

Benin has a hydroelectric power station on the Oti River with a storage capacity of 350 million m^3 and the capacity to produce 15 MW. Additionally, a hydroelectric power station is planned at Pouya (Natitingou) on the Yéripao.

II.5.2 Groundwater Resources, Recharge, and Quality

The geological characteristics of the basin show that the rocks have no inherent porosity. Formation of aquifers, therefore, depends upon secondary porosity created as a result of fissuring or weathering. Muscovite or hornblende can weather to approximately 30 m, whereas the Birimian formation can weather to a depth of approximately 73 m, thus

giving rise to a thicker aquifer. The hydrogeological characteristics of the basin in Ghana are described in **Table 39** below:

Table 39: Hydrological Characteristics of the Groundwater aquifers in the Volta Basin of Ghana

	Run-off Coefficient (%)	Borehole Yields (m ³ /h)	Mean Borehole Yields (m ³ /h)	Specific Capacities (m ³ /h/m)	Depths to Aquifer (m)	Mean Depth to Aquifer (m)	Depth of Boreholes (m)	Mean Depth of Borehole (m)
White Volta	10.8	0.03 – 24.0	2.1	0.01 – 21.1	3.7 – 51.5	18.4	7.4 – 123.4	24.7
Black Volta	8.3	0.1 – 36.0	2.2	0.02 – 5.28	4.3 – 82.5	20.6		
Oti	14.8	0.6 – 36.0	5.2	0.06 – 10.45	6.0 – 39.0	20.6	25.0 – 82.0	32.9
Lower Volta	17.0	0.02 – 36.0	5.7	0.05 – 2.99	3.0 – 55.0	22.7	21 – 129.0	44.5

Source: MWH (1998)

The Table indicates that run-off coefficients are in general low. Direct recharge of aquifers from precipitation is less than 20% across the basin. These figures do not give a good outlook for the recharge of the groundwater resources.

Borehole yields are quite variable with a mean for all the sub-basins between 2.1 and 5.7 m³/h. These figures suggest that the groundwater yields in the basin are low. The figures in the table show that the region has low hydraulic transmissivity.

The depth of aquifers is also variable in the basin. Studies have shown that there is no correlation between depths to aquifer and borehole yields. The results indicate that groundwater resources are not abundant in the basin and face threats if not properly managed (UNEP, 2002).

i. GHANA

(a) The Hydrogeology

Ghana has two major hydrogeologic provinces. These are: (1) the basement Complex composed of Precambrian crystalline igneous and metamorphic rocks, and (2) Paleozoic consolidated sedimentary formations (Figure 10). Minor provinces consist of (1) Cenozoic, Mesozoic, and Paleozoic sedimentary strata along narrow belts on the coast; and (2) Quaternary alluvium along the major stream courses.

The basement complex underlies about 54% of the country and is further divided into sub-provinces on the basis of geologic and groundwater conditions. Generally, these sub-provinces include the metamorphosed and folded rocks of the Birimian system, Dahomeyan system, Tarkwaian system, Togo Series, and the Buem Formation (Figure

17). The basement complex consists mainly of gneiss, phyllite, schist, mihnatite, granite-gneiss, and quartzite.

The Paleozoic consolidated sedimentary formations, locally referred to as the Voltaian Formation, underlie about 45% of the country and consist mainly of sandstone, shale, arkose, mudstone, sandy and pebbly beds, and limestone. The Voltaian Formation is further subdivided on the basis of lithology and field relationships into the following sub-provinces: (1) Upper Voltaian (massive sandstone and thin-bedded sandstone); (2) Middle Voltaian (Obusum and Oti Beds); and (3) Lower Voltaian (**Figure 18**).

The remaining 1% of the rock formation is made up of two coastal provinces (the coastal Block-Fault Province and the coastal-Plain Province) and the Alluvial Province. The coastal Block-Fault province consists of a narrow discontinuous belt of Devonian and Jurassic sedimentary rocks that have been broken into numerous fault blocks and are transacted by minor intrusion. The coastal plain hydrogeologic province is underlain by semi-consolidated to unconsolidated sediments ranging from Cretaceous to Holocene in age in south eastern Ghana and in a relatively small isolated area in the extreme south western part of the country. The Alluvia hydrogeologic province includes narrow bands of alluvium of Quaternary age, occurring mainly adjacent to the Volta River and its major tributaries and in the Volta delta (Ministry of Works and Housing, 1998; Dapaah-Siakwan and Gyau-Boakye, 2000).




Figure 16: Hydrogeological provinces and river systems of Ghana (Geological Survey of Ghana 1969)




Figure 17: Hydrogeological subprovinces of the Basement Complex (Ghana Geological Survey 1969)




Figure 18: Hydrogeological subprovinces of the Voltaian System (Ghana Geological Survey 1965)

(b) Aquifers found in Ghana

The rocks that underlie 99% of Ghana (the basement complex and the Voltaian formation) are essentially impermeable and have little or no primary porosity. Therefore groundwater occurrence in Ghana is associated with the development of secondary porosity as a result of jointing, shearing, fracturing and weathering. This has given rise to two main types of aquifers: the weathered zone aquifers and the fractured zone aquifers. The weathered zone aquifers usually occur at the base of the thick weathered layer. The weathered layers vary, from 0 m (outcrops) to about 100 m. The weathered layer is thickest in the wet forested south-western part of the country where it reaches an average thickness of 60 m and is thinnest in the semi-arid zone in the extreme northeast where the mean thickness is 10 m. The fractured zone aquifers are normally discontinuous and limited in area. Due to the sandy clay nature of the weathered overburden, the groundwater occurs mostly under semi-confined or leaky conditions. The yield of these aquifers rarely exceeds 6 m³/h (Ministry of Works and Housing, 1998).

Three aquifers occur in the remaining 1% of Ghana, mainly in the extreme south eastern and western part (with cenozoic and mesozoic sediments formation). The first aquifer is unconfined and occurs in the recent sand very close to the coast. It is between 2 m and 4 m deep and contains fresh meteoric water. The intermediate aquifer is either semi-confined or confined and occurs mainly in the red continental deposits of sandy clays and gravels. The depth of this aquifer varies from 6 m to 120 m, and it contains mostly saline water. The third aquifer is the limestone aquifer. It varies in depth between 120 m and 300 m. The groundwater in this aquifer, which occurs under artesian condition, is fresh. The average yield of the limestone aquifer is about 148 m³/h (Ministry of Works and Housing, 1998; Dapaah-Siakwan and Gyau-Boakye, 2000).

(c) Borehole Yields

Yields from boreholes are highly variable because of the lithological varieties and structural complexities of the rocks. In 1994, the Water Resources Research Institute analysed borehole yields for the various geologic formations in the country. The least explored geologic unit is the Voltanian system (underlying also the Volta basin). Table 40 gives a summarizes borehole yields for the various hydrologic units in the country.

Table 40: Summary of borehole yields of hydrologic provinces and sub provinces

Hydrogeologic province and subprovince	Borehole-completion success rate (%)	Range of yield (m ³ /h)	Average Yield (m ³ /h)
Basement Complex			
Lower Birimian System	75	0.41-29.8	12.7
Upper Birimian System	76.5	0.45-23.6	7.4

Dahomeyan System	36	1-3	2.7
Tarkwaian System	83	1-23.2	8.7
Togo Series	87.9	0.72-24.3	9.2
Buem Formation	87.9	0.72-24.3	9.2
Voltaian System			
Lower Voltaian	55	1-9	8.5
Middle Voltaian (Obusum and Oti beds)	56	0.41-9	6.2
Upper Voltaian	56	1-9	8.5
Cenozoic, Mesozoic, and Palozoic			
Sedimentary Strata (Costal Provinces)			
Coastal Block-Fault Province	36	1-5	3.9
Coastal-Plain Province	78	4.5-54	15.6
Alluvial Province	67	1-15	11.7

Source: Dapaah-Siakwan and Gyau-Boakye (2000)

Again, in 1994, the Water Resources Research Institute prepared a borehole-yield map of Ghana based on available data on borehole yields; static water level and other vital information (**Figure 19**). This map indicates the borehole yield to be expected in any area within the country.




Figure 19: Distribution of borehole yield in Ghana (Water Resources Research Institute 1994)

(d) Groundwater abstraction and distribution

Groundwater is abstracted from all the geological formations in the country. In 1994 there were over 45,000 abstracting systems made up of approximately 10,500 boreholes, 45,000 hand-dug wells and some dug out, all over the country (Kortatsi, 1994). Current available information shows increase in the number of abstraction systems, possibly due to increase in population, which has resulted in a higher demand for water for various uses particularly domestic. As at march 1998, the number of hand dug wells had risen to about 60,000 while the number of boreholes reached 11,500 in the year 2000 (Dapaah-Siakwan and Gyau-Boakye, 2000); making a total of over 71,500 systems. From the borehole and well figures, it could be inferred that the rate of construction of wells in Ghana (1994-1998), and that of boreholes (1994-2000) was 10 per day and 1 every other day respectively. A hand dug well is a cost-effective device for extracting shallow groundwater bodies and it is a technology that has found extensive use as a traditional water supply system in many rural and urban communities throughout Ghana.

A typical hand dug-well in Ghana consists of three components, the intake, a shaft and the wellhead (DANIDA, 1993). Boreholes are also found in use in several areas in Ghana though the cost involved limits its use (the average cost of drilling a borehole in Ghana, including pump testing is about US \$3,920). Most boreholes have been drilled through one project or the other for community use. Few private organizations and very few individuals own their own boreholes.

There is limited data on hand-dug wells in the country, except the Volta region where an inventory has been done and from this inventory, it can be inferred that hand dug well yield varies from 0 (dry well) to 26 m³/day with a mean of 6 m³/day (Kortatsi, 1994). Therefore, the estimated total abstraction of hand dug wells per year is 1.3x10⁸ m³. Figure 20 gives the regional distribution of borehole (NB) and the estimated annual abstraction (AA) of groundwater based on 12 h of pumping per day. The estimated total annual abstraction of boreholes is 1.41x10⁸ m³.




Figure 20: Regional distribution of borehole and annual abstraction

(e) Aquifer recharge

Little information is available on groundwater recharge in Ghana. Recharge to all the aquifer systems in Ghana is mainly by direct infiltration of precipitation through fracture and fault zones along the highland fronts and also through the sandy portions of the weathered zone. Some amount of recharge also occurs through seepage from ephemeral stream channels during the rainy seasons. Some indirect recharge mainly occurs in the lower rainfall, low relief and low permeability areas. This happens when runoff from watershed outside the areas or a particular storm event is of sufficient magnitude to cause runoff. The drainage courses or stream which act as conduit for the overland flows are generally weak fissured zones which allow a greater part of the runoff to infiltrate through their beds to the groundwater table.

Data on water level fluctuation are scarce but support the contention of high recharge in some areas; observations carried out in the Upper Regions between 1976 and 1979 show oscillation of 0.3 to 5.4 m between the dry and wet seasons with the peaks normally in September/October. Wells monitored by the Water Resources Research Institute between 1980 and 1989 show generally irregular movements of groundwater levels and may have been affected by pumping either of the monitoring well itself or

other wells in the vicinity. Nevertheless, there are some indications that the groundwater system is active rather than passive, and is affected by significant recharge and discharge on an annual cycle (World Bank Country Report, 1992).

The minimum recharge for the sub-basins within the Volta Basin of Ghana and replenishable groundwater capacities are presented in Table 41.

Table 41: Minimum Recharge and Replenishable Groundwater Capacities

Sub-Basin	Minimum Recharge (mm)	Replenishable Groundwater Capacities (x 10 ⁹ m ³)
White Volta	151	6.6
Black Volta	205	3.4
Oti	175	3.7
Lower Volta	205	8.2

(f) Groundwater quality

Previous studies (Nathan Consortium studies, 1970; Amuzu, 1978; Andah, 1993; Kortatsi, 1994; Ministry of Works and Housing, 1998; Darko *et al*, 2003) revealed that the quality of groundwater in Ghana is generally good for multi-purpose use except for the presence of low pH (3.5-6.0) waters, high level of iron, manganese and fluoride in certain localities as well as high mineralization with TDS in the range 2000-14,584 mg/l in some coastal aquifers particularly in the Accra plains. In Tamale and Atiave, fluoride concentration levels could be as high as 5.0 mg/l and 20.0 mg/l respectively. About 30% of all boreholes in Ghana have iron problems (Ministry of Works and Housing, 1998). High iron concentration in the range 1-64 mg/l have been observed in boreholes in all geological formations. This iron originates partly from the attack of low pH waters on corrosive pump parts and partly from the aquifers (Ministry of Works and Housing, 1998). The percentage of iron derived from the aquifers is however unknown. **Table 42** gives the mean values of chemical analyses of many water samples in the various geologic formations in Ghana.

The waters in many hand-dug wells look turbid and polluted as they contain high levels of nitrate in the range of 30-60 mg/l and abundant coliform (Kortatsi, 1994). This is probably due to improper construction and inadequate protection of wells sites from surface runoff and animal droppings.

Table 42: Chemical analysis of water samples in the geologic formations of Ghana (all values except pH are in mg/l)

	Gneiss	Granitic formation	Phyllites	sandstone	Mudstone and shale	Sand and gravel	Limestone	Quartzite
pH	7.5	6.99	6.83	6.95	7.64	7.53	7.7	6.36
Total dissolved salts	4888	387.38	211.19	533.45	424.66	632.04	932.04	398.26

Calcium (Ca)	595	49.38	32.09	25.08	26.10	68.72	58.08	42.06
Magnesium (Mg)	207.2	19.06	15.67	7.57	9.12	33.50	36.14	23.37
Sodium (Na)	720	47.99	11.67	262.55	125.39	134.45	296.77	24.53
Chloride (Cl)	1790	73.48	9.90	70.42	42.04	173.56	196.86	103.61
Sulphate (SO ₄)	1800	10.60	7.16	65.17	11.18	101.19	77.25	60.06
Bicarbonate (CO ₃)	34	81.17	104.14	97.49	189.29	154.59	149.66	67.05
Total Iron (Fe)	0.1	1.01	2.15	1.95	0.645	1.84	0.467	2.87
Maganese (Mn)	0.05	0.44	0.39	0.17	0.10	0.22	0.16	0.45
Flouride (F)	0.25	0.35	0.315	0.775	0.57	0.60	1.76	0.23
Nitrate nitrogen (NO ₃)	0.5	1.605	0.59	0.75	0.135	2.22	1.79	2.32
Total hardness	2340	172.49	123.70	70.76	222.77	230.35	229.94	179.61

Source: Kortatsi (1994)

ii. Burkina Faso

(a) Aquifers in Burkina Faso

The types of aquifers found in Burkina Faso can also be classified as continuous and discontinuous. The continuous aquifers are the continental terminal aquifers located in the Southern part of the Gondo plain and the weathered zone aquifers found in Bobo Dioulasso and Tenkodogou. The discontinuous aquifers are the Precambrian bedrock, the Gourma formation (y doubam group) and the Gres premier and intracambrain.

The top part of the continental aquifer is mainly made up of clay, sand, and *gres*. The bedrock consists of fractured and weathered schists and dolomite. The depth of the water table in this type of aquifer is between 10 and 80 m. Area covered by this aquifer is about 11,000 Km² with a recharge of 430 million m³ (equivalent to annual infiltration of 38 mm). The saturated depth of the aquifer decreases from southwest to northeast, from about 50 to 5 m.

Assuming a draw down of 1/3rd of the saturated depth, it is estimated that about 1-3 billion m³ of water can be extracted from this aquifer. Generally, the quality of water from this aquifer is very good, but it becomes poorer as one approaches the middle of the Gondo plain. In the Northern part of Bobo Dioulasso, this aquifer has a depth of 10-30 m and is mainly used with shallow wells. Not much is known about its recharge in Bobo area but most recharge is done through preferential flow. Water from this aquifer is also used for industrial and irrigation purposes though there was no information on the volume involved (CIEH, 1976; BILAN D'EAU, 1993).

The weathered zone aquifer is found in the Bobo Dioulasso region (where it is part of the "zone de socle" aquifer system), Tenkodogou region and other areas of Burkina Faso. In

general, this aquifer has a depth of 10-50 m. Many studies have shown that the depth is a determining factor in the success of landing water. That is, borehole or well success rate increases with depth. In Bobo Dioulasso, this aquifer has 10-30 m of weathered layer which gives an average yield of 0.5-5 m³/hr with transmissivities between 1.9-5.5x10⁻⁴ m²/s. The piezometric head is between 10 m and above the bedrock. Some 83 % of the flow in this aquifer is found in up to 50 m depth; 93 % within 60 m; and 90 % within 70 m. Optimum depth is between 45-65 m (CIEH, 1976; BILAN D'EAN, 1993).

The Precambrian bedrock type of aquifer consists of crystalline, volcanic and metamorphic rocks; granite, schist, green rock, etc. Availability of water resources in this aquifer is linked to the fracturing or weathering of these rocks. The weathering of schists produces clay which makes it impervious, thus water can only be found by drilling beyond these layers to the fractured layers. The coverage area of this aquifer is 225,000 Km² and recharge is estimated at between 3-4x10⁹m³ (equivalent to annual infiltration of 17 mm) (CIEH, 1976; BILAN D'EAN, 1993).

The Gourma formation (y doubam group) type of aquifer is characterised by dolomite and calcareous rocks that contain water in fractured zones. Recharge by rainfall is very low. Much is not known about this aquifer and needs research to understand its characteristics.

Gres premier and intracambrian aquifer is not well known but is important. It has a good yield and is located in areas with many perennial rivers in the Bobo Dioulasso area. It covers an area of 30,000 Km² and has a discharge of 1.9 billion m³ (equivalent to annual infiltration of 60 mm). It has a transmissivity of 1.4-4.8x10⁻³ m²/s and a yield of 1 m³/hr/m. The thickness of the aquifer in this area is estimated at 100 m. In general, discharges increases with depth. Research is needed to know more about this important aquifer (CIEH, 1976; BILAN D'EAU, 1993).

(b) Groundwater availability

Given the climatic conditions of Burkina Faso, surface water is in limited supply and groundwater is therefore an important resource. Rural water supply projects rely mainly on groundwater, although this is also scarce in many areas. Dammed river courses typically provide urban supplies. Only around 25% of the population have access to safe drinking water. Traditional sources of water are hand-dug wells and ponds used during the rainy season. Currently, groundwater is also abstracted from a number of tubewells, typically equipped with handpumps. Tubewells generally source groundwater from fractures within the basement complex, while hand-dug wells are largely sourced from the weathered overburden layer. Yields from tubewells and dug wells in the basement areas are generally low and many dry up seasonally (UN, 1983).

Groen et al. (1988) noted groundwater levels in north-western Burkina Faso of some 10–60 m below surface with a seasonal variation of around 1–2 m. In the sedimentary rock formations along the northwest border region, groundwater availability is also limited, although the dolomitic limestones form the best aquifers. The younger sedimentary

formations cropping out in the north constitute a small aquifer which is connected with underlying dolomitic limestone.

Based on many drilling works carried out in Burkina Faso, it was possible to classify areas of the country according to their groundwater potential. From this, it is noted that about 45 % of the territory of Burkina is in the category described as bad or poor potential (Table 43)

Table 43 : Groundwater potentiality of Burkina Faso

Groundwater potential	Surface area (Km ²)	Proportion (% territory)
Bad	74,000	27
Poor	52,000	19
Good	112,000	41
Very good	36,000	13
Total	274,000	100

Source: BILAN D'EAN (1993)

(c) Aquifer yield

Geology largely determines the yield of aquifers. Thus in the crystalline basement zones, which constitutes more than 80 % of Burkina, yields obtained in drillings are about 2 m³/h. Consequently in these zones, the large needs (urban water supply, industries and irrigation) are difficult to satisfy. On the other hand, in the sedimentary zones, significant yields are frequently obtained, sometimes more than 100 m³/h Figure 21 shows discharges of wells drilled in the entire country.




Figure 21: Map showing discharge of wells across Burkina Faso

In an IDRC project, 2 out of 16 (test-) drillings with depths to 80 m were dry, whereas the yield of 14 wells was between 0.6 and 6 m³/h. Success rate of drilling varies but usually it is quite low (70% for wells with a flow from 0.5 to 5 cubic metres per hour; 15% for wells with a flow of more than 5 cubic metres per hour and for those with a higher discharge is only 2.5%). Low success rates have been attributed partly to that drilling is local and lacks modern exploration methods (Faruqui, 2003).

Figure 22 shows the distribution of piezometric heads of different wells drilled all over Burkina Faso.




Figure 22: Distribution of Piezometric heads of different wells in Burkina Faso

(d) Recharge Mechanisms

Two mechanisms of recharge of the groundwater table have been identified in Burkina Faso. These are the direct and indirect recharge mechanisms. With the direct recharge, rainwater infiltrates directly into the ground and percolates towards the water table. With the indirect recharge, infiltration takes place in the low points (hollows, streams, alluvial valleys) where water concentrates after runoff. The environment plays a great role in infiltration. Indirect infiltration is more significant in the areas affected by the impoverishment of the soils where the recharge of the water tables occurs primarily around the hollows. The annual recharge of the water table is estimated at 5 mm in the North and 50 mm in the South. Depending on the permeability of the soil, this recharge varies between 0.1 and 10 % of annual rainfall. For the whole of Burkina Faso, recharge is estimated at 9.5 billion m³ per annum (FAO, 1995).

(e) Groundwater Abstraction and Distribution

The groundwater resources available in Burkina Faso are mainly provided by boreholes. In all, there are about 24,350 boreholes spread across the entire country. **Figure 23** shows the spatial distribution of boreholes in the various Provinces in Burkina Faso.

- **Nitrogen species**

Shallow groundwater has significant potential for being contaminated by pollutants from domestic (latrines, drains, waste tips) agricultural and industrial sources. Groen *et al.* (1988) noted the occurrence of nitrate at concentrations above the WHO guideline value (50 mg/l) in several groundwater samples from tube wells and open dug wells in north-western Burkina Faso. In a study of 168 tube well samples, 15% were found to have NO₃-N concentrations greater than 10 mg/l. In addition, of 123 samples of well water investigated, 36% had concentrations above this value. The maximum observed concentration was 90 mg/l (as N).

Nitrate concentrations were highest in areas with increased housing density and were also elevated in groundwater down gradient of the housing areas. Increased nitrate was found to be accompanied by increased electrical conductivity, which suggests that the concentrations of many other major ions (notably chloride, sulphate, sodium, potassium) were likewise increased as a result of the pollutant inputs. Yameogo and Savadogo (2002) also noted high concentration of nitrate in shallow aquifers in Ouagadougou particularly between August and early September. This, they attributed to high infiltration resulting from heavy rains.

However, in the dry season, the concentration level goes down significantly. Water samples of 5 out of 6 wells taken in the dry season showed nitrate concentrations below the acceptable WHO standard except the sample from Kossodou well which had higher concentration (250 mg/l). High nitrate concentrations are likely to be a feature of shallow groundwater in many areas of the country. The concentrations of ammonium and nitrite are less clear.

- **Salinity**

Groen *et al.* (1988) found variable salinity in groundwater samples from their study of north-western Burkina Faso. Electrical conductivity values were in the range of 50–2700 µS/cm. The highest values observed are relatively saline and unsuitable for potable use. However, median values determined (270 µS/cm and 370 µS/cm respectively for dug wells and tube wells) indicate the presence of predominantly fresh groundwater. The salinity variations in groundwater from the sedimentary formations along the north-west and northern border areas of Burkina Faso are not known. Yameogo and Savadogo (2002) found salinity range of 100-350 µS/cm from 25 wells sampled in Ouagadougou and in the Baskuy community the figure ranges between 326 and 1595 µS/cm in water sampled from 15 wells with permanently high levels of nitrate.

- **Fluoride**

Fluoride data are not currently available for Burkina Faso groundwater. Smedley *et al.* (1995) and Ghana Ministry of Works and Housing (1998) reported often high fluoride concentrations (up to 5 mg/l) in some granites and some meta-igneous horizons in the Birimian basement from the neighboring Upper Regions of Ghana. This suggests that areas of granite areas especially and possibly other areas of the crystalline basement of

Burkina Faso can carry groundwater with unacceptably high fluoride concentrations (greater than the WHO guideline value of 1.5 mg/l). Fluoride concentrations are likely to be higher in tube wells abstracting from the basement fractures than in shallow dug wells.

- **Arsenic**

Data for arsenic in groundwater from Burkina Faso were not available as at the time of writing. Many accounts suggest that arsenic problems exist in parts of Burkina Faso, although the extent and scale of the problem is not well defined. De Jong and Kikietta (1980) noted the occurrence of locally high arsenic concentrations in a village close to Mogtedo in central Burkina Faso. Appelo and Postma (1993, p248) also indicated from an original investigation carried out by IWACO (1975) that arsenic problems occur in some Burkina Faso groundwater. The high concentrations were taken to be the result of oxidation of sulphide minerals (e.g. pyrite, arsenopyrite) and are likely to be concentrated in and around the gold mineralized areas.

Arsenic presence in the drinking water is said to have given rise to the serious chronic arsenicosis condition known as blackfoot disease. If so, this would imply the occurrence of very high concentrations in affected potable supplies (several hundreds of µg/l) and is a cause for concern. Verification of the location(s) of the noted health problems and instigation of a water-testing programme for arsenic, especially in the mineralized areas, will help to define the nature and scale of the problem in Burkina Faso and form the basis for providing low-arsenic alternative water supplies.

- **Iodine**

As with many other chemical constituents, analyses of iodine are not available for the groundwater. However, the region is arid and remote from the influences of maritime rainfall. Thus, iodine inputs from atmospheric sources is likely to be low. This, combined with the dominant hard-rock geology of the region, means that iodine concentrations in groundwaters (as well as soils and locally produced crops) are likely to be low (of the order of a few µg/l). Indeed, iodine concentrations were found to be low or very low in many groundwater samples from the neighboring Bolgatanga area of northern Ghana (Smedley et al., 1995). Therefore, it is possible that iodine-deficiency problems occur in the Burkina Faso population if dietary iodine supplements are not used.

II.4.2 A geographical triptych

A. Agro-Ecological Zones in the Volta Basin of Ghana

The natural vegetation of Ghana is closely related to the ecological zones. Six agro-

ecological zones, defined on the basis of climate, reflected by the natural vegetation and influenced by the soils are recognized in Ghana (Figure 24).




Figure 24: Map showing the agro-ecological zones of Ghana and Burkina Faso

These consist of the Sudan, Guinea and Coastal Savanna Zones, the Forest-Savanna Transitional Zone, the Semi-deciduous Forest Zone and the High Rainforest. In all these zones, the natural vegetation has undergone a considerable change as a result of human activities. Also considerable variations exist between successive rainy seasons in time of onset, duration and amounts of fall. Rainfall is also generally accompanied by high intensities and energy loads and is therefore erosive. Generally, Alluvia soils (Fluvisols)

and eroded and shallow soils (Leptosols) are found in all the agroecological zones (FAO-RAF, 2000/1).

- **The High Rainforest**

The rainforest covering an area of about 7500 km² is located in the south-western corner of the country. This zone is characterised by a bimodal rainfall distribution pattern with a mean annual total ranging from 1750 to 2200 mm. The major rainy season occurs from March to mid-July with a peak fall in June. The minor rainy season starts from early September and tails off at the beginning of November. The major dry season commences from mid-November and lasts till the end of February. Temperatures are generally high throughout the year. The mean annual maximum temperature varies from 28.7 to 31 °C with a corresponding minimum temperature of 20.6 to 23.2 °C. The mean temperature is about 26.5 °C. Relative humidities in the morning are over 90 %. In the dry season, the value falls below 80 %. Potential evapotranspiration in this zone is about 1350 mm/yr (FAO-RAF, 2000/1).

The vegetation in the rainforest is generally evergreen although some species common to the semi-deciduous forest may be found. Such species tend to shed their leaves during the dry season. The zone is characterized by the *Cynometra-Lophira-Tarrietia* association with *Cynometra ananta*, *Lophira alata* and *Tarrietia utilis* as indicator trees (Lane, 1962). The topography is undulating to rolling with numerous fresh water swamps potentially suitable for rice cultivation occupying the low lying valley bottoms. The swamp vegetation consists of *Raphia* palms with shrubs such as *Alchornea cordifolia*, *Caropa procera* and *Macaranga* spp. entangled by various climbers. The soil types in this zone are: Acrisols, Nitrisols and Gleysols (FAO-RAF, 2000/1).

- **The Semi-Deciduous Forest (SDF)**

The Semi-deciduous forest zone is about 66300 km² in extent and forms about 90 % of the total forest zone. Like the HRF zone, the SDF zone is characterized by a bimodal rainfall distribution pattern with mean annual totals ranging from 1400-1750 mm. Conditions of temperature potential evapotranspiration and relative humidities are similar to that in the HRF zone (FAO-RAF, 2000/1).

The characteristic associations are Celtic-Triplochiton and Antiaris-Chlorophora. The indicator trees for the former consist of Celtic milbraedii and Triplochiton scleroxylon whilst the latter is characterized by Antiaris Africana and Chlorophora excelsa. It is within this zone that most food crops and cocoa cultivation takes place. Most of the timber for both local needs and export comes from the zone. As a result of these activities the vegetation outside forest reserves consists mainly of forb regrowth, thicket, secondary forest and swamp thicket. Soils present in this zone are: Acrisols, Nitrisols and Gleysols (FAO-RAF, 2000/1).

- **The forest-Savanna Transitional Zone (Derived savanna)**

This zone covering about 8300 km² occurs as a normal strip of about 48 km wide along

the north and the north easting limits of the semi-deciduous forest. This zone is also characterised by a bimodal rainfall distribution pattern with mean annual totals ranging from 1200-1400 mm. The mean temperature is about 26.5 °C. Morning and mid-day relative humidity values range between 85 % and 88 % and 70 % and 74 % respectively. Potential evapotranspiration in this zone is about 1350 mm/yr (FAO-RAF, 2000/1).

Most tree species of the forest zone occur in this area in addition to such species as *Daniella oliveri*, *Borassus aethiopum* and *Terminalia macroptera*. These trees occur in association with tall to medium grass such as andropogon and *Pennisetum* spp. The soil types here are: Lixisols, Nitrisols, Plinthosols and Cambisols (FAO-RAF, 2000/1).

- **Guinea Savanna Zone**

The guinea savanna zone which covers almost the northern two-thirds of the country is the largest ecological zone. Its aerial extent is about 147900 km². This zone is characterized by a unimodal rainfall regime lasting from April to October with a mean annual rainfall varying between 1000 and 1200 mm. Monthly total increases gradually from March until a maximum is reached in August or September, after which monthly total falls rapidly. It is dry November and March. This is the period when the desiccating effect of the harmattan is strongly felt. Mean annual maximum temperature ranges from 33 °C to 35 °C with a minimum of about 22 °C and a mean of 27.8 to 28.5 °C. Relative humidity is about 40 % in the dry season but may reach 84 % during the peak of the rainy season in August. Potential evapotranspiration varies between 2000 and 2300 mm/year (FAO-RAF, 2000/1).

The vegetation consists typically of a ground cover of grasses of varying heights interspersed with generally fire resistant, deciduous, broad leaved and gnarled trees at the forest margins in the south. This grades into a more open grassland with widely spaced shorter trees towards the north. Owing to the tussocky nature of the grasses bare patches of land are common. During the dry season, November to March, the dry grasses are highly inflammable. In the less eroded areas, *Andropogon gayanus*, the commonest grass, may be replaced by *Hyparrhenia* and *Heteropogon* spp. While *Aristida* and *Cymbopogo gigantus* dominate the badly eroded sites. *Vetiveria nigriflora*, *Seteria anceps* and Sedges occur in alluvial sites. The common tree species, including *Lophira lanceolata*, *Anogeissus*, *Azadirachta africana*, *Prosopis africana*, *Pterocarpus erinaceus*, *Parkia clappertoniana*, *Butyrospermum parkii* and *Antiaris africana*. Fringe forest and woodland may be found along the water courses. Soils present in this zone are: Lixisols, Acrisols, Luvisols and Gleysols (FAO-RAF, 2000/1).

- **Sudan Savanna Zone**

The Sudan savanna zone covers an estimated area of 1900 km². It is characterized by a unimodal rainfall pattern with a mean annual rainfall varying between 900 and 1000 mm. Mean annual maximum temperature ranges from 27.8 to 28.5 °C. Relative humidity and potential evapotranspiration are similar to that of the Guinea Savanna Zone (FAO-RAF, 2000/1).

This zone consists of short drought and fire resistant deciduous trees, interspersed with open savanna grassland. Grass cover is very sparse and in most areas, the land is bare and severely eroded. Tree cover is very low. However, in the densely settled and cultivated areas, important economic trees such as *Adansonia digitata*, *Ceiba pentandra*, *Butyrospermum parkii*, *Parkia clappertoniana*, *Tamarindus indica* and *Acacia albida* still remain. The soil types here are: Lixisols, Acrisols, Luvisols and Lithosols (FAO-RAF, 2000/1).

- **Coastal Savanna Zone**

The Coastal savanna zone covers an estimated area of 4500 km² with mean annual rainfall varying between 600 and 900 mm. This zone has a bimodal rainfall pattern with a characteristic distribution similar to that of the forest zone. The mean annual maximum and minimum temperatures are 30.5 and 22.9 oC respectively. Relative humidity varies from 55 to 65 % during the day and fall to about 40 % during the major dry season (FAO-RAF, 2000/1).

The vegetation consists of mainly grassland interspersed with dense short thickets often less than 5 m high with a few trees like *Antiaris Africana*, *Ceia pentandra* and *Milicia excelsa*, *Albizia zygis* and *Azadirachta indica*. Short and medium grasses are the dominant plant species, notable among which are *Andropogon gayanus* and *Hyparrhenia dissolute* in upland areas and *Vetiveria fulbibarbis*, *Brachiaria falcifera* and *Bothriochloa bladhii* in low lying areas. Soils present in this zone are: Acrisols, Luvisols, Cambisols, Gleysols, Vertisols, Solonetz and intergrades (FAO-RAF, 2000/1).

Exception of the Tropical Rainforest zone, the Volta Basin of Ghana falls in all the agro-ecological zones within the country. The largest proportion is in the Sudan/Guinea Savanna zones (24.2 %). The rest of the distributions are shown in Table 45 below:

Table 45: Proportion of Volta Basin in Agro-ecological zones of Ghana

Agro-ecological zone	Proportion of Volta Basin in the zone
The Semi-Deciduous Forest	5.4
The forest-Savanna Transitional	11.2
Sudan/Guinea Savanna	24.2
Coastal Savanna	0.5

B. Major Farming Systems in the Volta Basin of Ghana

The term farming-system refers to a particular arrangement of farming enterprises (e.g cropping, livestock-keeping, processing farm products) that are managed in response to the physical, biological and socio-economic environment and according to the farmers' goals, preferences and resources. Farming is used here in a wide sense to include not only crops and livestock but the other natural resources available to the farm household, including resources held in common with others (Reijntjes *et al*, 1992).

Two farming systems are dominant in Ghana: the bush fallow system (temporary system) and the permanent system. Some variants of both systems are also found in the Volta basins: the HUZA and the mixed farming systems.

- **The bush fallow system**

This is a system of land rotation between crops or fields and bush. A plot of land is cultivated for a number of farming seasons and abandoned when necessary to revert to secondary vegetation. It allows the soil to keep its nutrients qualities. The length of this resting period depends on how pressing the need is for land for cultivation. In the sparsely settled areas, the length rotation may reach 15 years, whereas in the densely settled areas, the period may be as short as 3 years. As soon as the farmer abandons a plot, he starts to cultivate the new plot. Thus he moves from one plot to another plot in different farming seasons. This system could be named shifting cultivation instead of bush fallowing. The latter term implies the movement of both settlements and farming to new areas when old farm lands are abandoned. In Ghana, farm settlements are usually permanent.

The average size of food farm cultivated under the bush fallow system is 1.10 hectares. During the dry season, men clear the land by hand. The vegetation is cut down and burnt. All trees of economic importance, such as shea butter, dawa-dawa and the oil palm trees are left standing. Sowing begins with the first rains. For planting yams, plantain and cocoyam, the hoe is used. In the case of cassava, maize, guinea corn and millet, cutlass is used. To improve the soil occupation and the yields, several kinds of crops are grown together on the same farm. Indeed the nutrient requirements are not the same for all the crops, so it is possible to obtain high yields from all the crops grown on the same farm or plot. Moreover, the different crops grow in different ways, this means that all the different crops together provide a better cover for the soil and as a result, help to fight the erosion.

This system of bush fallowing has some advantages in the peculiar environmental conditions of the tropics:

- The rotation allows the growing crops to make use of the plant food that accumulates in the soil as a result of the decay of leaves and twigs from the fallow vegetation. Consequently the harvest is satisfactory in the first seasons of cultivation. After 2 or 3 years, the farmer abandons the food plots to allow it to regenerate fresh supplies of plant nutrients.
- The burning of cleared vegetation on plots intended for cultivation saves time and labour. It also improves the soil. The ash contains carbonates and phosphates which are washed into the soil by the first rains, and so increase the soil fertility. The danger of this burning method is to affect the protecting vegetal cover and thus increases the risk of erosion.

The disadvantage of the system is that it is generally thought that a fallow period of 25 to 30 years is desirable. When the pressure on the land is too high this period cannot be respected. As a result, the length of the fallow period has consequently decreased and this has resulted in low crop yields. For the last two decades, the fallow period has been shortened to approximately 2-3 years. This has led to decline and deterioration of cultivated soils and yields. Using fire for cleaning vegetation exposes the soil to the sun and torrential rains until the first crop forms an effective protective cover.

- **The HUCA farming system**

It differs from the bush fallow because of the peculiar system of land ownership, which gives rise to the strip pattern of land. A co-operative regroups all the financial resources of the farmers. The company is organized for the sole purpose of collecting land. The land so acquired is called 'huza' described as "a tract of land bought by a group of people, often but not necessarily kinsmen, under an elected leader (Hutze) who makes all negotiations with the seller (Gyasi, 1976)". When the land is acquired, it is divided into strips for each farmer. The width of the strip is proportional to the farmer's financial participation. After two or three seasons of cultivation, when yields decline, the field is left fallow. Due to the pressure of population and the great demand for land, the length of the fallow period is now short (4-6 years).

- **The permanent systems**

Contrary to the bush fallow system, these systems are intensive and a piece of land is cultivated continuously. In Ghana, there are 2 permanent systems of food farming: the compound farming system and the Anloga-Keta system. In the Volta Basin, only the compound farming system is used.

- **The compound farming system**

It is used in the densely settled areas of northeastern and northwestern Ghana. This system centres on the household compound. The land immediately surrounding the compound house is intensively cropped with vegetables and staples using organic soil regeneration techniques, which involve using household refuse and manure from livestock. The average size of a compound farm is less than an acre. These pieces of land are used for cultivating okra, tomato, pepper, maize, cocoyam and plantain. These farmers cultivate also larger fields at some distance away from the household. In these fields, they adopted the bush fallow system. These outfields provide the main bulk of the farmers' food supply.

- **The Mixed Farming System**

It is characterised by a combination of cultivation with keeping of livestock. The latter provides power and manure on the farm. This system was introduced in the 1930s to check the rapid deterioration of soils as a result of population pressure on land and to increase agricultural production. In addition to using manure to increase production, many technical innovations were introduced such as the use of bullock plough and planting on ridges. Cow-dung, compost of household refuse, kitchen refuse and goats pens are applied to compound farms. Mixed farming is restricted to areas which are free from tsetse fly. The growing season is based on when rainfall is more than one-half of the potential evapo-transpiration, and ends when there is less than half the potential

evapo-transpiration.

In the Sudan Savanna Zone, the climate, which is characterised by the alternation of clear-cut wet and dry seasons, has a direct effect on soil forming processes in the area. Although the prevailing climatic conditions permit accelerated chemical decomposition and deep weathering of rocks, the sudden and torrential rainfall following a prolonged dry season, during which the vegetation cover is burnt, induces great soil erosion.

C. Land tenure characteristics in several agro-ecological zones of the Volta basin of Ghana

- **Sudan and Guinean Savanna Zones**

Various ethnic groups live within the zones. The village chief or the earth priest (Tendana) is in charge of administrating the land. According to traditional tenure rights, land is only allocated to the farm households. The usufruct rights can be passed on to the descendants so that land tenure is secure as long as the land is cropped. When the land is fallow the Tendana can redistribute it. There is no fixed rent paid to the chief or to the Tendana. A part of the harvest is given to them more or less as a sign of recognition of his authority. As the household is not the owner of the land, they cannot sell it. Rights of usufruct have considerable religious implications unless the occupier of the land stands in breach with the chief or the Tendana, he cannot be deprived of them.

Land transactions are limited to leasing which is rare. Land can temporarily be exchanged for bullock or tractor services. When population density rises and land becomes more scarce, the land which is fallow will be relocated. Thus, smaller households that do not have enough labour to cultivate all the land but who still want to keep the land at their disposal also try to lease the land to neighbours or relative in the meantime. Land security is very high as measured by rights over land and low incidence of dispute, except for migrant farmers from other regions. Customary law does not recognise female right of *usufruct*. This is likely to influence the type of crops cultivated by women.

As soon as the land is fallowed, it can be redistributed. There is virtually no restriction on the use of land for food crop production. Where land is leased to migrant farmers for agriculture, the control over trees and other perennial features on the land remain under the control of the original owner. Ownership of economic trees on farms belongs to individuals who develop the land but for economic trees in the bush, the first person to gain access to a tree can harvest its fruit.

The main difference between the Sudan and the Guinean Savanna Zones is the current land tenure. In the Sudan zone, the traditional land tenure systems have been generally maintained. Land is bought and sold around larger urban. In the Guinean Savanna Zone, recent developments and pressure on land have resulted in the flagrant violation of traditional regulations on land. Now land is sold for cash or in kind or both depending on the contractual arrangements.

- **Forest Savanna Transitional Zone**

Access to land for farming can be gained from family, from outright purchase of land, by paying fee to chiefs or local district assemblies or by hiring and share cropping.

Table 46: Access to land

Method of lease	Percentage
Hire land	35
Sharecrop	14
Hire out	8
Share out	4

In the food crop sector, leasing is usually based on short-term sowing contract. The maximum period of loan is usually three years, but often for one year. With the sharecropping tenancy, 1/3rd of the harvest (generally maize) or 1/3rd of the proceeds from sales is restored to the landlord while the tenant retains the rest. With the increase of land pressure, the landlord, in some areas, increases the share to half.

The Hiza system land tenure (as described below) is practiced in the southeastern part corner of the forest/savannah transition zone.

- **Deciduous Forest Zone**

In the Deciduous Forest Zone, land in the community belongs primarily to the chief. However, most of the land is now in the custody of families. The chief has direct control over virgin forest only. Land is often given to migrants on share cropping terms. In some areas the migrants develop new plantations that are later shared between the tenant and the landlord. Like in the Forest Savanna Transitional Zone, the landlord gets 1/3rd of the harvest. With the increase in land demand the landlord has been able to demand a larger part. In some cases, farmers hire land. With this system, the control of the farmer on his land is minimal. Tenants are worried that landlords would take the land from them when plantations mature. Cocoa has become the dominant landscape in the zone. In giving land to migrants, landowners place restrictions on the use of their land. They may oblige settler farmers to plant cocoa because of economic reasons.

D. Major farming activities in the various agro-ecological zones of the Volta Basin in Ghana

i. Cropping systems in the Volta basin of Ghana

- **Sudan Savanna Zone**

The basis of the cropping system throughout the zone consists mainly of pearl millet. The early millet is inter-planted with late millet or sorghum in fields close to compounds where fertility is highest. Maize was highly available at subsidized rates. Since fertilizers was no longer available and soil fertility had declined, crop planting of crop fell drastically in 1997.

There has been a spread of European vegetable cultivation with the gradual decrease of

some minor indigenous crops. Carrots, cabbage, lettuce and peas are now available and grown on irrigated plots close to towns and were probably originally planted for expatriates.

- **Guinean Savanna Zone (GS)**

Maize is the major cereal crop produced in this zone. More than 80% of the small-scale farmers grow maize. Maize has a special position because of government support for the crop. Improved varieties are used. Every farm family cultivates sorghum either as a sole crop or as an inter-crop. Cereals like maize, sorghum and millet cover most of the cultivated land. However, areas under maize decrease with increasing population density, while areas under millet or sorghum increase. Cotton, a cash crop, is found at specific sites throughout the Savanna Zone.

Table 47: Proportion of area allocated to major crop species and yield per hectare in the GS

Crops	Average area (%)	Yields (Mt/Ha)
Cotton	2.28	-
Groundnut	38.28	0.52*
Cowpea	31.05	0.60#
Yam	14.31	2.98*
Cassava	23.36	3.9*
Sorghum	48.18	0.70*
Millet	62.10	0.68#
Maize	59.21	1.00+
Rice	3.78	0.90+
Tomato	-	1.90+
Pepper	-	0.9+

*Source: *MOFA 1999, + Runge-Metzger and Diehl 1993, # Diehl 1992*

Cash crops are groundnuts, cowpeas, maize and rice. Mainly women are responsible for marketing the cereals. They pass on the money to their husbands. The women also sell products to the market when the quantity harvested is large.

- **Forest Savanna Transitional Zone**

Food crop production dominates the farming system. Animal production is higher in this area than in the forest. It is the zone for major commercial food production: maize, cassava, groundnut and yam. Oil palm is also important as it is reserved in fallow land rather than grown plantations. Cotton and tobacco are important cash crops.

- **Deciduous Forest Zone**

Cassava and plantain are the important food crops while cocoa and oil palm are important as cash crops. Vegetable production is increasing in importance in the zone. Cocoa is the most important cash crop in this area. The indigent and settlers grow it.

ii. Crop husbandry processes and activities

- **Sudan Savanna Zone**

Population density has a great influence on cultivation systems. Until recently, shifting cultivation was the dominant system with the increase in land pressure, farmers

started to practice permanent cultivation. Compound farms and bush farms are found in this zone. On the bush farm, no manure is applied and it consists in land rotation. The plots take place 2-4 kilometres away from the farm. In the compound system, the land cultivated is directly around the homestead and is fertile because household and farm refuse are used as manure. Tobacco, gourd, melon, okra, tomatoes, pepper and sweet potatoes are usually cultivated in the compound lands. Further away is another zone planted with early and late millet, guinea corn, bambara beans and cowpea. This second zone is fertilized with farmyard manure though this is often inadequate. The rest of the compound area, usually the largest, has no manure application and is cropped to guinea corn and late millet.

The compound farms produce higher yields than the bush farm. In the first case, the soil fertility is maintained due to manure whereas in the bush farm, soil is left to nature to restore through the fallow system. With increase in land pressure, the compound system has its advantage. The bulk of cereals are however produced in the bush fallow farm. Usually the system of cropping is guinea corn and late millet mixed together, or both planted singly and often inter-cropped with groundnut, bambara beans or cowpeas. Rice may be planted in poorly drained soils, or grown inter-planted with early millet.

The pressure on land and the need to produce cash crops for sale have gradually brought about innovative farming techniques. Migrants like Muslim, seem to be the first to practice horticulture in rivarian areas. Many of the larger rivers are not exploiting in this way because of a lack of inadequate methods of lifting water. However, most of the shallow rivers and seasonally flooded land are now given over to dry-season gardening. In addition, gardens have been established on the edges of dugouts, after they had been excavated. There is still an opportunity to develop some of the larger rivers for irrigation, given appropriate water-lifting technology.

Onion cultivation is particularly popular and probably represents one of the most important agricultural exports from the zone. Cereals produced are locally consumed and so, are not open to the commercial market. Sheanuts are bought and exported on small scale. Tomatoes and onions are produced for sale as cash crops and are exported to Southern Ghana.

• **Guinea Savanna Zone**

Animal production is more important than it is in the rest of the Savanna Zone. However, food production dominates. Bullock is also used for ploughing although some farmers are not able to afford it. Tractor may be used but at a higher price (between ₵25000 and ₵30000 per acre for the bullock, and ₵35000 for the tractor.) Like in the Sudan Zone, farming systems practiced are bush fallow and compound farming.

The major cropping system is mono-cropping of early maturing maize within the compound. The following groups of cropping systems may be distinguished in the zone:

- Maize, sorghum, groundnut and cowpea with root crops, namely yam and cassava that occur in the central portion of the zone;

- Sorghum based but mixed with maize or cowpea and yam, occurs in the western part of the zone;
- Yam, maize, sorghum, groundnut based system, occur in the southeastern part of the zone.

Table 48: *Cropping patterns in the Guinea Savannah zone*

Cropping pattern	Percentage area (%)
Sole	4.0
Mixed cereal	15.4
Cereal/legume	62.2
Roots/others	12.0
Others	5.0

The choice of soil tillage is influenced by ecological and economic factors such as soil type, land use of the preceding year, crop that is actually to be sown or planted, and the available technological options. Tractor is used for heavier lowland soils whereas hoe and bullocks till sandy upland soils. Soil preparation is done by hoe.

Table 49: *Percentage of use of each technology*

Hoe	77.4 %
Bullock	14.6 %
Tractor	8.0 %

Soil nutrient stocks are replenished by fallowing, the use of organic manure, biological processes, rainfall, sedimentation and mineral fertilisation. The application of fertiliser and manure is still not a common practice and it fluctuates from year to year. The majority is applied to crops which show an elastic response to the fertiliser, such crop as maize, rice and vegetables.

Men and women have distinct roles: men usually carry out land, clearing and ploughing, while women gather and burn the cleared weeds and later plant all crops. Women do most of the marketing and are responsible for the daily cooking and childcare. Almost every farmer in the zone has some livestock. About 90 % of all women have 5-10 chickens; about 29 % have 2-5 goats. Some 89 % of all men have sheep, particularly in the Dagbon area and 10 % have cattle.

- ***Forest Savanna Transitional Zone***

In this zone, permanent mechanized cultivation of food crops is common. Many farmers have adopted technologies based on ploughing, permanent cultivation and use of chemical fertilisers. The transitional character of the ecosystems, the ethnic and cultural diversity resulting from migration led to a considerable diversity in farming systems and crops. The widest variety of crops is grown in the transition zone. Mixed or sole cropping is used and the major cropping systems in the forest area are sole maize, maize/cassava, maize/cassava/plantain and /maize/pepper. In the Savanna area of the zone the cropping systems are sole yam, sole groundnut, rice/cassava and yam/cassava.

If the maize is planted in the first season, the inter-cropped culture (cassava or plantain) is relayed to the maize field at maize tasselling stage. To lower costs, maize is cropped

twice a year or inter-cropped with cassava. Groundnut is planted during the first season, it normally follows yam. In this zone, single stand maize gradually displays the maize-cassava intercrop. Sole yam is the second most important cropping system. Mixture of pepper, garden egg and okra is the third. Legumes like cowpeas and groundnuts are rotated with cereals and yam.

Table 50: *Percentage of farmers practicing cropping system in Wenchi*

Cropping system	Percentage
Cocoa (with food crops)	0
Plantain intercrop (cocoyam, yam, cassava)	3
Maize cassava intercrop (cocoyam, vegetables, pulses)	7
Vegetables (pepper, garden egg, okra)	17
Sole cassava	5
Sole maize	26
Sole tomato	5
Cowpea	5
Groundnut	10
Rice	0
Yam	19
Others	3

- **Deciduous Forest Zone**

In this zone, the systems all have combination of food crops and at least one tree crop. For example cocoa or oil palm is combined with food crops like plantain, cassava, cocoyam and some other minor crops. The farming system involves permanent cultivation of tree crops, and rotational bush fallow of food crops. The first crop usually planted is maize, which is planted in almost every part of the farm. Two or three weeks later, trees (cocoa or oil palm) are planted, followed by plantain, cassava or vegetable in the same piece of land. Farm sizes vary from 1 acre to 15 acres with the most recurring farm size being 2 acres followed by 1 acre.

iii. Livestock production

- **Sudan Savanna Zone and Guinea Savanna Zone**

Livestock may be owned individually or by a family. Sheep, goats, fowls and guinea fowls are kept by many household.

Table 51: *Percentage of farmers keeping livestock, 1992*

Cattle	Sheep	Goat	Chicken	Guinea fowls	Duck	Turkey	Pig
58.3	60.0	68.3	73.3	48.3	33.3	5.0	28.3

The animals are free during the dry season and tethered to a post in uncultivated patches of grass near the farm in the rainy season. Grazing lands are poor and are those obtained under natural conditions.

- **Forest Transitional Savanna Zone**

In this zone, poultry, sheep, ducks and goats are kept in extensive and or semi-intensive

management systems, whilst pigs are kept under an intensive system. Chickens are kept in coops during the night and left on free-range during the day. The animals are seen to be liquid assets, which can be sold easily on the local markets. Pigs are kept in the relatively urban settlements.

- **Deciduous Zone**

As a result of the susceptibility of livestock to Trypanosomiasis and other diseases, the zone keeps very few livestock. It also results from a difficulty of integrating livestock with arable farming particularly where farmers have to walk long distances to farm. Small livestock are allowed to roam and graze around the village.

E. Constraints to agricultural production in several agro-ecological zones of the Volta basin in Ghana

- **Savanna Zone**

- a long and intense dry season
- the pattern of rainfall is uncertain: the single wet season may start as early as the beginning of March or as late as the end of June;
- the major proportion of the early rains is lost due to surface runoff;
- annual bush burning aggravates the low levels of organic matter in the topsoil;
- debilitating human diseases such as *Onchocerciasis* and malaria
- livestock pest and diseases such as *Trypanosomiasis*, are a major constraints to increase livestock production;
- scarce drinking water;
- poor and deteriorating infrastructure and communications systems; and
- remoteness from major domestic and overseas markets.

- **Savanna Transitional Zone**

The major problems are:

- high costs of farming input and fluctuation in their prices;
- high interest rates by credit institutions;
- erratic rainfall;
- long-term management strategies for the land use are difficult because the landlord may deprive the farmer of the land when he wishes;
- the aridity of the zone in the dry harmattan produces stress in cocoa, thereby resulting in such diseases as swollen shoot; and
- prevalence of livestock diseases.

- **Deciduous Forest Zone**

The major problems are:

- tsetse infestation limits the exploitation of livestock potential;
- cocoa suffers from black pod and the spread of the swollen shoot virus;
- although the soils are fertile in this zone, when the forest is cleared for cultivation, the nutrient status is drastically reduced as a result of the organic matter being quickly oxidised, the topsoil dries hard when exposed to the sun;

the yield of each crop was estimated.

Table 52: Estimated Average National wholesale prices of crops/tonne (1998)

Crop	Price (cedis)
Maize	704,670
Cassava	304,473
Millet	751,807
White yam	620,793
Sorghum	616,110
Cowpea	1,123,018
Groundnut	1,515,232
Yam	670,793
Rice	1,047,070

Source: Ministry of Food and Agriculture, Ghana, field estimates.

The revenue from each component crop in the crop mixtures taking account of the spatial and temporal arrangements on a piece of land when added up, give an estimate of the total revenue per area for the cropping or farming system. This gives an indication of the sustainability of the different cropping systems. To determine the economic profitability of different cropping or farming systems, the cost of input and activities is considered.

Examples of cash flow analyses are illustrated for food crop intercrop with limited rotational bush fallow system (sorghum, maize and yam) at low, intermediate and high input levels on Ferric Acrisols over a 4-year period.

These results show that improving in the farming/cropping system could be done for better yields and better revenue. Increasing the input levels from low to intermediate and from intermediate to high brings a significant increase in the net cash flow.

Farming on all the soils is however relatively sustainable for all cropping systems at both the intermediate and high input levels.

• Conclusion

From the analyses, we conclude that in the Savelugu Nanton District, the most profitable farming/cropping system is the one involving the mixture of sorghum, maize, groundnut, cowpea, cassava and yam. It may be used at all level of input and on all the soils. However, since profitability increases with increasing level of input used, the most sustainable farming system in the Interior Savanna Zone is the high input level. Farmers in this zone should be encouraged to increase their use of input where possible.

i. The Forest-Savanna Transitional Zone



Farming systems

The Techiman District in the Brong Ahafo Region of Ghana is discussed here as typifying

the farming system in this zone. The main tree crops include cashew, oil palm and teak. A quarter of the farmers grow tree crops. The three most common food crops grown are maize, cassava and yam. Two dominant cropping systems are used:

- Permanent tree crop system
- Rotational bush fallow of food crops system (Cassava + yam + maize or Plantain + yam + maize)

The average farm size in the zone is 1.4 hectares. Another important element of the farming system is the main rotation of annual cropping patterns. Most of the Transitional Zones that is Techiman Districts falls within the cropping systems zone with the length of growing period being between 210 and 240 days. Sowing and planting of crops start in mid- April, yams are planted from November to December and harvested from August onward.

The sequence of planting different crops in a year on a plot in the Techiman District is shown in the following Table 53.

Table 53: Sequence of planting in the Techiman District of Ghana

Position in sequence	Crops planted	Planting Period/Season
1 st	Maize	From mid April
2 nd	Cashew	March-September
3 rd	Plantain	April-September
4 th	Cocoyam, vegetables	April-May
5 th	Yams	November-December
6 th	Cassava	Throughout the year

There is also the rotation/sequencing of crops from year to year. A plot may be farmed for up to four years. Maize and vegetables are harvested the first year, the other food crops the second year. Cassava and yam are replanted in the third year and the plantains continue to be harvested up to the fourth year. Normally, farmers plant food crops with the trees until the tree crops form a closed canopy, which prevents light from reaching the food crops underneath them. Food crops are cultivated for 2 to 4 years and then the land is left to fallow for 2-3 years.

Table 54: Types of farming systems in the Techiman District of Ghana

National types	Agro-ecological zones types of cropping systems	Specific farming systems
Permanent tree crop systems	Cashew permanent tree crop systems = cashew + food crops (plantain, maize, cassava, cocoyam + other minor crops)	Cashew cropping systems without livestock Cashew cropping systems with some limited livestock = cashew + food crops + limited semi-intensive livestock (poultry, goats and sheep)
	Oil palm permanent tree crop systems = oil palm + food crops system (e.g. maize, cassava, cocoyam, plantain + other minor crops)	Oil palm permanent tree crop systems without livestock Oil palm permanent crop systems with some livestock = oil palm + food crops + limited semi-intensive livestock
Rotational bush fallow of food crop system	Plantain based rotational bush fallow of food crop system = plantain intercropped with other food crops	Plantain based rotational bush fallow of food crop system without livestock

(usually excluding cassava), in a rotational bush fallow system	Plantain based rotational bush fallow of food crop system + limited semi-intensive livestock (poultry, goat and sheep)
Cassava based Rotational Bush fallow of food crop system = cassava inter-cropped with other food crops (usually excluding plantain), in a rotational bush fallow system	Cassava based rotational bush fallow of food crop system Cassava based rotational bush fallow of food crop system + limited semi-intensive livestock

- **Interface between land evaluation and farming systems in the Transitional zone**

The main soils occurring in the Techiman District were identified to be Haplic Lixisols, Ferric Lixisols, Ferric Acrisols and Haplic Acrisols. The suitability of these soils varies from marginally suitable to very suitable. The Haplic Lixisols and Ferric Acrisols are suitable or very suitable for maize, cassava and cowpea cultivation. Haplic Acrisols are not suitable for maize at low input levels but are suitable for yam, cassava and cowpeas.

The use of input is generally intermediate (soil fertility maintenance levels of fertiliser or organic manure). Increasing input may be determined by whether the soil is suitable for a such crop or not; the description of each crop yield for each soil has been done. To determine the revenue from output of the crops in the farming system, estimated wholesale prices of outputs per tonnes, based on National Average Wholesale Prices were used.

Table 55: Estimated average national wholesale prices of crops/tonne (1998)

Crop	Price (cedis)
Maize	704,670
Cassava	304,473
Plantain	377,188
White yam	620,793
Cocoa	2,250,000
Oil palm	1,400,000
Groundnuts	1,515,232
Citrus	500,000
Cashew	650,000

Source: Ministry of Food and Agriculture, Ghana, field estimates.

Revenue from the yield of each crop was estimated by multiplying wholesale market price of crop by yields.

The costs of inputs for farming system in the Techiman District have been estimated. Thus the example of cash flow analyses may be done. In the Table 55, we can see the

levels on Haplic Lixisols for a period of 9 years

For all the soils types at low input level, the most sustainable cropping system is cassava food intercrop with rotational bush fallow. It gives the highest annual profitability on all soil types. For all the cropping systems, increasing the input level mean increasing cash flow. The amount of increase however varies with soil type and cropping system. The exception is the food crop systems on Haplic Acrisols, where an increase of inputs from intermediate to high level leads to a decrease in profitability.

• **Conclusions**

Currently, in this zone, farmers apply organic and inorganic fertiliser on their farms. However, an increase in the input level should be encouraged to move to the high input level. The most sustainable crop systems are cashew permanent tree with plantain intercrop on Haplic Acrisols and cashew permanent tree crop with cassava intercrop, when using high levels of input. On the other three soils, cassava food crop intercrop with rotational bush fallow is the most profitable.

iii. The Deciduous Forest Zone

• **Farming systems**

The Atwima District in the Ashanti Region is discussed here as typifying the farming systems in the zone. The main tree crops include cocoa, oil palm, coffee, rubber and avocado. The food crops are maize, plantain, cassava, cocoyam, yam and vegetables (tomato, okra, garden egg).

Farm sizes range between 0.45 and 6.8 hectares with a modal size of 0.9 hectares.



The two dominant systems are:

- The permanent tree crop system (cocoa + food crops or oil palm + food crops)
- The rotational bush fallow of food crop systems (mainly plantain, cocoyams and maize, white yam and vegetables)

Plantain and cassava are not usually intercropped. The food crop farms are fallowed after 2-4 years of farming. Most farmers in the district keep local fowls or poultry, whilst an average of farmers keep two goats and a sheep.

Table 56: Types of farming systems in the Atwima District of Ghana

National types	Cropping systems	Farming systems
Permanent tree crop systems	Cocoa cropping systems = cocoa + food crops (e.g. plantain, cassava, cocoyam + other minor crops)	Cocoa cropping systems without livestock = cocoa + food crops (e.g. plantain, cassava, cocoyam + other minor crops)

		Cocoa cropping systems with some limited livestock = cocoa + food crops (e.g. plantain, cassava, cocoyam + other minor crops) + limited semi-intensive livestock (poultry, goats and sheep)
	Oil palm permanent tree crop systems = oil palm + food crops (e.g. plantain, cassava, cocoyam + other minor crops)	Oil palm permanent tree crop systems without livestock = oil palm + food crops (e.g. plantain, cassava, cocoyam + other minor crops)
		Oil palm permanent crop systems with some livestock = oil palm + food crops + limited semi-intensive livestock (poultry, goats and sheep)
Rotational bush fallow of food crop systems	Plantain based rotational bush fallow of food crop systems = plantain inter-cropped with other food crops (usually excluding cassava), in a rotational bush fallow system	Plantain based rotational bush fallow of food crop system without livestock = plantain inter-cropped with other food crops (excluding cassava), in a rotational bush fallow system
		Plantain based rotational bush fallow of food crop system with some livestock = plantain inter-cropped with other food crops (excluding cassava), in a rotational bush fallow system + limited semi-intensive livestock
	Cassava based rotational bush fallow of food crop systems = cassava inter-cropped with other food crops (usually excluding plantains), in a rotational bush fallow system	Cassava based rotational bush fallow of food crop system = plantain inter-cropped with other food crops (excluding plantain), in a rotational bush fallow system
		Cassava based rotational bush fallow of food crop system = plantain inter-cropped with other food crops (excluding plantain), in a rotational bush fallow + limited semi-intensive livestock

In almost all cases, the use of purchased inputs like fertiliser, organic manure, lime and pesticides, and veterinary drugs is very low.

- **Interface between land evaluation and farming systems in the Deciduous Forest**

Zone

The main soils occurring in the Atwima District are Ferric Acrisols, Haplic Alisols, Dystic Fluvisols and Rhodic Nitosols. The suitability of these soils varies between unsuitable to very suitable. The low input level is the current level used in the district.

In order to determine the revenue from outputs of the crops in the farming system, estimated wholesale prices of the outputs of the crops per tonne, based on National Average Wholesale Prices, obtained from the Ministry of Food and Agriculture (MOFA).

Table 57: Estimated Average National wholesale prices of crops/tonne (1998)

Crop	Price (cedis)
Maize	704,670
Cassava	304,473
Plantain	377,188
White yam	620,793
Cocoa	2,250,000
Oil palm	1,084,226
Groundnuts	1,515,232

Source: ministry of food and agriculture, Ghana, field estimates.

As it was for other districts in the other agro-ecological zone, a typical cash flow analysis has been done. The example done is for cocoa with food crops intercrop system at low and medium input levels on Ferric Acrisols over 9 years.

- **Conclusion**

After analysis on Ferric Acrisols, Dystric Fluvisols and Haplic Alisols, the most sustainable system is plantain food crop intercrop with rotational bush fallow. For Rhodic Notisols, the most sustainable cropping system is the cocoa with food crop intercrop.

Using Dystric Anvisols for any cocoa with food crop intercrop farming system or cassava food crop intercrop with rotational bush fallow system is actually not sustainable and should be discouraged. For Rhodic Notisols, intermediate input level is always more sustainable than the high input level.

B. The Agro-ecological zones in the Volta basin of Burkina Faso

Five agro-ecological zones have been identified by the CNRST, based on agro-climatic, socio-demographic conditions and regional constraints and potentials. These zones have the following characteristics:

- **The Eastern zone**

This zone covers an area of 60,600 sq.km with a population density of less than 20 inhabitants per sq. km. It has an annual rainfall of between 500 mm and 1000 mm. It has great potential in animal husbandry, fishery and wildlife resources (5 reserves). The farming system is dominated by sorghum and millet.

- **The Sahelian zone**

This zone covers an area of 36,896 sq. km with a very low population density of 10 to 17 inhabitants per sq. km; it has a rainfall of between 300 mm and 600 mm and highly degraded vegetation cover due to drought. Its agriculture is subsistence and consists mainly of cereals. The main source of income is animal husbandry.

- **The North-western zone**

This zone has an area of 30,870 sq. km., a rainfall of between 500 mm. and 800 mm. It has an average population density of 41.1 inhabitants per sq. km. Although the farming system is based on cultivating sorghum and millet, the zone has great potential for irrigated crops such as vegetables, rice and maize. Animal husbandry is also significant.

- **The Central zone**

This zone covers an area of 94,000 sq. km with a high population pressure resulting in severe degradation of the plant cover and soils. Rainfall varies between 600 mm and 900 mm. The agro-pastoral and forestry potentials are limited and the farming system which consists of agriculture combined with animal husbandry is dominated by small ruminants and poultry.

- **The Western zone**

This zone has an area of 52,000sq.km, and a rainfall of between 700 mm and 1200 mm. The zone has great natural resources potential. It is the converging area for immigrants from the Central and Northern parts of the country. Its agriculture is much diversified with high output in cotton, fruits, vegetables and dominance of cereals (sorghum, maize, millet, rice and fonio). It has a high potential for animal husbandry, and possesses a large stock of cattle.

Subsistence farming is very widespread and is essentially manual with not much external input. Animal traction is mainly used in the cotton-growing tracts (cash cropping) where modernisation (mechanisation and use of agrochemicals) is well advanced.

C. Agricultural regions and production systems in Burkina Faso

Many regional divisions of Burkina Faso were made by the technical services of ministerial departments (MARA, MEE, MAT...) following precise specific objectives. The agricultural zones described here are those determined by INERA. They cover the entire agroecological zone described under this section and the entire Volta Basin in Burkina.

- **The Northern Region: The SAHEL**

It covers most of the Burkinabè Sahel and includes the Sahel provinces of Oudalan and Soum. It is the driest region of the country. The rainy season, which lasts approximately three months, extends from June to September. Rains are erratic and the total rainfall in a year is hardly more than 600 mm. Evapotranspiration there is very high and is combined with high amplitudes of temperatures during the day and at night.

By tradition, it is a livestock zone. Millet is the main crop, while white sorghum follows. There are almost no rotation crops. Fallowing decisions depend on the displacement of animal habitat or penning (KAFANDO, 1995). Night penning of animals on plots after harvests is the main form of soil fertilisation. The input of chemical fertilizer is negligible. Animal traction, which has been introduced by vulgarization agents is not yet generalized. However, in the Soum there are a few hitches

with donkeys and camels. Usually agricultural work is manual in the region where the margin of manoeuvre of producers is narrow as far as the choice of cropping and production system is concerned.

- **The Central Region**

It covers the following provinces: Sanmatenga, Namentenga, Oubritenga, Boulkiemdé, Sanguié, Kadiogo, Ganzourgou, Bazèga, Zoundwégo, Sissili and Nahouri. It extends to almost all the central plateau, with an annual rainfall ranging from 600 mm in the north to 900 mm in the south. Unequally distributed, rains spread over 4 to 6 months. Agriculture in this region is mainly rain fed.

Due to its high population density, this region experiences serious problems of environmental degradation resulting from the overexploitation of its meager resources. The population pressure in the centre is such that there is practically almost no fallow anymore. Therefore, soil fertility is not restored, accelerating in this way its degradation and aggravating the adverse effects of wind and water erosion. The input of fertilizers to make up for and restore crop exploitation is weak. This system of land use gradually leads to soil depletion, hence the notion of land overexploitation.

Farming systems in the regions are based on cereals like in the east. Sorghum and millet come first, i.e. about 80 % acreage, followed by groundnuts and quite far behind is maize. The introduction of animal traction dates back to the 1960s. In general, traction equipment is limited to ploughing before planting, particularly for cash crops (groundnut and cotton). It cannot be said that animal traction is a characteristic of this region as most farming activities are still done manually. Local varieties of sorghum and millet are still preponderant. Recourse to improved seeds concerns only groundnut and rice.

As a result of the many water bodies in the central regions, market gardening is developing. Since it is an out of season activity, producers have the opportunity to get to work and increase their incomes.

- **The North-Western Region**

It includes the provinces of Bam, Passoré, Yatenga and Sourou. This region is characterized by a rainfall which varies between 600 mm in the north and 800 mm in the south. The dominant economic activity is livestock raising, with, however, fewer number of animals than those of the Sahel and the central regions. But the degradation of the climatic conditions obliged producers to adapt themselves. In this sense farming systems in this region are now based on the couple millet-sorghum (white). Groundnut comes in third position. Peco-climatic conditions offer producers in this region little choice in crop diversification. Croplands are lacking and they are continuously used under rotations: millet-sorghum-groundnut. However, it can be observed that farmers are making efforts to overcome this hostility of nature. It is one of the regions, where using organic manure (animal manure and excreta), in combination straw is quite common.

In provinces like Yatenga and Passoré, the Zai (improved traditional technique) is used to restore deteriorated land. Although the introduction of animal traction dates far back to the beginning of independence (BDPA and SATEC intervention), the practice also constraints for many reasons: soil fragility, high costs, maintenance of draught animals. As a result, farming activities still remain manual. In the north-western region, it is worth noticing the existence of the Sourou valley, which provides great opportunities for cultivating irrigated rice and maize and for market gardening. With irrigation, producers undertake two forms of campaign for rice and earn substantial incomes. It must be noticed that in the irrigated areas, agricultural intensification techniques are used.

- **The Eastern Region**

It covers the provinces of Boulgou, Kouritenga, Gourma, Gnagna and Tapoa. Some provinces (Gourma, and Tapoa) in this region are the least populated and thus the least deteriorated in Burkina Faso. The annual rainfall varies between 600 mm and 900 mm. The region shelters the country's big fauna reserves. It is considered as a cereal producing region. Farming systems there are characterized by the predominance of sorghum and millet in rotations. Groundnut comes next. In recent years, the penetration of cash crops such as cotton has been noticed due to political incentives.

- **The Western Region**

It covers the provinces of Kossi, Mouhoun, Houet, Kéné Dougou, Bougouriba, Comoé and Poni. The rainfall is in the range of 900 mm and 1100 mm. It is the region with the best agricultural potential. Maize is the main food crop. Growing of rain-fed rice is also being developed. It is the chosen zone for the main industrial and cash crops (sugar cane and cotton).

The western region grows yam. Mainly cultivated in the Comoé and Poni provinces, the position of yam is relatively important in the farming system. Its cultivation demands rich soils, hence the need to clear new plots as fields become poor. Degraded soil is a destructive factor of biological diversity.

The western region, nevertheless is the modernization of agriculture is fast with the use of improved seeds and grain drills, mechanical ploughing and weeding and treatment with insecticides. In addition to the large adoption of animal traction, favored by the cultivation of cotton, an experiment of mechanization occurred resulting from the financial facilities provided by cotton cultivation. This intensification of agriculture is limited in the short term by the fragility of soils whose fertility conservation is not guaranteed. The low rate of organic matter in the soil and the need to restore soil depletion constitute challenges for most producers in the region despite their satisfactory technical level.

In the region, fallowing is a practice still in force, because of the relative availability of lands. But the land pressure, which is growing with the flow of migrants, tends to make it disappear.

III. HISTORICAL DEVELOPMENT OF VOLTA BASIN

III.1 Changes in water supply and use

III.1.1 Water Institutions and Legislation

Institutional structures and legal frameworks have been established to some degree in the riparian countries for environmental management, as documented in the various country reports. A summary of the various national institutional structures and legal frameworks is presented in the Tables below. The institutions charged with transboundary water resources issues are then discussed in greater detail. A short summary of legal and institutional constraints follows.

Benin

Benin has developed a number of laws and institutions to address the environmental impacts of activities in the country, which are outlined in the following Table 58.

Table 58: Ministries and Departments for managing water and Land resources in Benin

Ministry	Departments	Responsibilities
Ministry of Environment, Settlements, and Urban Development	Environment, Sanitation, and Urban Roads, Administration of Territories	Management of the Environment
Ministry of Agriculture, Livestock, and Fishing	Rural Development, Forest and Natural Resources, Agriculture, Fishing, Livestock	Management of Natural Resources, Water, and Soils
Ministry of Mines, Energy, and Hydraulics	Mining, Beninois Society of Electricity and Water, Hydraulics	Management of Mineral Resources, Management and Distribution of Water Resources at the National Level
Ministry of State in Charge of Coordination of Government Act, Forecasting, and Development	-	Identification of projects and programmes that will have positive impact on the environment. Followup of projects and programmes and their actual impact on the land, and in particular the environment
Ministry of Interior and Security and Decentralization	Department of Territorial Administration; Department of Local community; National Commission of State Affairs; Department of Prevention and Soil Protection	Environmental issues
Ministry of Law and Justice and Human Rights	Department of Law and Codification	Support of the legal Framework
Ministry of Finance and Economy	-	Development of policies for improving the environment, e.g., tax Incentive
Ministry of Public Health	Department of Hygiene and	Implementation of national policies in

Ministry of Higher Education and Scientific Research	Sanitation National University of Benin; Committee of Man and Biosphere; Beninois National Commission of UNESCO; Beninois Center of Scientific Research and Technique	matters of hygiene and health Concern about environmental policies
--	--	---

Apart from the Ministries playing some roles in managing water and land resources, the Beninois government has put in place measures that give roles to the local communities concerning sanitation, public health, and roads. Several legislative texts have been passed for rational management of the natural resources of the country. Some of these include:

1. Decree No. 82-435; December 30, 1982 against bushfire and setting fires to plantations.
2. The Law No. 87-016; Water Code.
3. The Law No. 98-030; February 12, 1999 on legal framework for the environment of the republic of Benin.

Burkina Faso

The overall vision of the country's framework for managing natural resources, expressed in the document "Policies and Strategies in Water Matters" adopted by the Burkinabe government in 1998 is of sustainable human development. This means providing economic security, health, food security, and a sound environment, among other things. The following laws are in existence for managing the environment and land resources:

1. Act No. 005/97/ADP
2. Act No. 0014/96/ADP of 23 May 1996

Table 59 shows the responsibilities of the various government ministries and departments associated with the management of land and water resources in Burkina Faso.

Table 59: Departments for Managing Land and Water Resources in Burkina Faso

Department	Responsibilities
Foreign Affairs	To implement framework of agreements of international cooperation
Administration of Territories/Lands	Land administration
Energy and Mines	Production of hydropower and utilization of mineral resources
Tertiary and Secondary Education and Scientific Research	Education and training
Public Works, Settlements and Urban Development	Road infrastructure and urban development
Agriculture	Enhance irrigation development
Animal Resources	Management of pastoral zones
Health	Public Health
Transport and Tourism	Collection of climatological data
Social Action and the Family	Management of risk linked with
Authorities for managing valleys	AMVS, MOB

Other actors associated with the land and water resources management include NGOs, the private sector, and development partners. Some difficulties encountered in institutional management include sectoral management of natural resources and inadequate management of human and financial resources.

Côte d’Ivoire

In Côte d’Ivoire, numerous institutions are charged with the responsibility of managing and using water resources. This situation has led to fragmentation and dispersion of functions among the institutions. The functions of main actors in water resources are divided into two main groups: managers and users. The law that created this division was Law No. 98 – 755. There are two codes regulating, protecting, and guiding the use of water: environment and water codes. The ministry with the overall responsibility for managing water resources is the Ministry of Water and Forests. It is the authority in charge of policies for managing water resources. Additional ministries include:

Table 60: *Ministries and Their Responsibilities in Côte d’Ivoire*

Ministry	Responsibilities
Resources	
Ministry of Mines and Energy	Generation of hydroelectricity
Ministry of Transport	River and maritime transport
Ministry of Construction and Urban Development	Sanitation
Ministry of Environment and Life	Protection of water
Ministry of Economy and Finances	Financing of water projects
Ministry of Planning and Development	Scheduling of projects
Ministry of Public Health	Protection against diseases associated with water
Ministry of Industry	Industry
Ministry of Tourism	Tourism
Ministry of Interior and Decentralization	

Table 61: *Ministries for the Management and Use of Land Resources in Côte d’Ivoire*

Ministry	Responsibilities
Ministry of Water and Forests	Protection of soils and fight against desertification
Ministry of Environment and Life	Protection of Ecosystems
Ministry of Construction and Urban Development	Management of urban areas
Ministry of Agriculture and Animal Resources	Agricultural development and management of rural areas
Ministry of Mines and Energy	Mineral exploitation

A legal framework that came into force in December 1998, the Rural Land Code, allows for the registration and security of rural land.

Ghana

Policy framework

Details of the Ghana water policy are presented here to illustrate the positive steps being

taken in the basin to create a policy environment that promotes sustainable water resources management.

Water vision: Water is essential to the man and all living things. Health, nutrition and food production, for example depend on its availability in adequate quantities and good quality. Population increase and concentration, rapid urbanization and industrialization resulting in an increase in individual and collective needs have made water increasingly scarce and often of low or reduced quality. In addition to these, current accelerating climatic change processes are expected to increase both the spatial and temporal unpredictability of water availability. The water resource base is, therefore, under increasing threat. Ghana's Water Vision for 2025 has the main objective which to *'promote an efficient and effective management system and environmentally sound development of all water resources in Ghana'*.

Policy Objectives: The water policy of Ghana aims at achieving an efficient and effective management system for the sustainable development of water resources in Ghana to assure full socio-economic benefits for present and future generations.

This would be achieved by ensuring:

- Availability of water in adequate quantities and quality to sustain nature, biodiversity and the aquatic ecosystem.
- Access to safe drinking water and sanitation facilities for the entire population, both rural and urban.
- Availability of water in sufficient quantities for cultivation of food crops, watering of livestock and sustainable freshwater fisheries to ensure sustainable food security for the country.
- Availability of water for hydropower generation, industrial use, water transport, and recreation.
- An effective management system for sustainable use of water and fully integrated into the socio-economic development of the country and national development planning.

Guiding Principles for Ghana Water Policy

On the basis of national aspirations and convictions, as indicated in the Constitution of the Fourth Republic, the National Development Framework, Ghana Water Vision 2025 and the National Environmental Action Plan, as well as other international guidelines, agreements and conventions, Ghana's water policy will be guided by the following principles:

- The principle of fundamental right of all people to safe and adequate water to meet basic human needs.
- The principle of recognising water as a finite and vulnerable resource, given its multiple uses.

- The principle of integrating water resources management and development with environmental management in order to ensure the sustainability of water resources in both quantity and quality.
- The precautionary principle that seeks to minimize activities that have the potential to negatively affect the integrity of all water resources.
- The principle of coordinating water resources planning with land use planning.
- The principle of adopting the river/drainage basin as a planning unit.
- The principle of polluter pays, to serve as a disincentive to uncontrolled discharge of pollutants into the environment.
- The principle of subsidiarity in order to ensure participatory decision-making at the lowest appropriate level in the community.
- The principle of solidarity, expressing profound human companionship for common problems related to water.
- The principle of recognizing the economic value of both raw and value added water and making users bear the appropriate cost.
- The principle that international co-operation is essential for sustainable development of shared basins.
- The principle of integrating river basin management with management of the coastal zone.
- The principle of the greatest common good to society in prioritizing conflicting uses of water.

Acts establishing new institutions and strengthening existing ones for managing water resources in Ghana, and in particular the Volta Basin are as follows:

. Act 46 of 1961 (Volta River Development Act) sets up the Volta River Authority (VRA).

The VRA has the mandate to plan, execute and manage development of the Volta River. The primary function of the VRA is to generate power for the country's industrialization.

. Act 490 of 1994 establishes the Environmental Protection Agency.

. Act 522 of 1996 establishes the Water Resources Commission.

. Act 462 of 1993 establishes the District Assemblies.

Government departments and agencies charged with responsibilities of usage or management of water resources are presented in Table 62.

Table 62: *Ministries and Departments Responsible for Water Resources Development and Use in Ghana*

Ministry	Department/Institutions	Responsibilities
Ministry of Works and	Water Resources *	Planning and regulation of the

Housing (MWH)	Commission	development and use of freshwater resources in Ghana
Ministry of Environment and Science (MES)	Environmental Protection Agency (EPA) *	Management of the country's environment, collaborating with relevant state institutions and international bodies in ensuring sustainable development of the country's natural resources
Ministry of Lands and Forestry	Forestry Commission *	Control and planning of forestry resources
Ministry of Mines	Mineral Commission*	Granting of mining rights
Ministry of Energy (ME)	Public Utilities Regulatory Commission	Regulate the supply, Transmission, and distribution of treated water
	Volta River Authority (VRA) **	Plan, execute and manage the development of the Volta River for hydropower generation
Ministry of Food and Agriculture	Irrigation Development Authority **	Development of irrigation in the country
Ministry of Works and Housing	Ghana Water Company Ltd. **	Provision of potable water for urban settlement
Ministry of Works and Housing	Community Water and Sanitation **	Provision of potable water for rural communities
Ministry of Roads and Transport	Meteorological Services Division ***	Assessment of data
Ministry of Environment and Science	Water Research Institute of CSIR ***	Assessment of surface and groundwater resources in quantity and quality.

* Organization involved primarily in the regulation of the environment and natural resources.

** Organization involved mainly in the development and use of water resources.

*** Organization involved mainly in the data collection and processing of data/information for water resources management.

Mali

In Mali, the water sector is under the Ministry of Mines, Energy, and Water. Its function is carried out by the National Department of Hydraulics, which was established by the law No. 99-023 of June 11, 1999.

The functions of the department, among other things, include the assessment of potential water resources in the country, supervision of works and appraisal of projects in the water sector, and promotion of sub-regional cooperation in the domain of water resources management.

A number of departments are also involved in managing of water resources. To avoid duplication and harmonize activities, a Committee of Interministerial Coordination of Water Sector and Sanitation was established by decree No. 95-447/PM-RM. The composition of the Committee reflects the ministries involved in water issues:

- . The Ministry in Charge of Hydraulics
- . The Ministry in Charge of Planning

- . The Ministry in Charge of International Cooperation
- . The Ministry in Charge of Public Health
- . The Ministry in Charge of Agriculture
- . The Ministry in Charge of Livestock
- . The Ministry in Charge of Environment
- . The Ministry in Charge of Territorial Administration
- . The Ministry in Charge of Finances
- . The Ministry in Charge of Industry
- . The Ministry in Charge of Cottage Industry

The Ministry in Charge of the Environment is responsible for all issues affecting the environment. However, the management of the environment is shared among ministerial departments, which include Rural Development, Health, Hydraulics, Transport, Urban Development, Industries, Education, Public Works, and Territorial Administration.

A number of laws regulate economic and social activity in order to protect the environment. The Preamble to the Malian Constitution states that the people of Mali must insure the cultural inheritance and environmental protection. Some of the laws and regulations governing the water sector in Mali include:

- (i) The Water Code
- (ii) National Policy on Water
- (iii) The Code on Decentralised Territorial Communities and
- (iv) National Strategy on Development of Potable Water Supply and Sanitation.

Other laws governing the environmental sector include Law No. 91-047/an-rm and the Law No. 89-6/an-rm. Other laws govern land ownership and the management of forest resources.

4.6 Togo

In Togo, many institutions are involved in managing of water and soils. The various ministries and departments involved are presented in Table 63.

Table 63: Ministries, Departments, and Institutions Responsible for the Management of Water in Togo

Ministry	Department/Institution	Responsibility
Ministry of Equipment, Mines, and Hydraulic Resources	General Department of Hydraulics	Implementation of programmes, formulation of laws and regulations on respect to water resources and sanitation
Ministry of Public Health	Division of Public Health and Sanitary Engineering	Public hygiene and sanitation
Ministry of Public Health	National Institute of Hygiene	Analysis of water
Ministry of Agriculture, Animal Husbandry, and Fishing	Department of Managing Rural Equipment	Management of surface water, Agro land laws
Ministry of Environment and Forest Resources	Department of General Ecology and Rehabilitation of the	Control of withdrawal of water from water courses, aquifers, lagoons, and the sea for industrial and agricultural

Ministry of Agriculture, Animal Husbandry, and Fishing	Environment Togolese Institute of Agricultural Research (ITRA)	purposes Conservation, studies, and mapping out of soil types
Ministry of Planning and Management of Territories, Habitat, and Urban Development	Department of Urban Development and Habitat	Control of the management of urban lands
Ministry of National Education, University of Lome		
Ministry of Cooperation and Foreign Affairs		

The legal framework governing the management of land and water resources is the Code for the Environment, the decree of 5 February 1933, and Code for Water, which is to be finalized under the management of water resources, and the Mining Code.

Overview of national institutional and legal framework for integrated management

In the riparian countries, many institutions are responsible of managing water and soil resources. This results in the overlapping of responsibilities and difficulties in coordination. Coordination of activities among the institutions was found to be generally weak, and in some cases is only on an *ad hoc* basis for crisis situations. In order for the management of water and soil resources to be effective, it should be integrated at the local and national level, with emphasis on intersectoral coordination.

The effectiveness of the laws governing resources constitutes another problem as the laws and regulations established for managing water and soil resources appear to be weak and ineffective. In some instances, the laws are adequate but they are not adhered to or enforced either due to lack of institutional capacity or political commitment. The knowledge base of the state of natural resources, rate of depletion, and consequent future impact is poor, and probably contributes to the weak political commitment on the part of governments and general apathy on the part of the populace.

- **Regional Coordination**

Several initiatives have been undertaken at the regional level to manage water resources. One such process initiated by the Government of Burkina Faso, with the support of DANIDA (Danish International Development Agency), brought together official delegations from 16 West African countries to form the West African Regional Action Plan for Integrated Water Resources Management (WARAP – IWRM). This regional cooperation arrangement, which started in 1997 within the Economic Community of West African States (ECOWAS) has proposed the establishment of a regional structure for coordinating and monitoring the West African Regional Initiatives for Integrated Water Resources Management. Some of the IWRM country initiatives identified for support include:

- GIRE (*IWRM*) in Burkina Faso
- Water resources management in Benin
- WRIS project for water resources monitoring in Ghana

- Establishing the Water Resources Commission in Ghana
- The sub-regional action plan for combating desertification adopted in 1999 by the environment ministers

Another regional cooperation effort for integrated management of water resources is being developed by the Global Water Partnership and its technical group, the West African Technical Advisory Committee (GWP/WATAC). Their aim is to prepare regional Programmes of Action to implement the West African Water Vision for the 21st century.

A sub-regional initiative, Comité Permanent Inter Etats de Lutte Contre la Sécheresse (CILSS), limited to the Sahel region, considers how to fight drought and desertification with a view to promoting food self-sufficiency in the region.

Green Cross International, with its sub-regional head in Burkina Faso, is also undertaking a basin-wide initiative with the objective to develop basin principles, agreements, and management policies in order to promote peace.

Other initiatives in the region include:

- . GLOWA Volta Project on Integrated Assessment of Feedback Mechanism between Climate Land Use, and Hydrology
- . World Bank
- . Agence Francaise de Développement
- . West and Central Africa Action Plan for Abidjan Convention (WACAF)
- . Land-Ocean Interactions in the Coastal Zones (LOICZ Afribasins project)
- . Center for Africa Wetlands (CAW)

Two other regional initiatives have direct bearing on the Volta River Basin environment. The New Partnership for Africa's Development (NEPAD) is a comprehensive integrated framework for the socio-economic development of Africa, and contains a strong environmental component. Also, two basin countries (Côte d'Ivoire and Ghana) participate in the African Process. Implemented through a GEF Medium Sized Project by UNEP, the African Process has developed a series of concrete projects that effectively address problems identified as having adverse effects on the sustainable development of the marine and coastal environment in sub-Saharan Africa. This TDA and the ensuing Strategic Action Programme draw on the interventions developed under these two programs.

Bilateral cooperation also exists among the riparian countries for mitigating some environmental issues and problems. Such cooperation efforts include the Burkina Faso–Ghana Joint Committee for managing the water resources of the Volta Basin; Burkina Faso–Côte d'Ivoire Committee for the development of programmes for integrated management of water and cooperation in matters of the environment and forest, etc. Table 64 provides details of some of the existing bilateral cooperation efforts.

Table 64: Bilateral Cooperation among Riparian Countries

Countries	Areas of Cooperation
Burkina Faso - Côte d'Ivoire	. Demarcation of borders . Cooperation in matters of the environment and forest . Harmonization of geological cartography of border zones . Development of programmes of integrated management of water resources
Burkina Faso - Ghana	. Transhumance and sanitation issues . Finalization of border demarcation . Fight against epidemics . Pipeline project . Creation of joint commission on the management of water in the Volta Basin
Burkina Faso - Benin	. The dam project of Nounbiel . Rehabilitation of transnational highways . Transhumance and sanitary issues . Poaching
Burkina Faso - Togo	. Transhumance and sanitary issues . Demarcation of tripartite boundary . Exchange of experience in matters of soil and water
Burkina Faso - Mali	. Fight against bushfire . Protection of elephants of Gourma and the management of their movement . Fight against desertification . Roads

Regional institutions, such as the Economic Community of West African States (ECOWAS) and the Economic and Monetary Union of West Africa (UEMOA), all have within their purview the promotion of the integrated management of the natural resources of the region for social and economic development. At the moment, though, a coordinated framework for holistic management of the natural resources (water and land resources) and the ecosystem of the Volta Basin for sustainable development does not exist.

• **International Cooperation**

The riparian countries are also involved in a number of international agreements that relate to the protection of the environment in the Volta River Basin. All the six of the Volta countries are parties to the conventions listed in the table on the next page Table 65. Further, Côte d'Ivoire, Ghana and Togo are involved in Tropical Timber Agreement 83 and Tropical Timber Agreement 94. Benin, Mali, and Togo are involved in the Convention on Conservation of Migratory Species. Finally, all the riparian countries except Togo and Ghana are involved in the Basel Convention on Hazardous Waste. None of the countries, however, is involved in the Convention on Environmental Impact Assessment in a Transboundary Context (Espoo Convention).

Table 65: Dates of Ratification of Major International Environmental Conventions

CBD	Ramsar	Biological Diversity	Climate Change	Montreal Protocol	CITES	World Heritage	Desertification
Benin	30/06/94	24/05/00**	30/06/94	01/07/93*	28/02/84*	14/06/82	29/08/96
Burkina	02/09/93	27/10/90**	02/09/93	20/07/89	13/10/89*	02/04/87	26/01/96

Faso Côte d'Ivoire	29/11/94	27/06/96**	29/11/94	05/04/93*	21/11/94*	09/01/81	04/03/97
Ghana	29/08/94	22/06/88**	06/09/95	24/07/89	14/11/75	04/07/75	27/12/96
Mali	29/03/95	25/09/87**	28/12/94	28/10/94*	18/07/94*	05/04/77***	31/10/95
Togo	04/10/94*	04/11/95**	08/03/95	25/02/91	23/10/78	15/04/98***	04/10/95***

* accession

** entry into force

*** acceptance

Surface water development and use

Throughout the Volta River Basin, dams and reservoirs have been created to mobilize water for agricultural, industrial, and electricity-generating purposes. The number of large and small dams continues to increase as population pressure grows. Increasing use of water and probably decreasing precipitation in the region threaten the continued sustainable management of the waters in the basin. Figure 1 shows the location of dams in the Volta Basin.

Several large dams have been constructed all over the Volta River Basin with the primary purpose of generating electricity. The damming of the Volta River at Akosombo has created one of the largest man-made lakes in the world, covering an area of approximately 8500 km². A smaller and shallower impoundment, the Kpong Headpond, covering an area of roughly 38 km² with a storage capacity of 2000 x 10⁶m³, was completed in 1981 when another hydroelectric dam was constructed at Kpong, 20 km downstream of Akosombo.

Benin has a hydroelectric power station on the Oti River with a storage capacity of 350 million m³ and the capacity to produce 15 MW. Additionally, a hydroelectric power station is planned at Pouya (Natitingou) on the Yéripao.

In recent decades there has been a great push in Burkina Faso to increase the number of dams in the Volta River Basin to meet the country's need and demand for fresh water. As a result, there are now estimated between 600 and 1400 dams and lakes with a total storage capacity of 4.7 billion m³ and about 10,484 wells of which 8020 are in good condition. The volume stored annually in these reservoirs is about 2,490 billion m³. On hydro-power, 13 locations have been identified in the country and 10 are located within the Volta Basin. A total of 125.9 GWH/yr is expected to be generated from these hydro-power stations. A recent survey by DIRH indicates that several planned irrigation projects covering a total area of 1045 ha will require about 65.3 million m³.

There is not a reliable set of data on water demand and use prior to 2001, in Burkina Faso. Water demand for various uses was estimated in 2001 by DGH when formulating the Burkina IWRM programme. For domestic water use the following parameters were used:

- Ouagadougou: 651 l/day/pers.
- Bobo-Dioulasso: 50 l/day/pers.
- Urban area: 40 l/day/pers
- Semi-urban area: 30 l/day/pers
- Rural area : 20 l/day/pers

The total demand for domestic water in Burkina Faso is estimated at 103,500,000 m³/yr; 37,000,000 m³ /yr in urban areas, 2,500,000 m³/yr in semi-urban areas and 64,000,000 m³/yr in rural areas.

Water demand for agriculture was estimated using the following parameters:

- large irrigation schemes (double cropping with rice): 20,000 m³/ha/yr
- small irrigation schemes (double cropping rice and vegetables): 15.000 m³/ha/yr
- developed inland valley (supplementary irrigation) 5500 m³/ha/yr
- Vegetable cropping: 8000 m³/ha/yr

Total estimated water demand for irrigation in Burkina is 323,000,000 m³/yr

Table 66: Water demand during year 2000 (km³)

Volta Basin in Burkina	Domestic water demand	irrigation	livestock	Industrial and mines	Hydropower
Mouhoun	34,89	133,17	21,60	1,31	0
Nakanbé	47,93	69,68	24,80	1,31	2 000
Total	82,82	202,85	46,40	2,62	2 000

Source : Programme GIRE/OTEG

Table 67: Projected water demand in Burkina Faso (km³)

years	1990	2000	2010	2020	2025
Domestic and Industrial water	66,6	85,4	105,6	132,3	149,0
Irrigation	43,1	202,9	383,7	554,0	639,1
livestock	37,2	46,4	60,8	77,8	87,9
TOTAL	146,9	334,7	550,1	764,1	876

Source : Programme GIRE/OTEG

Table 68 : Large dams in the Volta Basin of Burkina Faso

	Basin area (km ²)	Storage capacity (km ³)	Lake surface (km ²)	Annual inflow	Evaporation (mm/year)	10 mm Evaporation (km ³)
Bagré	34 000	1 700	255	946	1900-3000	2,55
Kompienga	5 800	2 050	210	631	1900-3000	2,10
Sourou	11 000	360	400		1900-3000	4,00

Ziga	20 800	200	72	315	1900-3000	0,72
NAKANBE Basin	82 000	391	57 791	2 444	1900-3000	578
MOUHOUN Basin	91 000	470	24 685	2 640	1900-3000	247
VOLTA Basin	173 000	861	82 476	5 084		825
Akossombo	398 390	149 000	8 500	36 900	1600-1800	85

Cote d'Ivoire has any major dams in the Volta Basin since her basin is small and is on the border to Ghana. The following minor dams are located in Cote d'Ivoire.

Table 69: Information on Dams in the Volta Basin of Cote d'Ivoire

Name of the Dam	Year	Manager	North	West	Use	Surface of the Basin (Km ²)	Height of the Dike (m)	Storage Capacity (1000 m ³)
Sorobango	1994	Sodepra	8°09	2°43	Livestock	2.50	4.75	30
Kamala	1994	Sodepra	8°24	2°44	Livestock	3.00	5.00	36
Yerekaye	1994	Sodepra	8°21	2°49	Livestock	7.00	4.50	64
Kiendi	1994	Sodepra	8°11	2°42	Livestock	6.00	5.00	73
Poukoube	1994	Sodepra	8°23	2°42	Livestock	6.00	5.00	30
Tambi	1994	Sodepra	8°13	2°35	Livestock	6.00	4.50	37
Borombire	1989	Sodepra	8°44	3°08	Livestock	4.00	4.25	73
Imbie	1988	Sodepra	9°13	2°54	Livestock	5.50	3.90	73
Lankara	1988	Sodepra	9°11	3°02	Livestock	5.00	4.25	73
Niandeqe 2		Sodepra	9°13	2°54	Livestock		5.00	73
Syaledouo	1988	Sodepra	9°03	3°01	Livestock	4.50	4.25	73
Tidio	1980	Prive	9°16	2°57	Livestock		4.50	73
Angai	1988	Sodepra	9°35	3°17	Livestock	4.50	4.25	73
Bikodidouo	1983	Sodepra	9°34	3°04	Livestock	6.00	4.25	73
Bouko	1990	Sodepra	9°28	3°13	Livestock	4.00	4.20	73
Bouna	1979	Sodepra	9°17	2°58	Livestock	6.00	4.00	73
Bromakote	1988	Sodepra	9°21	3°03	Livestock	9.50	4.25	73
Danoa	1990	Sodepra	9°41	3°16	Livestock	7.00	4.25	73
Gnonsiera	1990	Sodepra	9°37	3°04	Livestock	5.00	4.05	73
Kalamon	1988	Sodepra	9°48	3°10	Livestock	7.50	4.25	73
Kodo	1980	Sodepra	9°41	3°18	Livestock	6.00	4.00	73
Kpanzarani	1988	Sodepra	9°25	3°05	Livestock	5.00	4.00	73
Kpoladouo	1988	Sodepra	9°30	3°19	Livestock	5.00	4.25	73
Nambelessi	1988	Sodepra	9°32	3°18	Livestock	5.00	4.20	73
Niamoin	1982	Sodepra	9°37	3°27	Livestock	7.00	4.25	73
Niadegue 1	1982	Sodepra	9°16	2°54	Livestock	6.00	4.25	73
Peko	1983	Sodepra	9°31	3°02	Livestock	5.50	4.25	73
Piri	1991	Sodepra	9°29	3°11	Livestock		5.00	73
Sepedouo	1982	Sodepra	9°40	3°24	Livestock	5.00	3.50	73
Sipe		Sodepra	9°40	3°24	Livestock			73
Sipirition	1983	Sodepra	9°25	2°54	Livestock	5.50	4.25	73
Tchassondouo	1988	Sodepra	9°35	3°25	Livestock	5.50	4.25	73
Timperdouo	1990	Sodepra	9°32	3°11	Livestock	10.00	4.25	73
Didre Douagre	1990	Sodepra	9°43	3°21	Livestock	5.00	4.25	73
Minichio	1990	Sodepra	9°46	3°29	Livestock	6.00	3.90	73
Nankele	1990	Sodepra	9°52	3°23	Livestock	9.00	4.25	73
Nikindjoka	1990	Sodepra	9°43	3°17	Livestock	4.00	4.20	73
Peon	1990	Sodepra	9°45	3°24	Livestock	4.00	4.00	73

Tinkalamon		Sodepra	9°49	3°38	Livestock	5.50	4.20	73
Yalo	1982	Sodepra	9°48	3°24	Livestock	7.00	4.25	73
Boromeredouo	1989	Sodepra	9°59	3°08	Livestock	4.00	4.25	73
Yonodouo			9°59	2°57	Livestock	5.60	4.25	73
Barriera	1982	Sodepra	9°53	3°27	Livestock	6.00	4.25	73
							Total	2,971

In the Volta Basin of Mali, Pont-barrage of Baye is the only significant dam. Togo has the following dams in the Volta Basin.

Table 70: Information on Dams in the Volta Basin of Togo

Dam	Volume (m³)	Uses
Dalwak	10,000,000	Domestic water supply, Irrigation
Tantigon	762,400	Agriculture, animal husbandry, domestic water supply
Namiete	600,000	Domestic water supply, animal husbandry, market garden
Magna	500,000	Domestic water supply, animal husbandry, market garden
Kozah	5,000,000	Domestic water supply, animal husbandry

Although it is believed that there are hundreds of dams in the Volta River Basin, the data on the locations and size of these waterworks are inadequate. Thus, it is difficult to quantify the effects of the dams on the Volta River Basin.

MAJOR DAMS IN THE VOLTA BASIN
BARRAGES PRINCIPAUX DANS LE BASSIN DE LA VOLTA




Figure 25: Location of Dams in the Volta Basin

Water use:

The basin demand for water is an aggregation of the demands from the riparian countries over a period of time. The country demand, in turn, is dependent on the types of economic activities undertaken, as well as the level of the country's development as more advanced economies will demand more water than less advanced ones. Population is also a factor in determining the quantity of water needed for domestic use. Projections for water demand are based on population growth and the activities envisaged to be carried out under the country's development plans.

Table 71 shows the water demand for domestic and industrial activities. These are projected to increase due to the rapid population increase and envisaged industrial expansion, both of which will require an increased use of water. The domestic/industrial water demand for Benin was, however, very high and may be due to planned economic development activities.

Table 71: Domestic/Industrial Water Demand of the Volta River Basin ($\times 10^6 m^3$)

Country	1990	2000	2010	2020	2025
Benin		56	196	336	448
Burkina Faso	67	85	106	132	149
Côte d'Ivoire	-	4	5	12	14
Ghana	82	138	192	272	284
Mali	5	9	13	16	18
Togo	51	68	92	123	145

Table 72 presents water demand data for irrigation in the basin. In Ghana and Benin, the increases expected are quite high. The percentage increases of year 2020 demand over year 2000 are 538% and 706% for Ghana and Benin respectively.

Table 72: Irrigation Water Demand of the Volta River Basin ($\times 10^6 m^3$)

Country	1990	2000	2010	2020	2025
Benin		152	548	1,225	1,600
Burkina Faso	43	203	384	554	639
Côte d'Ivoire	-	19	57	166	276
Ghana	75	565	1,871	3,605	3,733
Mali	126	180	219	291	311
Togo	43	50	91	133	171

The high projections of water demand for irrigation in the basin arise from the fact that rain-fed agriculture is becoming more precarious and less reliable under climate change and the ensuing variable precipitation. Further, the need to produce adequate food for the rising populations is a major concern of the countries in the sub-region.

Table 73 presents information on water demand for livestock production. It is observed again that the demand for water needed for livestock will increase several times by the year 2025 to meet the protein requirements of the basin population and for export.

Table 73: Water Demand for Livestock of the Volta River Basin ($\times 10^6 m^3$)

Country	1990	2000	2010	2020	2025
Benin		40	94	133	175

Burkina Faso	37	46	61	78	88
Cote d'Ivoire	-	1	2	3	3
Ghana	18	26	41	63	67
Mali	4	34	74	123	142
Togo	15	19	22	30	36

The information provided in Table 74 for the total water demand shows drastic increases of 62% to 1221% in year 2020 over year 2000 water demands. The sharp increases are, however, largely driven by the high irrigation water demand projected for the future.

Table 74: Total Consumptive Water Demand of the Volta River Basin ($\times 10^6 m^3$)

Country	1990	2000	2010	2020	2025	% Income 2020/2000
Benin	-	249	838	1,694	2,223	583
Burkina Faso	147	335	550	764	876	128
Cote d'Ivoire	-	24	64	181	293	1,221
Ghana	175	729	2,104	3,940	4,084	424
Mali	136	223	306	430	471	93
Togo	109	137	205	286	351	62

While significantly higher demands have been projected for the near future, current demands are not now being met in most countries. For example, the water resource supply for the Volta Basin in Ghana for 2000 was $245 \times 10^6 m^3$ (WARM, 1998). This implies that for a demand of about $729 \times 10^6 m^3$, only 34% was met. The problem of inability to meet the water consumption demand depends, to a large extent, upon inadequate infrastructure of water supply systems. This means that there are insufficient financial resources to store, treat, and distribute water.

The problem of not being able to meet the consumptive water demand depends, to a large extent, upon inadequate infrastructure of water supply systems. There are insufficient financial resources to store, treat, and distribute water.

Seasonal variations also hinder the ability to supply water on time. For example, in the upper reaches of the catchment, such as in Mali and some sections of Burkina Faso, river flows are not year round and some wells and groundwater boreholes go dry during certain months. Thus, water availability becomes a problem.

The overuse and misuse of water resources in the region also decreases water availability. In the water resources sector, an aspect of over-exploitation in the basin occurs through the excessive pumping of groundwater without due regard to the recharge characteristics of the aquifer system. This leads to lack of water during the dry season when water is scarce as in the drier parts of the basin in or near the Sahel Zone. Groundwater over-exploitation can lead to saltwater intrusion in the southern parts of the basin near the Gulf of Guinea Coast.

The inefficient use of water resources in the region has exacerbated the problem of scarcity. For example, flooding is the most common irrigation practice in the basin. This approach is very inefficient as it results in water losses through evaporation and deep

seepage. More efficient types, such as sprinklers and drip irrigation, may have to be introduced to cut the water use.

Water supply systems for domestic and industrial uses have large transmission losses due to leakages, which could be as high as 50%. The expansion of water supply systems for domestic/industrial use does not always match the water demands. The limitation in expansion is due to unavailable financial resources.

While each of the countries forecasts increased demand for water over the following decades, the trends in water use pattern among some of the riparian countries are quite different. For example, there has been a rapid expansion of irrigation in the last 15 years in Burkina Faso of about 934 %, while Ghana only experienced an expansion of 95m % (Andreini et al., 2002). Ghana on the other hand, plans to expand its hydropower generation by constructing the Bui Dam. Thus Burkina Faso, an upstream, agriculturally-oriented country hopes to develop the country's irrigation potential while Ghana, a downstream, aims to develop the use of hydropower. The trends in water use patterns can potentially generate conflict if the resources are not managed in an integrated fashion.

Although there is insufficient data on water quality degradation, it has been identified as an important issue in the basin. Some causes of water quality degradation include poor farming practices, improper land use, intensive grazing activities of cattle and sheep, and bushfire. Improper application of fertilizer to agricultural land promotes leaching into the waterbodies. These chemicals are transferred downstream into other countries without any possible restriction. Sediment transport across the riparian countries is the major source of degradation of shared water resources.

Discharge from untreated industrial effluents is not significantly present in the basin due to limited industrial activities, but some untreated sewage is discharged into the waters. Also, humans and animals defecating and bathing in rivers and water sources add to the degradation of water quality. Another significant cause of water quality degradation is the introduction of urban waste, particularly from run-off from inland port communities and urban settlements located near rivers banks and reservoirs.

Surface Water and health issues:

Closely related to surface water availability, access and usage is people's health. According to statistical analysis of GLSS data, the major health problems of communities in the Volta basin are malaria, measles, hernia and river blindness. Of these, malaria and river blindness are directly related to water. The incidence of malaria and river blindness in the sample is about 88 percent and 24 percent of the households respectively. This makes malaria the most significant water-related disease in the basin. Data analysis on malaria transmission intensity in the Nouna District in Burkina Faso points to *Anopheles gambiae s. l.* and *Anopheles funestus* as the principal vectors throughout the year, with high transmission intensities (sporozoite rates 5-15%) documented over the main transmission season (June until December), but very low transmission intensity (sporozoite rates <1%) during the dry and hot season (February

to May). There are considerable variations in malaria transmission intensity by village and season.

Groundwater development and use

Mali

Groundwater in most regions in Mali is put to multi-purpose use including domestic water supply, irrigation and watering of cattle. The distribution among types of use varies from one region to another, as the following table shows in Figure 2, but in most cases domestic water supply dominated other uses.

About 55% of the population of the capital city of Bamako uses water from aquifer resources. In the rural areas, groundwater is either the only source or the main source of portable water supply to the people. This source can be in the form of traditional wells (village sump wells, usually shallow with depth between 2 and 10 m), numbering an estimated 800,000 and not necessarily meeting standards e.g. sumps, pools, and other water-holes are often used or modern water sources e.g. drilled wells with pumps, cistern wells, or other modern wells, estimated at 3,000 modern wells and nearly 9,000 pumps.

A 1992 survey of 384 rural centers showed that three have a water supply; 211 are served by modern wells and sumps; 24 are equipped with solar pumps with mini-networks; and 146 (36 percent) are without modern wells. According to a WaterAid country report for Mali, the satisfaction rate for rural area "potable water" is about 49 percent and that access to potable water for most of the Malian population, especially in rural areas, will require a greater use of groundwater particularly in the desert and sub-desert regions (Mopti, Timbuctu, and Gao) of the country

Telmo (2002) determined that 48% of the households in Gouansolo had access to improved water supplies from groundwater aquifers (i.e., borehole pumps). Groundwater withdrawn for industrial use as a percentage of overall abstracted groundwater is 1; and per capita volume of water used in industrial production is 1 m³/p/year (CIA, 2004)




Figure 27: Regional distribution of groundwater use among sectors
 Source: After Mali National Office of Planning, National Direction Plan, 1991

Burkina Faso

The groundwater resource in Burkina Faso is used mainly for domestic water supply particularly in smaller towns and rural areas. Pipe borne systems based on groundwater are in use in 48 small towns and in the city of Bobo Diolasso (500 000 inhabitants), where drinking water supply relies entirely on groundwater, and in the capital Ouagadougou (> 1 Million inhabitants), where groundwater provides 15 % of the drinking water. This small percentage is however not unimportant as it represents a relatively reliable fresh water reserve in dry seasons (Bracken and Mang, 2002). While mechanized groundwater production for urban water supply has developed continuously since the 1980s in Burkina Faso, the country, like the other countries of the Sahel, suffers from lack of surface water because of geographical, geological and climatic conditions. This means that the country is often obliged to make use of groundwater to supply small towns, villages and the countryside with potable water.

In rural parts of the country, only 0.01 percent of Burkinabé have access to water through private taps; only two percent through public standpipes; 46.1 percent through public wells, 41.2 percent through boreholes, and six percent through sources like rivers, streams, and ponds. Only an urban pittance of 25 percent of city-dwelling Burkinabé access water through private taps, 47.7 percent access through public standpipes, 20.9 percent access through wells or boreholes, and 5.7 percent buy their water from vendors.

There is hardly any data on the amount of groundwater being used for agricultural and industrial purposes in Burkina Faso, though it is known that a limited amount from shallow aquifers is used for irrigation, particularly in market gardening and also for watering livestock.

Ghana

In Ghana, groundwater is mostly used in the rural areas and mainly for drinking and other domestic purposes; the use of groundwater for livestock and poultry watering and for irrigation of crops is limited. Generally, groundwater use is determined by the quantity available, the quality and the unavailability of other alternatives.

Due to the low yield of boreholes and the relatively good quality of groundwater compared to surface sources, boreholes in almost all the ten regions of Ghana with the exception of the Greater Accra region are exclusively used to supply water for drinking and other domestic purposes. Presently, about 52% of the rural inhabitants of Ghana have access to potable water mainly from groundwater sources (Gyau-Boakye and Dapaah-Siakwan, 1999). Since 56% of the 19 million Ghanaian population lives in the rural areas (Ghana Statistical Services, 2002), it means that about 30% of Ghanaians (in the rural areas) have access to groundwater as potable water source. This figure could be higher if those using groundwater for drinking purposes in urban areas are considered. **Table 75** gives an indication of the level of regional dependency on groundwater resources for domestic use.

Table 75: Dependency by region on groundwater supply for domestic use (1984)

Region	Regional Population	Source of groundwater supply by population				Groundwater Dependency (%)
		Borehole	Well	Dug-out	Total	
Western	1,157,807	196,582	134,661	17,343	348,586	30.1
Central	1,142,335	146,800	129,502	31,358	207,660	18.2
Greater Accra	1,431,099	9,135	65,063	21,847	96,045	6.7
Eastern	1,680,890	248,232	224,463	80,596	553,291	32.9
Volta	1,211,907	130,609	232,033	36,110	398,752	32.9
Ashanti	2,090,100	381,192	136,332	30,489	548,013	26.2
Brong Ahafo	1,206,608	103,458	146,439	57,187	307,084	25.5
Northern	1,164,583	23,203	207,009	100,613	330,825	28.4
Upper West	438,008	298,482	7,421	20,272	326,175	74.5
Upper East	772,744	491,177	86,982	15,206	593,365	76.8

Source: After Gyau-Boakye and Dapaah-Siakwan (1999)

III.2 Agrarian and environmental transformations

A. Technical change (agriculture, water mobilization...

1. AGRICULTURE

1.1 Agricultural changes in the pre-colonial era of Ghana

Emerging from the prehistoric era, men generally ceased to depend, as they had done, on wild fruits and root crops they had gathered and wild animals they had hunted. Instead they began to cultivate crops on farms. Ghana shared this experience. There is no exact knowledge about farming in Ghana in the very early days. However, archaeology, botany,

and the study of languages have provided some knowledge about the crops people cultivated and the methods farmers used. Ghana had three main different geographical zones: the savannah in the northern zone, the sandy swampy mangrove lands on the coastal belt, and in between these two zones, the forest belt. Some crops require much rainfall, others need ample sunshine, and other grow in zones with a combination of these climatic conditions. The crops grown in the above three zones have, until recently, depended very much on the nature of the land and other geographical conditions, including the amount and duration of annual rainfall.

The crops, which were cultivated in Ghana before the advent of the Europeans, were not as varied as they became later, following the introduction of new crops to the country by the white men. These foodstuffs include some species of cereals like millet (or guinea corn), yam, pepper, beans and other vegetables. The Portuguese records also indicated that a specie of rice was the staple crop in the country lying between the Gambia and the Cacheu River. Although there are no existing records to indicate that this cereal was in cultivated in Ghana, it was not unlikely that the country grew this crop at the time of the arrival of the white men. The surplus of their produce was sold to the growing populations around the European forts, and to the Europeans both for their own consumption and for replenishing the provisions of the vessels passing over West Africa on their way to the Far East.

When the early Europeans became firmly established on the Guinea Coast, they introduced tropical crops from other lands in the New World and in the Far East. Some of the important foreign crops introduced into Ghana and other parts of the Guinea Coast were cassava, pineapple, orange, tangerine, avocado pear, guava, sugar cane and coconut. These crops were introduced to ensure increased regular supplies of these imported crops for the consumption of the European merchants and for the crew en route to the East. Later, when they were widely cultivated, the new crops introduced a diversified diet to the people.

Before mechanized farming was introduced to the country in more recent years, the people of Ghana, as happened in most other countries at the time, employed shifting cultivation method. The implements used were mainly the cutlass and the hoe. In the absence of labour-saving devices, farming from planting and sowing to the harvesting, was all done by physical human labour. In consequence, the size of farms was generally small. For a very long time farmers mostly engaged in what is called subsistence cultivation, only growing enough for their own domestic need, with very little left for the market

With the increasing concentration of population in the growing towns, many city dwellers cut themselves off from the land. These people depended on the surplus farm products of the countryside. To meet the demand of the market, farmers expanded their farms and increased their yields and wealth. Following the abolition of the Atlantic slave trade,

more and more emphasis was placed on commercial agriculture in Ghana and other parts of West Africa. In Europe the transformation of industrial processes called the Industrial Revolution created a great demand not only for West African palm oil, but also for other tropical crops, like raw cotton, rubber, etc. From the second half of the 19th century onwards, many farmers changed to plantation farming, producing cocoa and coffee, which steadily mainly sustained the economy of Ghana.

1. Agricultural changes in the Volta Basin of Ghana

Long before Akosombo was constructed, the fertile banks along the Volta River were some of Ghana's richest agricultural land. Archeological findings show that the Volta Basin was once well populated (comments from Kaplan et. al., 1907). Much of the natural vegetation was burned down for agriculture over a period of more than a thousand years, which led to the eventual drying and erosion of the land. However, the floodplains along the Volta River provided a constant source of fertile agricultural land for local farmers.

Before the Akosombo Dam, local farming along the Volta was structured around the rise and fall of the river. The damming put an end to natural cycles, which deposited nutrient-laden silts along the flood plains. The river ecosystem was transformed into a lake ecosystem. Damming led to a drastic curtailment in subsistence agricultural production and animal grazing.

An important development in agriculture associated with the Akosombo Lake is farming on the land exposed by fluctuation in the lake level. This development has started on the initiative of the local lakeside dwellers, but research work has been focused mainly on the soils and cropping patterns. The annual fluctuation of the lake varies between 1.8 and 4.3m. It has been estimated that, at 3.0 m some 80,940 hectares would be exposed around the 4,828 km of the lake boundary. The soils in the drawdown area are mostly sandy and well drained. The eastern and southern banks are steep and expose only a narrow area. Wider stretches occur on the northern bank where population density is low.

The northern bank of the Afram and the western bank of the main lake also exposes moderately wide areas. According to various sources, about two-thirds of the periodically flooded land areas are suitable for agricultural use. In 1971, about 10-15% of these areas were under cultivation. In 1979-1980, conflicts with land use started in and around the major centres between the local population and fishermen as a result of the shortage of land in the shore area.

At present only small portions of the drawdown area, those adjacent to the areas with high population density are cropped, mainly with vegetables such as tomato, okra, pepper, garden egg, and maize. Some problems encountered include drought conditions and insect infestation.

Unfortunately, demands for electricity could not always be synchronized with the traditional flooding seasons. Furthermore, the reduced flow into the Gulf of Guinea

resulted in saltwater intrusion at the Volta River delta and estuary. Salt water destroyed clam beds and lowered drinking water quality. Many of the stream and clam fishermen downstream moved north of the lake, where they hoped to restore their career.

The Volta River Project also included plans for an irrigation network in the Afram Plains, which was considered Ghana's agricultural breadbasket. However, these plans were pushed down on the agenda for various economic, social, and political reasons. The possibility of using water from the lake for small, draw-down, lakeside irrigation schemes was further curtailed by the refusal of the VRA to release water from the lake and thus prejudiced the full production of power for the smelter and other consumers.

The Sahel drought did little to improve the prospects of such irrigation schemes. The sudden change in government in 1966 led to budget restructuring during the first year of Akosombo's operation, which affected many of the original project initiatives, including the irrigation network and the resettlement scheme. Teaching new farming techniques was expensive, and there was no guarantee that the traditional farmers would embrace them. In 1998, only 2,000 hectares of irrigated land existed in Ghana.

People who lived in flooded area were mostly subsistence farmers although some of them grew cocoa (2400 ha of cocoa were lost under the lake) and some were river fishermen. In general their standard of living was low. The typical farmer had 2.5 to 4 hectares of land under his control and 3 or 4 plots which he would use in turn in successive years. Only about 6% of the land area covered by the lake was used productively. The rest was unsuitable and unoccupied.

The main agricultural objective of the VRP (Volta River Project) was to improve the agricultural production systems to enable people to move from subsistence to a cash economy. To maintain the fertility of the soils, they choose an intensive mechanizing agriculture. The aim was to specialize agriculture as follows:

- Arable farms >4.9 ha mechanised;
- Tree crops 2 ha < surface < 6 ha was to be provided;
- Intensive livestock 1.2 ha < surface was to be provided;
- Pastoral 12.1 ha < surface was to be provided.

The VRA (Volta Resettlement Authority) had also hoped to introduce new farming methods. VRA controlled the allocation of seeds and fertilizers and the bulk buying of the products. They wanted to clean 41 600 ha for agriculture to provide 4.8 ha for each adult male settler, whereas 51200 hectares were used productively in the flooded area. Mechanized cultivation was to be used across several plots, to give the few farmers the advantage of modern farming methods. The farmers were responsible for weeding and harvesting their fields with the costs for the use of mechanized equipment which is shared on an acreage basis.

The VRA was to own the tractors, distribute and sell fertilizer and buy a significant quantity of the agricultural produce. It was expected that the average farmer would have an annual income of £(G) 350. Moreover, it was expected delivering irrigation water by gravity from the Volta Lake and by pumping from the Volta River, to irrigate about 180,000 hectares. The total area of the Accra plains that is irrigable and suitable for mechanized agriculture was estimated at 178,000 ha out of 335,000 ha of land and the economic potential has been found very high. The total investment required for the irrigation project was by the time about £ (G) 28 million in 10 years.

Annual fluctuation in the level of the lake varies between 1.8 and 4.3 m and as such it has been estimated that at 3m, some 80,940 ha of fertile land would be exposed around the 4828 km of the lake boundary, which could be farmed.

Irrigation opportunity

The Volta Lake represents a good opportunity for developing irrigation schemes in the Volta basin. According to many, about 178, 068 hectares out of 335,092 hectares of the Accra Plains could be irrigated by gravity from the Volta Lake and by pumping from the Volta River. A scheme had been constructed including the use of mechanized agriculture methods, fertilizer and pesticides. The types and properties of the soils were determined, and the socio-economics of the area found to be very high. Since the population of the Accra-Tema commercial region was growing very fast, the demand for food was also growing rapidly. However, the scheme appeared inappropriate and too expensive (£28 million). No moves have been made to implement it, so that the idea of irrigating the Accra Plains remains in abeyance.

VLR&DP successfully carried out drawdown demonstration farming complemented with sprinkler irrigation water from the Lake by using of small pumps at VRA's resettlement township of Ampem. People living in the whole of the Afram arm of the Volta Lake have broadly adopted this method of farming. The Afram area has thus become important for the production of tomatoes and other short maturing crops during the dry months from November to about May.

Kpong Farms Limited was created by the VRA in order to promote leadership in agricultural systems, irrigation practices and food processing technologies. The objective of the authority was mainly to prove that this venture (large scale irrigation) is economically viable and can be adopted to cover Accra Plains. The project which uses water from the Kpong headpond was initiated with the construction of a mainly gravity irrigated scheme covering an area of approximately 100 ha for the production of beans, rice, etc.

It has now developed into a full integrated agro-industry encompassing rice, livestock and pasture production as well as feed milling, abattoir services and meat processing. VRA hopes that the positive demonstration at the drawdown areas and the Kpong Farms Limited will encourage private entrepreneurs to move into irrigation farming.

Government has taken the initial step to irrigate the Asutuare lands for farmers using

water from the Kpong headpond.

II. WATER MOBILIZATION

Since the beginning of the 20th century, water development projects have been planned to make use of water and other natural resources in the basin. Thus, the idea of damming the Volta River began in the early 1900s with the conception of the Volta River Project (VRP) and ended with the creation of Volta River Authority (VRA) in charge of the creation and management of the Akosombo dam and later, the Kpong dam downstream.

The Akosombo dam is by far the most significant structure built in the basin and the Volta Lake is the largest man-made lake in the World. It is of strategic importance to the economy of Ghana. It generates 80% of the power produced in the country. The primary purpose of the project was to supply cheap electricity to smelt aluminium and the secondary one, the development of the country. Construction of the Volta Lake led to the resettlement of about 80,000 people from several hundreds of villages to fifty newly built townships (more than 1% of Ghana's population at that time). In addition to the resettlement of the river communities, damming affected local health, agriculture, fishing, and navigation.

In other riparian countries of the basin, small and large dams have been built by governments, NGOs and local people after the severe droughts that occurred in the 1970s and 1980s to secure food production. In the Nakambe sub-basin (Burkina Faso) alone more than 400 dams have been built most of which during that period. More recently power generating dams have also been built in some of the Volta main tributaries Bagre and Kompienga (Burkina) with generating capacities of 41.5 GWH and 31.0 GWH respectively and on the Oti River, at the border between Togo and Benin within a power generating capacity of 35GWH

***Animal production in the Volta Basin of Ghana**

1. Fisheries

After the formation of the lake, fishery has become an important branch of the economy in its employment and revenue potential. Contrary to expectations, the fish catch very rapidly exceeded 20,000 metric tons. A detailed study carried out in 1969-1970 located over 950 fishing villages, mostly newly established, along the shore of the lake. From these villages, some 12500 fishing canoes and approximately 20,000 fishermen were operating, and the total population of fishing communities was estimated at approximately 59,000 people in 1971. This number was estimated to be about 87,000 in 1975.

From May 1969 to April 1970 a statistical survey of the fish catch of these fishermen gave a point estimate of 61,000 metric tons fresh weight equivalent. The catch settled down to a level of 38,000 tonnes the following year, when the lake became

ecologically more stable. Before the construction of the dam, the annual take from the Volta Basin was of 10,000 tonnes. There had been 2,000 fishermen operating in the basin but now they are estimated to be 12,000 working on the lake. They and their families make up fishing communities, numbering about 59,000 in 1971, living in approximately 1,275 settlements around the lake (1,479 in 1975). Some 87 % of these persons were migrants and were already familiar with the techniques of fishing.

They were attracted by the excellent fishing grounds in the lake but also by the changes that occurred in the lower reaches of the river following the construction of the dam. 60 % of the households migrated from the new settlements and 8 % of them derived their income mainly from fishing in 1971. About 59.8 % of the settlements were located in the Southern area of the lake (46.1 % of the total shoreline), and only 55.4 % in 1975 because of the saturation of the free lands.

The selling price was about ¢0.25/kilogram (that is the fishermen received) so that the value of a year's catches is about ¢9.5 million (the total revenue obtained from Akosombo- produced electricity is of ¢28 million). The total marine catch was estimated as 119000 tonnes in 1969. The Ghana's total consumption of fish in 1969 was 197, 000 tonnes (18,700 tonnes imported at a cost of ¢5.8 million). The catch from the Volta Lake therefore satisfied approximately 20 % of the country's demand for fish. Most of the fishermen working in the Volta Lake operated elsewhere in Ghana before the Lake was created. A large proportion of them were engaged in clam and creek fishing on the lower Volta (Article 6). The creek fisheries were worth about £65,000 a year and the clam fishing no less than £25,000 a year. At the 1955 prices the loss was about ¢160,000 or at 1974 prices about ¢640,000.

About 62 % of the catch is consumed before being marketed and only 2 % ends on the market as fresh fish. The consumers' centres are not located in the immediate vicinity of the fishing grounds. The yields from the fishery in the Volta Lake cover about 15 % of the annual fish consumption (1974) and 29 % has to be imported. In 1977 the revenue from power produced by Akosombo Dam was ¢33.474 million that is 15 % less than the value of the annual fish production (¢38.719 million).

E. Health and Environmental Changes in the Volta Basin of Ghana

I. HEALTH

There have been serious health issues associated with the Volta Lake, in particular the two major diseases schistosomiasis (or bilharziasis) and onchocerciasis (or river blindness).

The dam virtually halted the rate of flow in the Volta River, increasing stagnant water

conditions and consequently, creating ideal breeding grounds for carriers of waterborne diseases. Before the Akosombo and Kpong Dams, malaria (from mosquitoes) was not much of a problem along the swift-flowing Volta River, but after it became a stagnant lake, malaria became a greater public health concern in lakeside villages. Likewise, only 1 - 5% of the population suffered from *schistosomiasis* (a disease transmitted by snails) before the dam was constructed. By 1979, urinary *schistosomiasis* had grown to become the most prevalent disease in the area, affecting some 75% of lakeside residents (comment from Gitlitz, 1993).

Volta Lake has had significant effects on public health situation. *Bilharzia* reached a prevalence rate 90% among children in certain localities. The lake has flooded out the riverine forests which constitute a breeding place for a specie of tsetse fly. The lake has also wiped out the breeding places of the vector of *onchocerciasis*. About 60,000 fishermen living mostly in isolated villages around the lake were exposed to the riverine and did not have access to health facilities.

f **Prevailing diseases before the Akosombo Dam Construction**

Among the many medical problems found in the area to be flooded, were the usual worm infections, particularly round worm, hookworm, yaw, leprosy, tuberculosis, malaria, cerebrospinal meningitis, smallpox and malnutrition and poliomyelitis in children. Before the creation of the Volta Lake, infection rates of *Schistosomiasis* in the area had been 1 to 5% . In the Asukwakwaw area, north of the Akosombo dam, the prevalence of *Onchocerciasis* could reach 90%.

f **Post construction period**

In 1968, the public health aspect of the Volta River Project was assigned to the UNDP (United Nations Development Program) assisted by VRR&DP of the VRA. It was to research into the public health situation and resettlement of the people displaced by the Volta Lake. Attention was paid particularly to the water-borne diseases (*Onchocerciasis*, *Schistosomiasis*, *Trypanosomiasis* and *Malaria*). The immediate objective was the control of the major parasitic diseases.

Between 1968 and 1977, much valuable health data were collected. The changing of familiar houses and environment to new villages with core houses and limited room space caused overcrowding and discomfort and created conditions for the spread of communicable diseases such as yaws, measles and smallpox. Lack of adequate farmlands and community life led to economic hardship, starvation and malnutrition for the children.

The communal latrines provided soon ceased to function due to lack of improper use and maintenance. The pipe borne water systems operated by diesel pumps soon fail, etc. This general low standard of living is usually associated with the parasitic

diseases such as malaria, dysentery, typhoid fever, hookworm and other intestinal worm diseases found in the resettlements. Yellow fever and dengue were the most significant arthropod-borne viral infections found during the early part of the resettlement program.

Increase in development of aquatic snail, host of *Bilharzia (urinary Schistosomiasis)* led to a great increase in the prevalence of the disease in many localities around the lake. Transmission occurred in the lake itself. Man is infected with urinary *Schistosomiasis* from water in which swim larval forms (called cercariae) of the parasite worm dwell. *Schistosomiasis* is a cumulative and debilitating illness. Surveys in 1964-1977 showed a 90 % prevalence rate in school children in fisher's population who live close the lake. The problem of *Bilharzia* in the lake basin must be seen as a disease, which embraces both the lake and the Volta delta. The migratory habits of the fishermen ensure the spread of the disease from endemic areas to other areas.

Onchocerciasis is transmitted from man to man by the bite of the black fly, which breeds in rapidly flowing streams and rivers. When the lake was formed, the major breeding sites north of the Akosombo dam were inundated and eliminated. The construction of the second dam at Kpong also eliminated the breeding sites downstream of Akosombo and therefore stopped the transmission of the disease in the vicinity. However, in the Asukwakwaw area, north of the Akosombo dam, the incidence of *Onchocerciasis* is still high because the tributaries of the Volta Lake and the seasonal streams continue to support the breeding of the black fly.

The 20-year *Onchocerciasis* Eradication Programme started in March 1974. The programme was a combined effort of seven West African countries (Ghana, Togo, Niger, Burkina-Faso, Benin, Mali and Ivory Coast). It has achieved much success using modern techniques and pesticides. The discovery of *Evermectin*® for mass treatment, has increased the hope for a total eradication of the disease in affected areas. The benefit to health of the construction of the Akosombo dam in 1965 and the Kpong dam in 1981 is undoubtedly the reduction of the incidence of *Onchocerciasis* in the Volta Basin.

The amount spent to combat these diseases is not well known. The health program has been part of the Volta Lake Research and Development Project, initiated in January 1968. This project was built to help develop fisheries on the lake, to research into the use of drawdown area for agriculture, to carry out socio-economic surveys of the resettlement villages and research studies on the public health aspects of the Volta Lake. This project was the result of cooperation between VLR&DP, UNDP/FAO and the VRA/Ghana Government.

What have been the effects of these diseases in terms of lost productivity? Very little literature has been produced on the subject of *Onchocerciasis*. Moreover, few reliable statistics are available on the prevalence of the disease in Ghana and for many public health specialists, it seems that Akosombo dam has caused no overall change in the number of people affected by *Onchocerciasis*. More work has been done on the subject of *Schistosomiasis*, both in Ghana and internationally. Several attempts have been made to estimate the cost of *Schistosomiasis* in terms of loss of productivity. A study in the Philippines (Farooq, 1963) divided the infected people into several categories:

Table 76: Categories of infected people according to Farooq, 1963.

Category	Loss of working capacity (%)	Proportion (%)
1) No manifest symptoms	0	61.5
2) Mild symptoms, no absence from work	25	22
3) Moderate symptoms, reduced capacity of work	50	15
4) Severe symptoms, frequent absence from work	75	1.5
5) Very severe symptoms, total absence from work	100	

It is possible to apply the scheme to the Volta Basin. Taking a low estimate of 100 000 as the number of people living around the Lake Volta, and using the average infection rate of 80% we may deduce that 80,000 people are infected.

Thus,

- 22% (or 17600) have their working capacity reduced by 25%
- 15% (or 12000) have their working capacity reduced by 50%
- 1.5 (or 1200) have their working capacity reduced by 87.5%

Ghana's GNP per capita in 1975 was \$590. The value of lost productivity per year was therefore $\$590 \times (0.25 \times 17600 + 0.5 \times 12000 + 0.875 \times 1200) = \$6\,750\,000$

During the construction of Kpong Dam in the early 1980s, flooding provided some health benefits. The dam was sited a few kilometers downstream from Kpong town, so that the backwaters would flood the Kpong rapids. This area was the largest breeding ground for the tsetse fly in Ghana. Sleeping sickness, carried by the tsetse fly, was a major problem to the British colonists, foreigners, and other people who did not have acquired biological defense. Tsetse flies breed in the dense bush bordering bodies of water, and are most prevalent in the lower part of the Brong Ahafo Region, the Ashanti Region, and parts of the Eastern Region into the Volta Region. Sleeping sickness may linger quietly in a person for many years, causing a loss of energy and reduced immunity to other diseases. Full-blown sleeping sickness leads to quick death.

II. ENVIRONMENTAL CHANGES

The impact of the Volta Lake on the environment has been found to be varied. These include enhanced fishing upstream and diminished fishing downstream, opportunities for irrigated farming downstream, proliferation of aquatic weeds upstream and downstream, increase in some water borne diseases such as bilharzia and malaria and reduction and elimination of other diseases such as Onchocerciasis in some areas. Also included seismicity, sediment load changes, morphological changes and microclimatic changes. During the pre-impoundment period a host of multi-disciplinary research workers from academic and specialist institutions studied the impact of the reservoir on agriculture, fisheries, public health and on conservation of biodiversity

Some studies showed that mineral resources flooded were mainly limestone, gravel and sand, which were of limited economic value. Initially sedimentation was not found to be a problem; but now due to deforestation on the edges of the reservoir some sedimentation is being observed. Meteorological data available shows that the reservoir suffers a significant loss of water through evaporation, which is balanced by gain from rainfall. No significant climate change has been observed around the reservoir except in the southern Afram area where a micro climatic change has been observed in the form of a relative dryness of the area. The Volta Lake has been found to be low in mineral nutrient except during the flood season when the annual inflows raise the level of mineral nutrients. The arms of the lake and the shallow areas, which have more nutrients, have been found to be more productive than the deep open waters.

(i) Deforestation

The activities of people living in uncontrolled human settlements have caused serious deforestation around the lake. It has resulted in intensive erosion leading to siltation and sedimentation, which adversely affected the life span of the Lake. VRA has therefore taken serious measures to avoid eventual silting of the lake. The authority has sponsored two afforestation projects intended to stop erosion around the lake. In the Adjena Gorge, VRA and the Forestry Department have declared the steep areas bordering the lake as protected lands. VRA has funded another pilot afforestation project in the low slope areas around Yeji in the Brong Ahafo region. The objective of this project is to encourage fishermen to plant wood lots and adopt the use of improved stoves, which use less fuel wood for fish smoking. It is hoped that this project would help to restore some 580 hectares of tree cover and reduce the dependency on fuel wood by 60 % in the Yeji area.

(ii) Aquatic weeds

The Volta Lake, like other tropical man made lakes, has been characterized by growth of aquatic weeds (Figure 27). The Volta did not experience that short-lived but sudden explosion of floating weeds, which populated Kariba immediately after impoundment. The weeds, which have occurred, are longer lasting. Among the important ones are Pistia, Vossia spp., Ceratophyllum. Pistia is flood resistant. Vossia spp., which is resistant to wave and wind action, is also resistant to herbicides. It has needle-like hairs, which make it unsuitable neither for grazing nor for manual cutting. It creates a good habitat for fish. Ceratophyllum's submerged beds house large populations of Bulinus snails the vector of Schistosomiasis. Attempts to control Schistosomiasis by controlling Ceratophyllum, which create a favorable habitat for the vector, has failed. The weeds have also provided favourable habitats for many disease vectors, notably the vector of *bilharzias* and that of *yellow fever* virus. The aquatic weed invasion had negatives affects on water transport, power generation, health, agriculture and fisheries.




Figure 27: Water Weed Encroachment

(iii) Sandbars

Before the dam construction, seasonal floods flushed out sandbars, which might have started during the dry season, thus the estuary was largely kept free of sandbars. However, after the construction of the dam, due to absence of annual floods, sandbars gradually formed at the estuary and in time virtually blocked it. The effect of this was that saline water, which during high tide flowed upstream into the river channel, completely ceased.

Moreover, with the absence of salt water into the river channel, soon the areas even closer to the estuary started to experience the presence of the schistome snail and Bilharzia became prevalent. But, by bringing a dredger permanently stationed at the estuary to cut a channel through the sandbar, VRA restored salinity to the river channel. As a result, Bilharzia is now wiped out of these areas.



Figure 28: Sandbar on the Volta River mouth

(iv) Seismicity

Reservoir induced seismicity (RIS) is one impact of the water project which has received attention in recent times because of its potential catastrophic effects. The creation of the Volta lake has added weight to the underlying rocks amounting to 165 million metric tons when the lake reaches its estimated capacity of 165 km³, and fluctuating through a range of 25 million metric tons during normal fluctuation of the lake level (Smithsonian Institute, 1974). Significantly, the weight build-up has occurred over an area, which is not very far from the seismically active area along the main Akwapim fault.

In November 1964, when about 28km³ of water had been impounded, an earthquake of an intensity of 5 on the modified Mercalli scale occurred with its epicentre located near Koforidua, about 40km south of the reservoir. In December 1966, another earthquake occurred when the total impoundment stood at 102km³, and a third occurred in February 1969 with storage at 162 km³, but the epicenters for these shocks were offshore from Accra and more remote from the lake. These incidents led Kumi (1973) to conclude that at least the 1964 incident, which had its epicenter within 40km of the reservoir, could be attributed to crustal readjustments associated with the initial stages of the lake's infilling.

(v) Sediment load changes

Before the construction of the Akosombo dam, the highest measurements of suspended sediment concentration of the Volta River at Ajena in 1956, at the height of the flood season, did not exceed 90mg/l near the surface (Volta River Project (VRP) Preparatory commission, 1956). Current measurements downstream of the dam at Tefle (Sogakope) by the Water Resources Research Institute (WRI) show that the average suspended sediment concentration is 38mg/l. This significant reduction in sediment load

downstream has been linked to increased channel-bed scouring and exposure of the piles on which the supporting piers of the Tefle bridge stand

(vi) Morphological changes

Some profound morphological changes are found along the stretch of the Volta river downstream of Akuse and in its delta zone. Since 1974, the point at which the Volta entered the sea is known to have shifted 12km eastwards in a channel, which runs parallel to the coast. This phenomenon of gradual eastwards shift of the mouth of the Volta River has been known in the past but the process seems to have been accelerated since the construction of the Akosombo dam and the flood-flushing of the river mouth stopped. The main concern with these changes has been the disastrous effect any rapid flood release could have on estuarine communities and the environment. It is largely due to the adverse changes that have occurred in the delta zone, which periodically prompts the VRA to artificially re-establish a direct flow into the sea. Also, the changes are probably linked to accelerated coastal erosion at Ada.

(vii) Microclimatic changes (weather changes)

The transformation of bare or vegetated land into water body causes significant changes in surface albedo and thereby alters the local heat budget. It also causes changes in surface roughness, which may tend to affect wind speed. Oral accounts of sudden change in wind speed on the Volta Lake and surrounding areas have been documented by Moxon (1984). Similar reports of observed changes in local weather conditions were corroborated by the residents during field survey of the impact of water projects in Ghana by the personnel of Water Resources Research Institute (Sam, 1993). The main scientific study of the impact of the Volta Lake on the local climate was pioneered by De-HeerAmissah (1969). He studied available climatic data from near and distant stations from the lake, such as Tamale and Kete Krachi in the northeast, and Akuse and Accra in the south, two years after the lake started filling up, and arrived at these conclusions:

(i) There was no evidence of significant changes in the monthly rainfall amounts at the four stations even though a tendency toward decreasing rainfall amounts at Kete Krachi and increasing amounts at Tamale was becoming evident.

(ii) The mean monthly maximum temperature since 1964 at Kete Krachi seemed to be lower than the pre-project 10-year mean, whereas the minimum temperatures were higher than the 10 year mean.

A recent work by Opoku-Ankomah (1996) on the rainfall pattern of Accra from 1935 to 1995 divided into decades, i.e. 1936–1945, 1946–1955, etc. showed that there was a gradual rise of rainfall total from the beginning of 1936–1945 to the 1956–1965 decade and a decline thereafter till 1986–1995. The mean annual rainfall for the 1986–1995 decade has not been high, however, the number of occurrences of very large (24-h

maximum) rainfall events have been relatively high

III. LAND TENURE SYSTEM AND AGRICULTURAL POLICY CHANGES IN BURKINA FASO

1. Changes in the land tenure system in Burkina Faso

According to Ouedraogo N., 1997, three land systems which coexisted in Burkina Faso can be distinguished. They are as follows: customary land system, colonial land system and post-colonial land system.

(i) The customary land system

The customary land system is almost the same everywhere in Burkina Faso. It is based on the collective ownership of land. The collective ownership of land is exercised by the land custodian (known as Tengsoba for the Mossi, Tarfolo for the Sénoufo, Susunnbaso tinibaso for the Bwaba etc. (Ouedraogo S. 1993). In all customs the land custodian is the closest descendant of the first settler. In this capacity, he administers the land patrimony of the group in the interest of all the community. He distributes land or he authorizes land use, following the indispensable rites, by households and individuals that require it and in accordance with their needs. Thus, the applicant acquires a user right, which must not be mistaken for ownership right in the Western sense of the term. However, after the death of the applicant his heirs will settle and exploit the same land without the land custodian intervening afresh. This land is available for the whole community for any possible use in case no heir claims it. Land is given provisionally to strangers (non-natives) Therefore the right accorded to the stranger is precarious, hence the notion of land insecurity for migrants. In this case, land is simply lent, often following royalties in kind or performances of various services. Security imperatives require that the applicant be first socially integrated. However, except in rare cases, land cannot be refused to an applicant according to customary law. Therefore the customary land system is complex in practice.

At the economic and social levels, it must be acknowledged that the customary land system, with all its utilization nuances, opposes creative investments or dissuades them through the almost permanent insecurity as far as individual use is concerned. In this system, land is not given to those who have the necessary means to develop it, but to those who win the confidence of the social group that owns the land, and moreover, on a provisional basis.

(ii) The colonial land system

This system is essentially based on private ownership while making provision for a public estate. Private estate is acquired through purchase, exchange, gifts or legacies, etc. Public estate is made up of natural properties such as hills, rivers, natural or artificial lakes (roads, artificial water bodies, etc.). This system was resisted by the customary land system.

(iii) The post-colonial land system

The land system in force now is based on the agrarian and land reform (RAF), adopted in 1984, which grants user rights to individuals and moral entities. It makes provision for a national land including all the lands within the borders of Burkina Faso, no matter their former status or legal systems. State property is inalienable, not to be seized and imprescriptible. In addition to ownership right, the State assigned itself that of management. In this way, it defines rights for groups and individuals. It should be pointed out that the settlement and exploitation of rural lands by peasants for subsistence are free.

The RAF so designed shatters the mystic aura of land and takes away from peasants their references and customary value systems by leading them to practice new farming techniques (protection and restoration of soils, fertilization, etc.). This law is particularly aimed at making land available to those willing to develop it, and at organizing the rational management of this resource which has become in the end scarce in Burkina Faso.

Therefore, this land system aims at favoring the development of productive agricultural forces. However, it does not encourage people to make sustainable investments and much resistance to its enforcement can be noticed.

2. Changes in Agricultural Policy in Burkina Faso

Burkina Faso has remained an agricultural economy, with agriculture providing a livelihood for 90% of the population. But the performance of Burkinabé agriculture is strongly constrained by the erratic limitation of land and water and by demographic pressure on the available of land.

Agricultural production in Burkina can be divided into "food crops" and "cash crops." Using 80% of the cultivated area, food crops, which consist of cereals (millet, sorghum, fonio), maize, and rice, are the most important products of the agricultural sector. Cash crops are basically cotton and groundnut. Cash crops are of fundamental importance to Burkina's economy, because they provide the main trade revenue of the country. Cotton in particular has been Burkina's main export commodity and has gained significant importance in recent years.

Since independence, the State has played a major role in the agricultural policy of Burkina Faso. It has devoted between 10% and 25% per annum of public investments to agriculture and achieved many Rural Integrated Development Programs, the most recent being the Valley of the Sourou River project.

The extension agencies of the Ministry of Agriculture have provided services such as training, agricultural technique development, and input distribution. As a seller of inputs and purchaser of harvest, the State has achieved another important regulation aim, through parastatals like SOFITEX in cotton production areas and the OFNACER responsible for cereal stabilization. The state has influenced the agricultural sector as a

policy maker. Agricultural policy changed in the decades preceding adjustment with regimes and their perception of the relative importance of rural versus urban dwellers. Through the tool of producer price setting, the State has provided both incentives and disincentives for the volume of agricultural output.

Some common characteristics emerge from the agricultural policies implemented by successive governments. The State has constantly reorganized its services in line with its prevailing objectives; it has managed to control the pricing policy of agricultural products and has attempted to organize producers and provide them with infrastructure.

To facilitate the development of the agricultural sector, newly independent Burkina Faso created 11 Regional Development Offices (RDO). The core of these offices is the Extension Service, responsible for all rural development activities in a given area. These activities include distribution of fertilizer, insecticides, and chemicals for the protecting stored grains; collecting of credit; purchasing of farm products; promoting cooperatives; collecting of statistical data; promoting community activities; and giving advice on planting techniques.

The first Five-Year Plan, launched in 1967, gave priority to increasing the output of cereals through better cultivation methods and the use of fertilizers. As in previous years, the objectives were diversification of crops, soil conservation, and improvement of the irrigation system. These improvements were expected to double agricultural output in 15 years. The long-run decision was also taken to establish a complementary system of agriculture and livestock. The plan foresaw the development of irrigated areas of intensive farming and the enlargement of regions of traditional dry farming. The total investment on rural development during that time amounted to 29.8% of total public investment.

The main objective during the second Five-Year Plan (1972-1976) was primarily to overcome the consequences of the drought the country had just suffered, which hit crop farming severely. In 1973 the existing parastatal RDOs were given the monopoly of purchasing grains from farmers. The importance of the RDOs enforced the role of the state as a buyer and purchaser of agricultural products. After four years of implementation of government measures, the objectives of monopolizing the grain trade and achieving price stabilization have not been reached. OFNACER, the national cereal board, was entrusted with greater responsibility for marketing cereals and supporting official producer prices at the level of farmers.

Like the food grain supplies, the livestock sector was severely affected by the drought of the early '70s. The drought-induced losses were estimated at 12-15% for cattle and at 8-10% for sheep and goats. From 1975-1977 the government undertook several measures to reconstitute the cattle stock. To facilitate these efforts, a new agency, the ONERA (*Office National d'Exploitation des Recherches Animales*) was set up. Its task was to develop domestic and export marketing of livestock and livestock products.

Given the substantial disparities in agricultural potential between the Mossi plateau and the peripheral areas, government policies aimed to redistribute the population to

alleviate population pressure on the Mossi plateau. The attempt to out-migrate people from the Mossi plateau into areas of lands with richer soil, better rainfall, and low population density in the peripheral areas was a means of promoting the country's long-term agricultural development. A government agency, the *Autorité d'Aménagement des Vallées des Voltas* (AVV), was created in 1974 to organize settlement in the White, Red, and Black Volta. The government also launched projects to eradicate the simulum fly, tsetse fly, and malaria in infected areas and also planned to mechanize production and begin resettlement schemes.

Since the 1980s, the government and NGOs have taken measures to reduce the heavy dependency on rainfall. Despite the high number of factors militating against overly ambitious irrigation development, it was imperative that the government formulate at least a moderate irrigation development program. The major objective of the program was to complement rain-fed agriculture at the margin and to gradually overcome the constraints identified earlier. Later, government introduced local low-cost water retention schemes and started irrigation programs such as the Sourou Valley Rural Integrated Development Program.

The agricultural policy of the years of the revolution years was distinguished by a great plurality of objectives: production and productivity gains; income stabilization; food security; foreign-exchange earnings; and fiscal revenues. The implementation of policies to meet these objectives involved a series of "classic interventions," such as minimum producer prices, maximum consumer prices, administration of agro-industrial output and input prices, stabilization levies or compulsory paybacks, marketing and stabilization boards, and regulation of markets and foreign trade. The revolutionary government also gave priority to constructing earth dams and digging wells for irrigation. Its first economic objective was to achieve food self-sufficiency in the decade that followed revolution.

But even while trying to improve the agricultural sector and the lot of the peasants during the first years of his administration, Sankara adopted the 1985 Agrarian and Land Reform Law, which basically hurt the interests of the peasants. The law nationalized the land that had been private property or owned according to custom. The program was meant to promote the rational use of land to increase productivity and social justice but was viewed with great hostility by peasants.

One major positive change in the agricultural sector brought about by Sankara's policy was that of self-help rural development. The best known of such projects is the Sourou valley agricultural project. In the first phase a 711-m canal was constructed in the Sourou valley. The canal was constructed entirely by local labor to bring water from the Mouhoun to the Sourou dam. The second phase involved the establishment of 40,000 hectares of irrigated land and the construction of a barrage at Samandéni. Of this land, 15,700 hectares were divided into plots worked by small shareholders and land for agro-industries. The rest was run by the State, with about 3,000 hectares devoted to sugar production and 5,800 hectares to cotton and oilseed producing crops. The Sourou project is an example of what can be achieved in an arid area that was considered to have little potential. Self-help schemes for soil improvement, especially in the Mossi highlands, where soil erosion due to over-cultivation was prevalent, were also developed.

In May 1987, the Ministry of Agriculture established new, decentralized development agencies, the Regional Agricultural Production Centers (*Centres Regionaux de Production Agropastorale*). They replaced the Rural Development Agencies (*Organismes de Developpement Rural*), which were abolished in 1986. Their essential function is to provide extension schemes.

Besides that, numerous initiatives to promote particular crops or activities were launched between 1982 and 1987, but were subsequently abandoned for lack of resources or follow-up. Government has consistently considered agriculture as the engine of growth but changed the nature of its development from being extensive, which has harmful environment consequences, to being intensive. In 1988, 11 Agricultural Production Units were set up, their aim being the operation of irrigation infrastructures.

The most pertinent concluding description of the Burkinabé agricultural policy was made by a World Bank report by the end of the 1970s, which said: "*A complex system of institutions, regulations and mechanisms which give the government a dominant role*" involving "*four key ministries, six marketing and stabilization boards, more than 20 decentralized agencies, and numerous parastatal enterprises for production, extension and research. Fiscal, price and trade regulations are both numerous and cumbersome.*" That system obviously introduced substantial economic inefficiencies that call into question its justification. Institutional reforms aimed at rationalizing the role of the public sector were initiated in the mid-1980s. These reforms started to restructure the institutions involved in the agricultural sector and to identify the need for greater coherence and eventual liberalization of price and marketing policies.

III.3 Changes in population, social and cultural patterns of water management

Population Distribution in Ghana

Population density increased steadily from 36 per square km in 1970 to 52 per square km in 1984; in 1990, 63 persons per square km was the estimate for Ghana's overall population density. These averages, naturally, did not reflect variations in population distribution. For example, while the Northern Region, one of the ten administrative regions, showed a density of 17 persons per square km in 1984, in the same year Greater Accra Region recorded nine times the national average of 52 per square km. As was the case in the 1960 and 1970 figures, the greatest concentration of population in 1984 was to the south of the Kwahu Plateau. The highest concentration of habitation continued to be within the Accra-Kumasi-Takoradi triangle, largely because of the economic productivity of the region. In fact, all of the country's mining centers, timber-producing deciduous forests, and cocoa-growing lands lie to the south of the Kwahu Plateau. The Accra-Kumasi-Takoradi triangle also is conveniently linked to the coast by rail and road systems--making this area an important magnet for investment and labor

By contrast, a large part of the Volta Basin was sparsely populated. Tsetse flies, the relative infertility of the soil, and above all, the scarcity of water in the area during the harmattan season affected habitation. The far north, on the other hand, was heavily populated. The 87 persons to a square kilometre recorded in the 1984 census for the Upper East Region, for example, was well above the national average. This may be explained in part by the somewhat better soil found in some areas and the general absence of the tsetse fly; however, onchocerciasis, or river blindness, a fly-borne disease, is common in the north, causing abandonment of some pieces of land. With the improvement of the water supply through well-drilling and the introduction of intensive agricultural extension services as part of the Global 2000 program since the mid-1980s, demographic figures for the far north could be markedly different by the next census.

Another factor affecting Ghana's demography was refugees. At the end of 1994, approximately 110,000 refugees resided in Ghana. About 90,000 were Togolese who had fled political violence in their homeland which began in early 1993 (see Relations with Immediate African Neighbours, ch. 4). Most Togolese had settled in Volta Region among their ethnic kinsmen. About 20,000 Liberians were also found in Ghana, having fled the civil war in their country (see International Security Concerns, ch. 5). Many were long-term residents. As a result of ethnic fighting in northeastern Ghana in early 1994, at least 20,000 Ghanaians out of an original group of 150,000 were still internally displaced at the end of the year. About 5,000 had taken up residence in Togo because of the strife.

Urban/Rural Disparities

Localities of 5,000 people and above have been classified as urban since 1960. On this basis, the 1960 urban population totalled 1,551,174 people, or 23.1% of total population. By 1970, the percentage of the country's population residing in urban centres had increased to 28%. That percentage rose to 32 in 1984 and was estimated at 33% for 1992.

Like the population density figures, the rate of urbanization varied from one administrative region to another. While the Greater Accra Region showed an 83% urban residency, the Ashanti Region matched the national average of 32% in 1984. The Upper West Region of the country recorded only 10% of its population in urban centres that year, which reflected internal migration to the south and the pattern of development that favored the south, with its minerals and forest resources, over the north. Urban areas in Ghana have customarily been supplied with more amenities than rural locations. Consequently, Kumasi, Accra, and many towns within the southern economic belt attracted more people than the savannah regions of the north; only Tamale in the north has been an exception. The linkage of the national electricity grid to the northern areas of the country in the late 1980s may help to stabilize the north-to-south flow of internal migration.

The growth of urban population notwithstanding, Ghana continued to be a nation of rural communities. The 1984 enumeration showed that six of the country's ten regions had rural populations of 5% or more above the national average of 68%. Rural residency was estimated to be 67% of the population in 1992. These figures, though

reflecting a trend toward urban residency, were not too different from those of the 1970s when about 72% of the nation's population lived in rural areas.

In an attempt to perpetuate this pattern of rural/urban residency and thereby to lessen the consequent socio-economic impact on urban development, the "Rural Manifesto," which assessed the causes of rural underdevelopment, was introduced in April 1984. Development strategies were evaluated, and some were implemented to make rural residency more attractive. As a result, the Bank of Ghana established more than 120 rural banks to support rural entrepreneurs, and the rural electrification program was intensified in the late 1980s. Government, moreover, presented its plans for district assemblies as a component of its strategy for rural improvement through decentralized administration, a program designed to allow local people become more involved in planning development programs to meet local needs.

Annual population growth rate in the Volta Basin of Ghana

The third demographic variable that was analyzed regarding its role in affecting agricultural land use is the annual population growth rate. Results of the annual population growth rate in the various sub-basins of Ghana in three periods, namely 1960-1970, 1970-1984 and 1984-2000, are shown in Table 77 and Figure 29 depicts the trends in the annual population growth rate.

Table 77: Annual population growth rate by sub-basins, 1960-2000

Sub-basin	Annual population growth rate		
	1960-1970	1970-1984	1984-2000
White Volta	5.4	2.7	1.5
Black Volta	3.4	3.3	1.2
Main Volta	4.9	2.7	2.5
Oti basin	7.9	3.1	3.0
Daka basin	6.6	2.4	2.8

Source: 1984 population census of Ghana, special report on localities by local authority and 2000 population and Housing Census, Summary Report of Final Results (GSS 2002b)




Figure 29: Annual percent population growth rates by sub-basins, 1960-2000

The annual population growth rate decreased for all the three periods in all sub-basin, the only exception being the Daka sub-basin, where it increased from 2.4 % between 1970 and 1984 to 2.8 % in 1984-2000. Annual population growth rates were very much higher between 1960 and 1970 compared to the periods in all the sub-basins

Agricultural land use in the sub-basins

Agricultural land use for 1992 and 2000 in the Volta river sub-basins, as shown in Figure 30 was characterized by a mixture of increases and decreases. While there were increases in the White, Black and Main Volta sub-basins during the period, the Oti and Daka sub-basins experienced a decline.




Figure 30: Average cropped area by sub-basins, 1992 & 2000

Source: Ministry of Food and Agriculture, Ghana, 2001: in Codjoe (2004)

The highest annual increase (89.9 %) in agricultural land use in the White Volta sub-basin is recorded in the Lawra-Jirapa Local council, and the lowest a 0.3 % increase in the Walewale local council.

Migrations and mobility in the Volta basin of Ghana

The spreading Lake behind the Akosombo Dam forced 739 villages along the banks to be moved. However, this was not the first time that Ghana was resettling citizens. In 1956, some Frafra people in the Northern Region first had their homes relocated because of overpopulation. The second resettlement was the Tema Manhean Project (1959), which resulted from the construction of the Tema Harbor seaport.

The Volta Resettlement Scheme, Ghana's third of its kind, had a budget of \$9.8 million. Before the 80,000 people could be displaced, economic, social, physical, political, and psychological factors had to be addressed. Since resettlement would drastically affect the lifestyle of the people, it was essential to make the transition as smooth as possible while preserving sacred traditions and rituals of life. Furthermore, there was a prevailing sentiment among the people to be displaced that the government "owed them something." Many of the people needed to regain a sense of worth and reestablish their contribution to society.

Everyone was given the option of either monetary compensation or resettlement into one of the 52 specially constructed townships. Over 70,000 people chose resettlement instead of monetary compensation. Many of the resettled people moved on immediately when confronted with shortages and inadequacy of housing, of cleared land, of money and of food. A study by FAO, revealed that by 1970, only five years after the move, only 25,000 of the settlers remained in the villages, illustrating clearly the collapse of the old way of life of the various communities in the area. Unfortunately, there are no studies on those who chose to leave the villages compared to the study on those who remained. However, it is fair to say that by and large it was men who left to work elsewhere, leaving behind women and children together with government officials (Graham, 1972).

Many case studies have shown that the move from old to new villages was itself a traumatic experience for many of the resettled people. Shortage of time, lack of money and inadequate planning combined to ensure that few of the villages were completed by the time people were arriving. By 1966, the disruption of existing social, cultural and economic relations was almost complete.

Before Akosombo, many of the people along the Volta lived in tiny scattered villages. The average village house was constructed with swish (soil-based) walls and thatched roofs. Subsistence farming, animal grazing, and river fishing were the most common practices in the area and these traditions were passed down from generation to generation. According to the 1956 preparatory studies, only about 6% of the land covered by the lake was "used productively," while the rest was "unsuitable for agriculture or unoccupied" with low cash incomes and many medical problems (Hart, 1977).

The resettled were referred to as subsistence farmers, with low levels of productivity,

which contributed little to the national development effort. According to Hart (1977), the resettled farmers were in fact, agricultural laborers for the VRA and not individual farmers acting in mutual cooperation.

The resettlement scheme offered a unique opportunity to consolidate the scattered villages into more organized communities and provide them with schools, improved sanitary facilities, and increased revenue potential through mechanized farming techniques and organized livestock breeding. Further this consolidation would facilitate future electrification of the area. Indeed the resettlement was seen then by the Government of Ghana as an opportunity to implement the ideas of “modernization theory” through wide scale social engineering. Isolated communities, living in harmony with the Volta River and practicing the most appropriate form of shifting agriculture, were not only expected to acquiesce to the resettlement scheme for the greater good of the nation, but were also expected to change their whole way of life to conform to the conventional models of development.

A resettlement house was constructed with landcrete walls (concrete made with local soil) and an aluminium roof. In terms of the sturdiness and durability of the building materials used, a resettlement house was superior to the average village house. Although each family was given a "core" house, overcrowding was common because previously, every household built as many rooms as they needed and, many of the houses were never completed.

The resettlement project also aimed to replace the common practice of subsistence farming with "cash crop" farming. In doing so, each farmer would be taught new farming techniques to produce enough for his family and some extra crops to sell for income. In order to support crop rotation, virgin forests were cleared for farmlands; but this was not always done without resistance (the government often clashed with traditional chiefs over who owned the land). As a subsidy, the government also provided chicks, piglets, and other young livestock for the people to rear on the new livestock farms and later, to sell for profit. Later, many of the animals began to die prematurely from disease and malnutrition (usually from improper care), thus, the government stopped giving out the young livestock at no cost to the farmers at no cost.

Many of the most crucial problems in the new villages stemmed from the land issues It has been planned that sufficient land be cleared to provide every farmer with 12 acres (4.86 ha). However, by 1966, when the deadline for the dam construction had been reached only 15, 000 acres (6,075 ha) had been cleared out of a target of 54, 000 acres (21,678 ha). By way of comparison, the flooded area had 128, 000 acres (51,830 ha) in productive use.

With only a quarter of the required land cleared it was clear that the resettled could not practice their traditional farming skills, or even continue growing their subsistence crops. Confronted by this land shortage, many of the resettled could not farm at all. The land clearing programme was initially based on the use of heavy machinery but by 1967, the scheme was abandoned due to lack of spare parts and lack of heavy earth moving. The

VRA had also hoped to introduce new farming methods to the people and also controlled the allocation of seeds and fertilizer and the bulk of buying the production. Farmers were assigned to various types of agriculture. Some were to become crop growers, others tree-crop farmers, and the rest livestock farmers. More specialized activities, like tobacco and cotton cultivation, and pig and poultry were later introduced. By 1968, it was clear that the new system of intensive, mechanized farming had failed.

The creation of the lake has also led to a massive escalation of disease among the lacustrine population. Among the feared consequences of the lake was an increase in three notorious afflictions: schistosomiasis (*bilharzias*) onchocerciasis (*river blindness*) and malaria. Although various efforts have been made to improve the health of the resettled, it is clear, despite the absence of statistics, that the health standards of the lacustrine population have been considerably lowered by the impounding of the river.

Many agree that the Volta Resettlement project improved the physical environment for the average rural Volta citizen however, the debate continues as to whether or not there were social and psychological improvements. VRA tried as much as possible to settle people of the same ethnic group in a village. But sometimes, there were cultural conflicts because a Fanti would not want to be governed by an Ewe chief, for example. Other problems arose from lifestyle issues.

Some who were seasoned fishermen did not want to become cash crop farmers. Many left their resettlement homes and constructed wooden shacks along the lakeside so that they could be closer to their best-known source of income. As more people encroached on the lakeside and the communities diversified their activities, illegal clearing and farming along the banks led to increased sediment deposit in the lake. With the receding perimeter of the lake due to the drought in 1998, the government made efforts to replant trees along the lakeside to control erosion.

IV.3.4 Legal Framework: Presentation of the Water Laws

A. Statutory Legal Framework for Water

In this section, we shall discuss the emerging consensus on water policy and the major institutions created by statute to deliver water in Ghana. We shall also discuss the institutions created to regulate the water sector.

A.1 National Water Policy

It is difficult to state that Ghana has a Water Policy because there is no national water policy document that states the intentions of government on the management of Ghana's water resources. What exists is a document by a number of consultants entitled '*Ghana's Water Resources, Management Challenges and Opportunities*', (to be referred to as *Ghana's Water Resources*) published by the Ministry of Works and Housing, which makes recommendations about how Ghana's water resources should be managed. It

seems however that government is following these recommendations. We shall discuss the salient recommendations in the document in the light of the difficulties that have faced the country in managing its water resources.

Until recently there was no institution in Ghana that had overall responsibility to manage water resources. Several single purpose state agencies like the Ghana Water and Sewerage Corporation (GWSC), the Volta River Authority (VRA), and the Irrigation Development Agency (IDA), pursued their individual mandate with little co-ordination and without any thought about what other water users required. Little emphasis was placed on the management and conservation of water. Second, water was provided in a way that did not capture the economic value of producing and conserving it. It was either provided free of charge or at a very subsidized rate. Third, there was no effective regulation of the water sector. The likely result was that the management of water could not be sustained.

The recommendations made in *Ghana's Water Resources* are supposed to correct these mistakes. It was recommended that Ghana should move away from a state dominated water sector to a private sector led industry where the private sector plays the central role in investing and delivering water. The recommendations argue that there are costs involved in managing water in a sustainable manner (*Ghana's Water Resources*, pp 55-57). A price must therefore be placed on the use of water resources to capture its value as a scarce resource. Raw water abstraction should attract a fee. The 'polluter pays' principle should be used to control discharge of effluents into water. People who use water for non-extractive purposes like fishing, transportation and recreation should be made to pay a fee for its use. The introduction of water charges should facilitate the promotion of water conservation, the protection of the aquatic environment and the maintenance of a database on water resources (*Ghana's Water Resources*, pp 46-51).

The objectives of regulation should be to ensure an efficient co-ordination and planning of water resources in Ghana. It should protect the economic life of existing investments in infrastructure and create the appropriate framework for people to invest in water and its resources. It is hoped that such an environment will safeguard the public interest against failure of hydraulic structures, flooding and river bed and reservoir sedimentation and facilitate the participation of all users in the management of water resources.

It also suggested changes in the institutional setting for managing water resources and controlling pollution. It argued for a multi-sectoral approach to water resources management. It recommended that a single agency should be established to co-ordinate the sector's activities to ensure that there is minimum conflict between different water users.

A.2 National Water Legislation

There are many agencies in Ghana that provide water under the statute law. However, the most important ones that provide water for domestic consumption are the Ghana Water Company Limited (GWCL) (formally Ghana Water and Sewerage Corporation) and the Community Water and Sanitation Agency (CWSA). By practice and

orientation, GWCL has concentrated on providing of water and sewerage services in urban areas, while the CWSA facilitates the provision of water in rural areas. The CWSA works in close collaboration with District Assemblies. Recently, a number of regulatory institutions have been established to regulate the delivery of water to consumers and also to protect the environment and conserve water resources.

A.3 The Ghana Water Company Limited

The GWCL was set up in 1965 as a public corporation and later became GWSC. A managing director who is accountable to a Board of Directors heads the Corporation. GWSC has 10 regional offices headed by regional directors.

Under the **GWSC Act 1965**, GWSC has authority to provide, distribute and conserve water for domestic, public and industrial purposes (section 4 of GWSC Act 1965, Act 310). It is also supposed to establish, operate and control sewerage systems in Ghana. The Corporation has power to make long-term plans for the provision of water and the operation of sewerage systems in Ghana. It has authority to conduct research into water and sewerage issues. It can also make engineering and survey plans and it can construct and operate water and sewerage works in the country. It sets the standards for water supply and the operation of sewerage systems.

Under LI 1233, the Corporation has enacted regulations through which water and sanitation facilities can be made available to Ghanaians. These regulations are meant to protect the environment, protect natural watercourses and provide for proper sewerage systems. Among its important powers, GWSC can by a notice published in the Local Government Bulletin, declare an area to be a 'connection area'. It can also in consultation with the Town and Country Planning Authority, establish planned connection areas. When an area is declared a connection area, GWSC can take over any private sewerage system there. Anybody who wishes to build a private sewerage system in a connection area must submit his plans and drawings to the GWSC for approval. The GWSC can also enter any land in a connection area and install or inspect a sewerage system (Sections 1-20 of GWSC Regulations 1979, LI 1233).

The GWSC has power to serve a notice on the owner or occupier of land in a connection area to apply for the installation of a water or sewerage system. A person can also on his own apply for the grant of a permit to construct a sewerage system. No private sewerage system can be connected to a public sewerage system without the approval of GWSC. The Corporation has authority to enter a person's premises or land and install or complete the installation of a sewerage system without the person's approval and at the person's expense. No one can receive a building permit in a connection area if he does not have a permit to install pipes and sanitary appliances in the building to be constructed.

The Corporation has regulatory powers to protect natural waterways. It is therefore illegal to pollute a watercourse or to cause damage to a sewerage system. Nobody has authority

to dig, excavate or remove earth around a public sewerage system without the prior approval of the Corporation. No one can build a structure over or near a public water way or sewerage system without the prior approval of the Corporation.

Customers of the Corporation are under obligation to pay for services provided by the Corporation. This has to be done within 14 days of the receipt of a bill. The Corporation has authority to disconnect a person who fails to pay for services it has provided. It can also institute action in court to recover the money owed to it.

A.4 The Community Water and Sanitation Agency

As we have already stated, GWSC has concentrated on the supply of water to urban areas. It failed to provide water for the rural areas of Ghana. In 1994 an autonomous division of the GWSC, the Community Water and Sanitation Division was created to facilitate the provision of water for rural areas in Ghana. By Act 564, the Community Water and Sanitation Agency became an institution in its own right and it is no more a division of GWCL. The CWSA is in the process of separating itself from GWSC.

Among the objectives of the CWSA is to facilitate the provision of safe water and sanitation services for rural communities and small towns (Section 2 of Community Water and Sanitation Agency Act, 1998, Act 564). It must support district assemblies in promoting sustainable safe water and sanitation services in rural areas. It must support district assemblies to encourage the participation of communities, especially women, in the management and construction of water and sanitation facilities. Among its responsibilities, it has to design strategies for mobilizing resources for the execution of water and sanitation projects. To achieve this objective, it must encourage the private sector to participate in the provision of water and sanitation facilities. It must also provide district assemblies with the technical assistance required for executing water and sanitation projects. It also co-ordinates the activities of NGOs involved in the provision of rural water, sanitation and hygiene education. It must collaborate with the Ministries of Local Government, Environment, Health and Education in increasing consciousness about water related health hazards. It must set standards for the provision of water and sanitation services. It has to charge fees for the services it provides and it must collaborate with International Agencies it considers necessary for implementing its programmes. It must collaborate with the Water Resources Commission, the Environmental Protection Agency, the GWSC and other public and private bodies that are involved in the providing of water and sanitation services for rural communities in Ghana (Mensah, 1999).

A.5 District Assemblies

Ghana has embarked on a decentralization programme that is aimed at allowing decisions affecting communities to be taken at the lowest appropriate level. Under the Local Government Act 1993, Act 462, district assemblies are the highest political and

administrative authorities in the district (Sections 10(1), 10(3)(c)(d)(e) of the Local Government Act, 1993, Act 462). District Assemblies must promote productive activity and social development in their districts. They must also remove all impediments to development in their districts. It is their responsibility to develop the basic infrastructure and provide the municipal works and services required in the district. They are also responsible for the development, improvement and the management of human settlements and the environment in their districts.

Under the second schedule 2 of the Local Government (Assin District Assembly)(Establishment) Instrument, 1988, LI 1380 some of the Assembly's duties include:

- To ensure the provision of adequate and wholesome supply of water throughout the entire District in consultation with the Ghana Water and Sewerage Corporation.
- To establish, install, build, maintain and control public latrines, lavatories, urinals and wash places.
- To establish, maintain and carry out services for the removal of night soil from any building and for the destruction and treatment of such night soil.

The lowest political authority in a district is the unit committee. Under schedule 5 of the Local Government (Urban, Zonal and Unit Committee) Establishment Instrument, 1994, LI 1589, unit committees are supposed to perform the following functions:

- Mobilize members of the Unit for the implementation of self-help and development projects.
- Monitor the implementation of self-help and development projects.

There is thus a close connection between the Community Water and Sanitation Agency, District Assemblies and the Unit Committees. The Community Water and Sanitation Agency is responsible for facilitating the supply of water and sanitation to rural communities. A District Assembly is responsible for ensuring that adequate and wholesome water is provided in the district. They are also responsible for developing infrastructure, the municipal works and human settlements. They are also responsible for managing sanitation. Unit committees are responsible for initiating and monitoring self-help projects. The provision of water and sanitation under the Community Water and Sanitation Programme (CWSP) for communities is supposed to be self-help (Mensah, 1999).

A.6 Regulatory Institutions

The effectiveness of the laws governing resources poses another problem as the laws and regulations established for the management of water and soil resources appear to be weak and ineffective. In some instances, the laws are adequate but they are not adhered to or enforced either due to lack of institutional capacity or political commitment. The knowledge base of the state of natural resources, rate of depletion, and consequent future impact is poor, and probably contributes to the weak political commitment on the part of the various governments and general apathy on the part of the populace. The legal framework

governing the management of land and water resources is the Code for the Environment, the decree of 5 February 1933, and Code for Water, which is to be finalized under the management of water resources, and the Mining Code.

Customary laws and practices in Ghana have existed over the years and cover water conservation, pollution control, protection of catchment areas and protection of fisheries. These laws are enforced through various sanctions, usually determined by traditional authorities. However, apart from issues on the protection of water for domestic use that is already present in existing statutes, many traditional regulations are rather localized and do not have any common features that can be readily incorporated into the common law. The Rivers Ordinance (CAP 226, of 1903) was the first attempt to comprehensively control the use of water other than for domestic uses. There was no follow-up to this ordinance. No regulations were made, and time and other enactments have since overtaken the ordinance. Parts 1 and II of the Rivers Ordinance have since been repealed by the Water Resources Commission (WRC) Act 522 (1996). Part III of the Ordinance is, however, still operative. A list of all statute laws that are related to water resources in the country is presented in Appendix 1.

Until the Water Resources Commission Act was passed in 1996, agencies in the water sector were set up by legal enactments that provided the legal framework for the management of the resource. Each of the legal enactments contain specific provisions which grant the agencies the powers, with the approval of government, to make general and specific regulations by legislative instruments to enable the objectives and functions of the agencies to be better discharged. This arrangement gave room to a fragmented approach of managing the resource.

Water, as an essential natural resource falls within the provisions of Article 269 of Ghana's Constitution, which seeks to protect water resources by setting up a Commission to regulate, manage and co-ordinate government policies in relation to it. Section 12 of the Water Resources Commission Act (1996) stipulates that *'the property in and control of all water resources is vested in the President on behalf of, and in trust for the people of Ghana'*. The vesting of the water resources in the President is to make water resources management consistent with general natural resources management in Ghana and the 1992 Constitution. The principle implies that there is no private ownership of water in Ghana, but that the President, or anyone so authorized by him, may grant rights for water use. It also implies that with good governance and practice, the principle is expected to ensure that water allocation for various uses will benefit to the public interest and be for the greatest good of society.

Though the WRC Act vests the ownership of water resources in the State, it recognises all existing uses of water prior to the enactment of the WRC Act. However, all existing claims to water uses are to be submitted to the Commission within twelve months after the coming into force of the Act. The WRC Act also provides for certain categories of water uses which are exempted from the requirement of prior permit. Section 13 (2) of the WRC Act provides for the non-preventive use of water resources for the purpose of fighting fire, while Section 14 (1) states that *'a person who has lawful access to water resources may abstract and use water for domestic purposes'*.

Many institutions have been established to regulate the delivery of water in Ghana.

A.6.1 The Water Resources Commission

The Water Resources Commission was set up under article 269 of the 1992 Constitution to be the single agency responsible for co-ordinating water policy in Ghana. Act 522 established the Commission.

The Commission is responsible for the regulation, management and co-ordination of policy in connection with water (Section 2 of Act 522). It is supposed to propose comprehensive plans for the use, conservation, development and improvement of water resources in Ghana. It has the mandate to initiate, control and co-ordinate activities connected with the development of water resources. It grants water rights. It can also require water user agencies to conduct research into water resources. It has also authority to monitor and evaluate programmes for the operation and maintenance of water resources in Ghana. It advises the government on issues that are likely to adversely affect water resources.

Under the Water Resources Commission Act, the property in and the control of water resources is vested in the President on behalf of and kept in trust for the people of Ghana (section 12). No person has authority to divert, dam, store, abstract or use water resources, construct or maintain works for the use of water resources without the authority of the Commission. A person who has been lawfully granted access to water resources can abstract it and use it for domestic purposes. However, no one has authority to abstract water for domestic purposes without a permit (section 13-24).

Where the Commission thinks that the use of a water resource poses a threat to the environment or public health, it may by notice order the person polluting the water resource to stop the activity. When an application for a water right is made, the Commission is obliged to consult the inhabitants of the area where the right is to be granted, to ascertain their views on the grant of the water right. It also has to publish the application in the *Gazette* to allow anyone who has an interest in the water resource to indicate the nature of his interest. The grant of a water right may be subject to conditions. No one has authority to transfer a water right without a written approval from the Commission.

The Commission has power to suspend or vary a water right if it thinks that the water resource in an area is insufficient for public purposes. Where it thinks that the water resource is needed for a public purpose it can also terminate or limit a right already granted. Compensation can be given to the right holder if this occurs. A water right can be terminated for breach of a condition attached to the right or for non-use. It is an offence to interfere or alter the flow of water, or pollute or foul water beyond the limits prescribed by the Environmental Protection Agency.

The WRC has just started its work. It is in the process of formulating its policies on regard to the granting of water rights.

A.6.2 The Environmental Protection Agency

The Environmental Protection Agency was established in 1994. Its functions include advising the Minister for the Environment on policies on all aspects of the environment (Section 2 of the EPA Act 1994, Act 490). The Agency is also supposed to co-ordinate activities of all bodies concerned with the environment and to be the link between such bodies and the Ministry of the Environment. It has responsibility to co-ordinate the activities of bodies that generate waste with the aim of controlling the generation, treatment, storage and transportation of industrial waste. It must protect and improve the quality of the environment. It has power to issue environmental permits and pollution abatement notices to control waste discharges and emissions and to prevent or reduce noise pollution. It is supposed to provide standards and guidelines in relation to air, water and other forms of environmental pollution. It also has authority to ensure that developers comply with environmental impact assessments of their development plans before they begin development. The EPA is supposed to implement the National Environmental Action Plan.

A.6.3 The National Environmental Action Plan

The objectives of the National Environmental Action Plan are to improve the surroundings, living conditions and the quality of life for all generations of Ghanaians. It aims to ensure that there is reconciliation between economic development and natural resource conservation. It is hoped that a high quality environment would become an essential element in Ghana's economic and social development plans (Ghana Environmental Action Plan, vol. 1, pp 1-18).

The principles on which the National Environmental Action Plan is based are that there must be optimum sustainable yield from the use of natural resources. It requires that the most cost-effective methods should be applied to achieve environmental objectives. This means that incentives and regulatory methods should be used to achieve environmental objectives. Consequently, the polluter should be made to pay for the cost of preventing or eliminating the pollution or nuisance he has caused. Decisions on the environment should be taken at the lowest appropriate level. The public must also be encouraged to participate in decisions that affect the environment. Ghanaians must co-operate internationally to achieve environmental objectives (Mensah, 1999).

A.6.4 The Public Utilities Regulatory Commission

We shall discuss the nature of water supply to consumers in Ghana in later sections of this paper. Thus we do that, we shall see that among the major complaints of consumers, are the high tariffs they have to pay for water and the deteriorating quality of service they receive from GWSC. The PURC was set up in 1997 to regulate the tariffs charged by utilities and to protect both consumers and utility companies. It is supposed to provide

guidelines on rates that can be charged by utility companies and to approve rates chargeable by the utility companies. It is supposed to monitor the standards and performance of utility companies (PURC Act 1997).

According to Part III of the Act, the PURC is supposed to set up a complaint procedure for consumers dissatisfied with the services or actions of a utility provider. The PURC has also just been set up and is now developing policies to fulfill its mandate.

B. Nature and Status of Customary Water Rights

Before the creation of GWCL and CWSA, customary law was the regime through which water was supplied to the people living in the rural areas of Ghana. Due to the fact that water has not been provided for all rural communities under the CWSP, customary water law continues to be important in enabling people to get water.

B.1 Description of Customary Water Laws

Traditionally, water is a treasured natural resource (see Ofori Boateng J. 1977). The major sources of water in the customary regime are wells, streams, rivulets, and rivers in that order of importance.

Most rural communities in the Assin Foso District have access to a surface water source. The desire to situate communities near water sources is commonsensical since water is essential for life. Human settlements are however situated quite a distance from water sources. This prevents human settlements from suffering from floods during rainy seasons. Situating settlements away from water sources has its drawbacks. It can take some time to get water. However, this compels people to conserve water and it prevents the use of rivers as sewers.

In most communities, no institution or office has responsibility to *provide* water. This maybe because water is acquired solely from natural sources and available technology is not sophisticated enough to get deep ground water. During dry seasons, when very little water is available, it is the responsibility of individual households to look for water. Communities however have institutions to ensure that water is conserved and its quality is maintained.

Water in its visible form - as sea, rivers or lakes - cannot be privately owned. It is unclear whether it is public property. In some communities, surface water is public property. In others, it belongs to the king. In reality the king holds the water in trust for his people. In other communities, water is said to be 'ownerless'. Generally, a private person will never be allowed to purchase or own a surface source of water.

Within the customary regime, it is not easy to get access to ground water. Ground water is acquired through digging wells. It is therefore unclear whether underground water is

considered public or private property. There are clearly public wells, when the community digs the well and uses it as its source of water. When a private person digs a well on his land, the water is likely to be considered as private property. In fact in most communities, people are encouraged to dig private wells to reduce the pressure on the community well during the dry season.

In rural Ghana digging a private well involves considerable expense. This can range between ₵250,000 and ₵400,000. This is far beyond the incomes of the average rural person. A person is allowed to charge members of the public who want to take water from his well. This can be at a commercial rate. The price of a bucket of water from such wells can range from between ₵10 and ₵40.

Generally no permit is required to dig a private or a public well. In small towns however, where sanitation has become a serious problem, one needs to acquire a permit from a health inspector to dig a well. This is to prevent the citing of wells near refuse dumps to prevent water contamination.

The chief, elders and priests control, manage and regulate the use of water sources. In many communities, the rules for collecting water are commonsensical rules. For example, one cannot use a dirty bucket to collect water. In others, especially where the water source is a river, the rules take on a religious character. For example, women are not allowed to collect water when they menstruate. Generally, there are particular days and months when no one can go into the river or the sea. Some of these rules are meant to protect water quality and prevent the catching of immature fish. Sanctions for breach of rules include slaughtering of sheep, the provision of schnapps or the payment of fines to the elders. Traditionally, water is used mainly for domestic purposes. Very little navigation is done on rivers and therefore the customary law of navigation is not well developed. Irrigation is also not a common form of farming technique (Mensah, 1999).

Water from traditional sources is usually contaminated and generally causes diseases like guinea worm infestation, bilharzia, river blindness and cholera etc. Enforcement of customary law depends on the cultural habits of the locality. Generally enforcement is good if the community is homogeneous.

IV. 7 Urban Water Management

A. Access to Clean Water in Urban Ghana

Most people in urban areas rely on pipe-borne water supplied by GWCL. According to the Ghana Living Standards Survey Round 3 (GLSS 3) 1991/2, 100% of households in Accra and 99.3% in Kumasi have access to pipe borne water (Rakodi C., 1996). Access to individual piped water depends on the degree of area planning and on the extent of one's wealth (Rakodi C, 1996, p 17). In 1991, in the Accra Metropolitan Area 59% of households had an indoor piped supply connection. All wealthy households had an

individual supply and less than a quarter of such people shared their pipe connections with other households. Some 85% of middle-income people and only 53% of low-income people had a private supply. Only 40% of middle-income people and 9% of low-income people had exclusive use of their taps. Some 12% and 23% of middle and low-income people respectively, shared their supplies with more than 10 other households. Public standpipes are generally uncommon. Only 8% of households had access to them. Some 28% of people had no direct access to a tap. Some 5% of middle-income people and 34% of low-income households purchase water regularly.

Water is purchased from neighbors who have an in house connection. Although the time spent fetching water is half of that in rural areas, households of Accra households which do not have an on-site connection spend about 8.4 hours per week on this task. As stated earlier on, under LI 1233, people have to apply to GWSC for an on site connection. The main constraint to getting a connection is the cost. In 1996, it cost about ₵ 200,000 to get an on site connection. Consequently, there are several illegal connections.

Irregular supply and frequent shortages are well known in Ghana. Water does not usually flow during the day. It flows in the night. Residents have to leave their taps on in spite of the fact that when air passes through the taps, it causes the meter to read. People are then forced to pay huge bills even though the water has not been flowing. Sleep is usually disrupted for people who do not have an on site connection or who have to share water with other residents. Typically, girls over 7 years and women are those who queue for water. Women spend about 18 minutes a day getting water, while it is 12 minutes for men. Boys between 7 and 19 spend about the same time as girls to fetch water. Older men rarely fetch water. In addition to the intermittent daily flows, water often flows only a few days week. In some areas, interruptions can last for months. Interruptions affect all areas, irrespective of socio-economic status.

Households have developed various strategies to deal with water supply interruptions. Some 96% of households store water. They use various types of containers including Jerry cans, pig feet containers, overhead tanks and buckets etc. Some 50% of wealthy people, 16% of middle income people and 3% of low income people use overhead tanks. It is possible to buy water from GWSC tankers or from private suppliers. This solution is available to the few people with overhead tanks and it is very expensive.

Another solution to frequent water shortages is purchasing water from areas where water flows constantly. This results in women walking long distances for water. The potential hazard to girls who enter the homes of strangers is a source of worry to many parents. Others use water from wells or collect stream water during rainy seasons. Some collect rainwater from roofs when it rains. Others recycle water within their homes.

Storing water has important health implications. There is evidence of faecal contamination in some samples of tap water consumed. This is because connections are illegal or poorly fitted. Contamination of water as a result of in-house storage is also

widespread. This is due to the fact that buckets used to store water are also used for other purposes. There is also irregular cleaning of storage containers. Failure to boil water and the frequent dipping of cups into water are other sources of water contamination.

A.1 Problems of GWCL

The major problem facing the Corporation is that it is a public corporation that has been subject to the dictates of politicians. Consequently water tariffs have for long been kept low with the aim of protecting consumers (*Ghana's Water Resources p45*). Politicians have also used it as a source of patronage and public policy towards the Corporation has been geared towards satisfying political objectives instead of assisting the Corporation to be managed efficiently. It has therefore been extremely difficult for the Corporation to meet its operational and capital plans. The result has been low investment and a general breakdown of GWSC 's water delivery system.

There has been very little investment in equipment. In most regions billing is done manually. This creates opportunities for fraud and illegal use of Corporation's funds. It is estimated that about 29% of the Corporation's customers are not billed. About 50% of meters do not work and the rate of revenue collection is low. About 30% of the Corporation's 207 systems are not operational and most of them have not been operational for more than a year. About 30% of the production systems are operating below capacity. By year 2000, it is estimated that daily demand for water will be about 1,200 MI/day. The Corporation's production is about 500 MI/day. The Corporation however has an estimated capacity of only 600 MI/day. There is thus significant under capacity to meet demand. Since about 50% of the Corporation's production meters are not working, which also makes revenue collection low, the Corporation's estimate of 43% of water unaccounted for is crude and is likely to be higher than is stated (*Halcrow Consultancy Report, 1995, p 11-13*).

A.2 Reform of Urban Water Sector

Government has been under pressure from donor agencies to reform water delivery in urban areas. This has resulted in the *Halcrow Consultancy Report* and the *Berger Report*. Until recently, the government was reluctant to privatize the whole urban water sector but was willing to commercialize certain operations including billing. It has also agreed to lease self-contained water systems from source to end users to private operators. Recently, however, there is evidence that under intense pressure from donors it has decided to privatize all utility companies including that of water.

The biggest problem facing these reforms is that there is very little debate about the reforms. Every step is shrouded in secrecy. It seems Government is being forced to take a direction that it does not approve of. The general public has been kept in the dark about the directions of the reforms and consequently, there is very little public support for these measures. Institutions that deal with the water sector are unsure of what to do. This is creating problems of institutional planning and co-ordination.

B. Access to Clean Water in Rural Ghana

B.1 Community Water & Sanitation Programme (CWSP)-objectives

The CWSP assists communities to improve their capabilities to meet their demand for clean water. It ensures that there is a minimum basic service of water that is protected all year-round, of 20 litres per capita per day, within 500 meters for consumers living close to a water point. A water point must not serve more than 300 persons (CWSP Implementation Manual, 1996 p1).

The CWSP is a demand driven programme. Communities are expected to participate fully in the programme in order to sustain it. They are supposed to organize themselves by contributing financial and human resources to ensure the success of the programme. Consequently, only communities that are willing to contribute to part of the cost of providing the service (generally 5%) and who are ready to pay for the maintenance of the facilities provided, become part of the programme. Women, as the primary collectors and users of water are given a central role in the design and management of water facilities. The private sector has the responsibility to design, construct, and maintain the facilities. NGOs play an important role by providing training in the management and repair of water facilities. The programme integrates health, sanitation, and hygiene education in order to maximize the benefits of the programme. As already indicated, local government authorities are closely involved in the programme. They provide infrastructure and organize the institutions necessary for effective and efficient use of water resources.

B.2 Access to Water under the CWSP

In most communities under the CWSP about 95% of people use the boreholes or hand dug wells provided under the programme. The other 5% use the traditional wells and other traditional sources of water. The water provided under the CWSP has to be paid for and the amount paid is determined by the WATSAN in consultation with the community. It ranges from ¢10 to ¢30 per bucket. The sale of water has been so successful that some communities have used the money from their water funds to embark on other development projects. No community has complained about a long interruption in their supply of water under the programme. When interruptions occur, it is due to the breakdown of a part. This is however quickly repaired because of the training given to women in repairing facilities. Problems may occur in the future when spare parts are not distributed as part of the programme. Generally, women spend less than 15 minutes to fetch water.

Storage of water continues to be a big problem. Water is stored in barrels, buckets and open tanks and this can lead to water contamination. While there has been a serious reduction in guinea worm infestation, there is still a problem with cholera and typhoid.

Another weakness of the CWSP is the weak institutional capacity of local governments. Local government has not shown the level of interest that was expected from it under the programme. The creation of DWSTs has been extremely slow. This has affected the progress of the programme.

B.3 Assessment of the Various Regimes

We now have to assess the various regimes for delivering water to people in Ghana. Do they take an entitlement view of water? Do they have mechanisms for alleviating poverty? How do they ensure that the environment is protected and water quality is maintained? Do they have a workable system of water conservation and what role is given to women in the managing water resources?

B.3.1 Assessing GWSC'S Performance

Before the 1990s, GWSC was basically a public corporation that was part of the public sector. Due to this role, it took an entitlement view of water. Service to most poor urban areas was free, while service for homes with on site connections was subsidized. This led to misuse of water and low cost recovery. With the advent of structural adjustment and the commercialization of its services, GWSC is now taking a very commercial approach to the delivery of water. Tariffs have increased substantially over the last few years and most standpipes in poor urban areas are being discontinued. GWSC has employed vendors who control the few remaining standpipes. These vendors sell water to customers.

Presently, if one does not have the means to purchase water, one simply cannot get it legally. For those who have no access to GWSC vendors, they have to purchase water from private vendors who make huge profit from selling water. Since there are no mechanisms for providing water at reduced tariffs to the poor, there is clearly an anti-poverty alleviation dimension to present urban water policy. Access to water for the poor has been greatly reduced through the combination of the phasing out of public standpipes and increased water tariffs. GWSC argues that this is necessary in the short term to save the Corporation from collapse. After it has corrected its balance sheet and it has been able to attract investors, it would be able to provide a more reliable service and also provide a subsidy for the poor in urban areas. There may be a basis for these arguments. In the short term however, it is fair to say that urban water policy has not been favored of the poor.

GWSC has also been unable to ensure water quality. This has been due in part to its lack of funds to meet its basic mandate. The second reason is even more fundamental. Ghana's population has grown rapidly since independence. In the last 15 years, there has also been intensified economic activity nation wide. This has led to severe pressure on land. Haphazard urban development has contributed to poor land use. These issues affect the environment and water quality. Unfortunately, GWSC has no control over such events. These problems have made its ability to control water quality an impossible one.

For the reasons stated above, GWSC has also been unable to play its role as a conservator of water. As indicated earlier on, the Corporation claims that 43 % of its water is unaccounted for. However, conservation has not been a major concern in Ghana because of the myth of an abundance of water. As the population increases and demand for water grows, there will be the need to provide strong conservation measures to protect the country's water resources. It is hoped that the Water Resources Commission and the Environmental Protection Agency that have just been set up, will combine effectively to deal with the environmental problems of water resource management.

In urban areas, the gender problem is not as acute as in rural areas. Women without on site connections still spend considerable time to get water. With increasing tariffs and with the closure of many standpipes, there is bound to be increased pressure on women. They may have to walk longer distances to get water. They will also have to invest time and money to purchase water. Another important issue is access to water and sanitation for the urban poor. The service they get from GWSC is also extremely poor. No arrangements have been made to capture the particular concerns of this socio-economic group.

B.3.2 Assessing the CWSP

How does the CWSP fare according to the issues raised earlier on? It seems that the CWSP is moving away from an entitlement view of water. Unless a community is ready to pay its contribution for water facility and is prepared to maintain it, it does not become part of the programme. Water from the facility is also sold to community members. Presently, there is no mechanism for allowing people who cannot afford to pay for water to get it for free.

The CWSA has an effective answer to this criticism. It claims that lessons from the past indicate that when facilities are provided for free, water and water facilities are misused. It is difficult to have an effective answer to CWSA's position on the entitlement problem. However, there is no doubt that there are people in rural communities who cannot afford to pay for water. If mechanisms are not provided to get water to them at subsidized rates, they will continue to use the traditional sources of water and may acquire infectious diseases. When this occurs, it would undermine the whole programme. NGOs have also complained about the rigidity with which the CWSA insists on the satisfaction of the finance side of the programme. They argue that a community's commitment to water and sanitation cannot be measured by the amount of money it can raise alone. Willingness to contribute in other ways like providing labor, etc ought to be used in assessing whether a community is committed to the programme or not. In fact, some NGOs claim that rigidity of the CWSA on the finance side, prevents communities that are ready to 'own' their water facilities from getting access to water (Interview with Country Director of WaterAid).

Has the CWSP assisted in poverty alleviation? Formally, we shall state that the CWSA is a realistic programme. It allows communities to prove their commitment to the programme. In this sense it does not provide free access to water. However, the conditions that have to be fulfilled are not too stringent. Only extremely poor communities will find it difficult to meet these conditions. Substantively, there is evidence that the CWSP has greatly improved the condition of life in rural communities. Communities under the CWSP have drastically reduced diseases like guinea worm infestation. The provision of water has also improved personal cleanliness and socio-economic life in these communities. Trained personnel like teachers are now willing to accept postings into these areas. There has been a marked improvement in school attendance by children (Ababio R. A., 'Rural Water and Sanitation: A Case Study of Eyisam, Abontsen and Ekumfi Dunkwa in the Mfantseman District of the Central Region of Ghana', 1996).

Environmental protection and water conservation are however still major problems in communities. Communities make rules to protect the quality of water that comes from the water facility. For example, one cannot wash near the water point, or use a dirty receptacle to collect water. However, communities cannot handle the complex problems of water conservation and environmental protection. There is also a serious problem of low capacity of District Assemblies to protect the environment and safeguard water quality at the local government level. They have neither the personnel, nor the funds, nor the equipment to handle these complex tasks. On the national scale water conservation and water quality protection have to be handled by institutions such as the Water Resources Commission and the Environmental Protection Agency.

The CWSP has major achievements in solving the gender problem. Women are given prominent roles in the design and management of water facilities. They are generally responsible for repairing water facilities. Since a water facility is not supposed to be more than 500 metres from the community, it has drastically reduced the distance and the time needed to get water.

B.3.3 Assessing the Customary Regime

As already indicated, there is no charge for collecting water from a customary water source. A person can take as much water as he/she can carry as long as he/she leaves enough for others. This is an entitlement view of water. This entitlement view is due partly to that very little investment goes into collecting water from customary sources. This is not the only reason. It is also due to the recognition that water is essential for human survival. And this can be seen in the different attitudes towards water during the dry and wet seasons.

In the wet season when water is plentiful, one can take as much as needed. During the dry season when water is scarce, water is still not sold but there are limits on the quantity that can be taken. Customary law is however not rigid. It recognizes that natural and public sources of water cannot satisfy a community's demand for water. It allows private

persons to invest in water. Such people can charge a fee for their investment. Presently, there is no regulation of water charges in the customary regime. Regulation may be unnecessary because there is not much demand for private water. Private persons may also have recognized the special nature of water and are not charging excessively for its provision. We shall therefore state that the customary regime takes a *realistic* view of water. It allows water from public sources to be available to all without charge. It recognizes however that demand can outstrip supply and therefore allows private persons to invest in water and charge a fee for supplying it to the market.

Does the customary regime solve the poverty problem? We shall argue that formally, it tries to solve the poverty problem. However, substantively, the customary regime actually aggravates poverty. Formally, water from natural and public sources is available free of charge. As long as water is available, a person is not denied access to it. The problem with the customary regime is that water from traditional sources is not available all year round. It is also the source of many diseases. It is also difficult to increase the supply of water to meet increased demand as populations grow. Sanitation under customary law is also seriously inadequate. Traditional methods of disposing waste cannot handle increased generation of waste. All these problems lead to diseases in rural areas and affect the economic lives of communities.

Within the traditional regime there are measures to control pollution as discussed earlier. Human settlements are located away from water sources. This, as we have already indicated, is to prevent water sources from being used as sewers. Within the limits of traditional knowledge, attempts are made to control pollution and maintain water quality.

How effective are these methods? The answer is that they are generally ineffective. With increasing population growth, many human settlements are now close to water sources. Poor land use methods have destroyed the environment. Mining and other industrial activities have generated waste that can be controlled under the customary regime. Generally as the population becomes educated and less homogeneous, taboos and religious sanctions that form the basis of the punishment under the customary regime are also ignored.

Under the customary regime, water conservation has been unnecessary because of the low level of technology that is used to get water. Traditional wells do not go deep enough to deplete the water table. The natural water cycle regulates the amount of water available. However, with the availability of sophisticated technology that can take water from underground, dam rivers and divert watercourses, traditional conservation methods are also likely to become irrelevant.

How does customary law deal with the gender problem? One would be tempted to say that customary water law is anti-female. Many of the rules affect women they are the primary collectors of water and these rules try to restrict their ability to get water. A critical look at the situation suggests that this is not actually the case.

However, while one can argue that customary law is not anti-female, it is definitely not profemale. In spite of the important role women play in collecting water, the customary regime gives them no special role in its management. Nothing is done to relieve the pain associated with collecting water. Women must trek long distances to get water especially during dry seasons. This affects their economic activity and quality of life.

The major strength of customary law is its attitude toward the issue of entitlement. It is an important lesson that has to be learnt. If we can graft it on to the CWSP, it would significantly improve its ability to solve the problem of poverty alleviation. Generally, we shall conclude that customary water law is not an adequate regime for supplying water to rural people in Ghana. It is unable to provide water all year round and is the source of diseases.

B.4 Impact of Water Law & Water Rights on Water Supply and Access

From the discussion, it is clear that the CWSP has been a relative success while the supply of water to urban areas has been a relative failure. Water supply under the customary regime still has serious challenges. The following questions should be answered ‘which conditions have enabled the CWSP to be a relative success?’

‘Why has the supply of water and sanitation to urban areas and under customary law been relatively unsuccessful?’

B.4.1 Enabling Conditions

The major enabling condition for the success of the CWSP has been the general consensus among stakeholders that increasing the supply of water and sanitation to the poor is essential if there is to be an improvement in their standard of living. Effective policies and strategies are being developed to implement the programme of providing water and sanitation for as many communities as possible. There is constant interaction among donors, the government and non-governmental organizations to review progress in the implementation of the CWSP.

As we have already indicated, the management of water resources is a multi-faceted enterprise that involves a number of users and institutions. Until recently, different

users and institutions pursued their mandates with very little co-ordination among themselves. There are now genuine attempts to integrate policies and strategies to ensure that the provision of rural water is managed in a multi-faceted manner. The CWSA consults with District Assemblies when it is developing its policies. Local communities are involved in the design and implementation of the programme. The private sector is given a prominent role to play in the supply of facilities. One of the major weaknesses of past water and sanitation programmes were attempts to separate water supply from sanitation issues. As part of the multi and inter-disciplinary approach to water management, water supply, sanitation and hygiene issues have been brought together to ensure that communities reap maximum health benefits from the programme.

What has increased public awareness on the importance of water and sanitation issues and has encouraged the government to tackle the problem seriously is the democratization process. Ghana has had two elections in the 1990s. The introduction of competitive politics has helped to improve the supply of water to rural areas in two ways. First, democracy has opened the political space and provided opportunities for civil society to contribute to debates on the future of the country. The press has highlighted the problem of access to water and sanitation for the poor in rural areas. Reports of outbreaks of guinea worm disease, cholera and typhoid fever are given prominence in the media. NGOs dealing with the provision of water are informing the public about its importance.

Democratization has also led to the need for government to develop new constituencies to bolster its position in the electoral system. Structural adjustment has generally had a negative impact on urban dwellers. Government is now developing its support in the rural areas. This has meant satisfying rural needs and one of the most important needs of the rural populace is the provision of clean water. Government has consequently taken seriously the supply of potable water to as many rural communities as possible (see Herbst J., 1993, for a discussion of the Politics of Reform).

Community participation in the provision of water and sanitation has also greatly improved the sustainability of water supply programmes. This bottom-up approach has many important dimensions. First it is a demand driven approach. Unless a community wants water and sanitation facilities, it does not become part of the programme. Communities must also be prepared to pay part of the cost of providing such facilities and they must be ready to pay for their maintenance. Water is sold to community members and the funds collected are used to maintain water and sanitation facilities. Tariffs are determined by WATSANs in consultation with communities. Community participation through the bottoms-up approach has been the greatest strength of the CWSP.

C. CUSTOMARY LAW AND TRADITIONAL WATER MANAGEMENT IN BURKINA FASO:

EXAMPLE OF THE YATENGA PROVINCE

Humankind and the environment were closely linked in traditional societies. In order to have a good understanding of environmental management as a whole in the province (kingdom) of Yatenga, it is necessary to consider the governing system, traditional religion, the origins of the migrations and the history of the settlement of villages.

1 - The governing system in the Yatenga kingdom

1.1- Traditional governance in the Yatenga kingdom

According to the Ramsa Naba (Minister of Youth and Protector of the kingdom's fetishes), the kingdom always had its own governing system, which is from top to bottom and from bottom to top. Such a system resisted the colonizer's attempts at reforms due to the great ascendancy of the Emperor over his faithful subjects, who in return must guarantee and defend the general interest of the kingdom. Thus the ambivalent application of the governing system justifies itself as follows:

from top to bottom: decisions are taken by the Emperor and communicated to the Zack-Naba (head of the family)

from bottom to top: the communities' concerns reach the Emperor through a hierarchical way. He will then gather the Ministers' Council to decide on solutions or sanctions to be applied. The Ministers' Councils gather every Friday at the royal palace in the presence of the following ministers

the Baloum, in charge of the King's protocol;

the Toogo Naba, Minister of Communication;

the Ouidi Naba, Minister of the Education and Defense of the Princes;and

the Rasam Naba, Minister of Youth and Protector of the fetishes, who is the central pillar of the system. He is considered as the second in hierarchy and is assisted by secretaries in charge of precise tasks.

1.2- Division of powers within the kingdom

In traditional Mossi societies, the principle of division of powers has existed since time immemorial and is regulated by the custom, the supreme law. Everybody knows about the division of powers as nobody ignores the rules enacted by the custom. The principle is applied as follows:

- *the Naba,* leader of the village, has administrative power over the subjects, at the level of the region. He is the intermediary between the Emperor of Yatenga and his people on one side, and, on the other side, he acts as an administrative relay, communicating messages and collecting the poll tax.

- *the Tengsoaba,* leader of the earth, is the guardian of the customary law of the village land. He organizes and chairs the ritual ceremonies linked to the earth. When a death occurs, he chooses the burial place in consultation with the Council of Elders. He acts as an intermediary between the people and the deities to ask for the community's

protection. In case of violation of the custom, sanctions are given to the guilty by the appropriate authorities (leader of the earth, Council of Elders) and expiatory and propitiatory sacrifices are undertaken in order to calm the spirits

- *The Council of Elders* acts as a consultative assembly to both authorities mentioned formerly; the opinion of the Council of Elders often influences the decisions of the Naba, leader of the village, and of the Tengsoaba, leader of the earth. The Council of Elders is composed of two "Chambers" the women and the men.

2. - Water in the Traditional Moagha Society of the Yatenga kingdom

2.1- Water as a universal myth

The water-life relationship is present in the founding myths of every culture. The flood theme is prevalent in all the three monotheistic religions, but, contrary to customary belief, it did not necessarily originate from, and is certainly not confined to, the Judeo-Christian- Islamic tradition. It is also found in the Hindu "vedas", in the Inca stories, in all the myths of Mesopotamia ("the earth between two floods") and in the animism of African societies. The founding myths influenced cultural behaviors and brought the notion of water as a sacred element. The cultural dimension is certainly indispensable and helps us understand the various conflicts surrounding around water and, eventually, to find pacific and fair solutions for its sharing and use.

2.2- Water and the animist religion

The traditional African societies were strongly animist before the implementation of Islam and Christianity. The animist religion is based on the natural elements, among which is water. These traditional societies have their own vision of water, which differs from the western water vision (art. 5 of the customary law of Yatenga). Water is used with a religious sense by users in order to evoke the spirits of the dead so that they watch over the community.

2.3- Water in the peoples' settlement

Lack of water in some regions drove the people to other cities where it is abundant. These last decades witnessed an important demographic upsurge on the riverbanks. The elders say that the search for water structured the people's settlement in the country. In the popular tales water often played a saving role in many societies.

2.4- Water as salvation for the migrant

Many societies show gratitude for natural elements, especially water, which at one point in their history granted them significant protection. The stories recalling these events are not exclusive to the traditional African societies. They are universal, as the holy scriptures or the popular tales of the oral societies testify.

There is the legend of Queen Abra Pokrou in the Gold Coast (now Ghana), in which water helped her to escape from her enemies after her designation to take over from her late father.

2.5- Village names with inspiration from water.

According to Sanon D. B. and Traore Y., (1999), the environment used to be a cultural source of inspiration in traditional societies. A toponymic approach reveals that the village names refer either to a plant-like entity, or to a watercourse, or to a particular relief existing before the people's settlement. In some societies, which consider water as a symbol of peace and life, certain villages have been given a name referring to water. The example mentioned hereafter refers to rivers or villages in western Burkina Faso, although all the communities in the country share the same vision with regard to their environment.

The "Zu" River has a mystic dimension, such as the bush or the mountain. It is considered as an expression of the world's dynamism, a strong, eventually suspicious, entity, hence the necessity for each village to make a pact with the river whose water it uses daily. The river is considered as the vital stream linking the present and past generations that lived on it. Apart from their mystic connotations, the various toponymies also convey a meaning, a projection of society and dictate particular behaviors.

Kofila: it is a village of the Madeira ethnic group in the north of the capital of the Houet province (Bobo), also the financial capital of Burkina Faso. This name comes from the local "Jula" dialect and is divided into two parts: "Ko", which means "river", and "Fila" which means "two". Effectively, the village is located at the confluence of two watercourses. These names of villages, places and regions represented projects of certain communities. In fact, the designated name act as a call for the defense and improvement of water as a natural element and of the conditions of life.

2.6- A system of water property

As an element in the democratization process that occurred during the 1990s, most African countries have established a legislative system governing various branches of industry, including water. Burkina Faso elaborated and implemented a law concerning the Land and Property Reform (RÈforme agraire et fonciÈre RAF). The 5th article of the RAF names the State as the owner of water. But in reality, this statement remains only a simple claim, as a real juridical prerogative will not be accepted without much resistance from the customary water users. Effectively, what characterizes African water law is the existence, parallel to written laws, of a corpus of customary norms which originated largely prior to colonization. In a country where the majority of the population is illiterate, these norms tend to supplant the official written legislation about to the exploitation of the water resources by the local inhabitants. During our interviews, the customary leaders affirmed that the State is powerless without those who know the true identity of the people. They therefore act as the intermediary between the population and administration. According to various customary leaders, no serious study, with the

ambition of creating acceptable juridical norms, could ignore the essential contribution of customary law.

2.7- Traditional water management: customary laws

In the traditional societies of the Yatenga kingdom, environmental management is the responsibility of the leader of the earth (article 9, 10 of Yatenga's customary code), who can be assisted by the elders or any other person designated to such as the Bouli Naba (customary leader in charge of water).

2.7.1- Prohibitions

According to the hierarchy of customary law, the prohibitions are above the totems, as they have a larger sphere of application. The prohibition represents what is not authorized by the community at the level of the village, that is what everybody forbids. Thus the following prohibitions can apply in a village and be severely sanctioned by the custom: Prohibition to sell water or refuse to let it be drawn from one's own well.

Prohibition to dirty the surroundings of the well

Prohibition to dig a well without the agreement of the relevant customary leaders.

2.7.2- Totems

They have a sphere of application restricted to the large family. Thus an ancestor, walking in search of a shelter, was saved by quenching his thirst at such a watercourse; he takes an oath about it, engaging his whole descent. This oath recommends that his descent watch-over the watercourse and carry-out sacrifices in order to thank the spirits of the place (Sanon D.B. & Traore Y., 1999).

2.8- Gender and traditional water management

In traditional societies the customary leaders were responsible for the management of water. Now the State also designates technicians for the same task. The distribution of the work now determines the participation by gender in the traditional societies of the Yatenga province. The males in the traditional societies are in charge of construction, maintenance and management of the work. During the work, girls and women constitute a stimulation force and organize in this perspective festivities on the building site in this regard. They praise the work accomplished, particularly of the young people who do the hardest tasks. Moreover, they play a predominant role in the domestic and horticultural spheres. In the traditional societies gender plays complementary roles

2.9- The causes of failure of water conveyance projects

In traditional societies the approaches that do not involve the population are very likely to

fail. Water management policy must take tradition into account in to achieve a sustainable vision. The ecological NGO, Green Cross Burkina Faso, understood this when it promoted its hydraulic policy through reevaluation of the traditional methods of water management. Several factors explain the causes of failure of water conveyance projects: the influence of ecological factors (advance of the desert, drought), demographic pressure (increase in water demand, contamination of water), economic and political aggression (inclusion of the economic dimension in social relationships, territorial conflicts). The mastery of local knowledge in water management, and the understanding of the sources of conflicts are necessary for the elaboration of sustainable projects and will favor partnership between traditional and modern water management.

Bibliography

Acquah, B. K. P. (2002) "Investigation of Groundwater Availability in the Volta Basin of Ghana, Based on Hydrogeological Field Data and Geological Maps Using GIS." Masters Thesis, Institute of Groundwater and Hydraulic Engineering, University of Stuttgart, Stuttgart.

Andreini, M.; van de Giesen, N.; van Edig, A.; Fosu, M.; Andah, W. (2000) « Volta Basin Water Balance. » ZEF – Discussion Papers on Development Policy, No. 21, ZEF, Bonn.

ACOPS. WGPI (Working Group on Programme on Interventions): WGPI-I (2001); WGPI-II (2002).

ACOPS. WGPIA (Working Group on Integrated Problem Analysis): WGPIA-I (2000); WGPIA-II (2001); WGPIA-III (2001).

Andreini M., Vlek, P. D. Nick van de Giesen, 2002. Water sharing in the Volta basin. Friend (2002) – Regional Hydrology: Bridging the Gap between Research and practice. IAHS Publ. No. 274.

Andreini M., van de Geisen, N., van Edig, A., Fosu, M. and W. Andah, 2000. Volta Basin Water Balance. ZEF Discussion Papers on Development and Policy No. 21.

Armah, A. K. and D. S. Amlao. (1998). Coastal Zone Profile of Ghana. Gulf of Guinea Large marine Ecosystem Project. Ministry of Environment, Science and Technology. Accra, Ghana.

Banoeng-Yakubo, B.; Skjerna, L. (2000) "Applications of Remote Sensing and Geographical Information System to Hydrogeological Studies in the Upper West Region, Ghana." In: Sililo et al. (Eds.), Groundwater: Past Achievements and Future Challenges, Balkema, Rotterdam.

BR5GM (1991) « Exploitation des Eaux Souterraines en Socle Cristallin et Valorisation Agricole. Pilote Expérimentale en Milieu Rural pour les Zones Soudano-Sahéliennes. Synthèse des Etudes de la Phase II (Rapport Final). » BRGM, Orléans.

Bromley, J.; Edmunds, W. M.; Fellmann, E. (1997) "Estimation of Rainfall Inputs and Direct Recharge to the Deep Unsaturated Zone of Southern Niger Using the Chloride Profile Method." Journal of Hydrology, 188-189, 1-4, 139-154, Elsevier.

CIDA (1998) "Basic Human Needs Performance Review. From Service Delivery toward Governance for Sustainability Report of the Ghana Water Program – Evaluation Report." CIDA, Quebec.

Danielson, J. J. (1996) "Delineation of Drainage Basins from 1 km African Digital Elevation Data." In: Pecora Thirteen, Human Interactions with the Environment –

Perspectives from Space, Sioux Falls, South Dakota, August 20-22, 1996.

Dapaah-Siakwan, S.; Gyau-Boakye, P. (2000) "Hydrogeologic framework and borehole yields in Ghana." *Hydrogeology Journal*, 8, 405-416.

Darko & Krasny, 2003

Deichmann, U. (1994) "A Medium Resolution Population Database for Africa, Database Documentation and Digital Database." NCGIA, University of California, Santa Barbara. Through: UNEP/GRID at Sioux Falls. Data available from:

<http://grid2.cr.usgs.gov/datasets/datalist.php3>.

Desconnets, J. C.; Taupin, J. D.; Lebel, T.; Leduc, C. (1997) "Hydrology of the HAPEX-Sahel Central Super-Site: surface water drainage and aquifer recharge through pool systems." *J. Hydrology*, 188/189, 1-4, 155-178.

DGH (Direction Générale de l'Hydraulique) (1997) "Inventaire des Points d'Eau Modernes au Burkina Faso." Ministère de l'Environnement et de l'Eau, Ouagadougou.

DGH (Direction Générale de l'Hydraulique) (2001) « Etat des Lieux des Ressources en Eaux du Burkina Faso et de leur Cadre de Gestion. » Ministère de l'Environnement et de l'Eau, Ouagadougou.

Dieng, B. ; Bazie, P. ; Schmitt, A. (1991) « Transfert d'Eau en Milieu Poreux non Saturé. Recharge des Nappes en Climat Soudano-Sahélien. » In : Utilisation Rationnelle de l'Eau des Petits Bassins Versants en Zone Aride, 131-137, AUPELG-UREF, Paris.

Diluca, C. ; Muller, W. (1985) « Evaluation Hydrogéologique des Projets d'Hydraulique en Terrains Cristallins du Bouclier Ouest Africain. » BGR, Hannover.

Edmunds, W. M.; Fellmann, E.; Goni, I. B.; Prudhomme, C. (2002) "Spatial and Temporal Distribution of Groundwater Recharge in Northern Nigeria." Hydrogeology Journal, 10, 205-215.

FAOSTAT, on-line database, FAO, Rome
(http://www.fao.org/waicent/portal/statistics_en.asp).

Ghana Geological Survey (1964) "Mineral Resources Map – North-West part of the Wa sheet (1:250,000)." GSD, Accra.

Ghana Geological Survey (1973) "Geological Map – Southwestern part of the Wa sheet (1:250,000)." GSD, Accra.

Ghana Geological Survey (1991 a) "Geological Map – Northern part of Bole sheet (1:250,000)." GSD, Accra.

Ghana Geological Survey (1991 b) "Geological Map – Southern part of Bole sheet (1:250,000)." GSD, Accra.

Gleick, P. (1996) "Basic Water Requirements for Human Activities: Meeting basic needs." Water International, 21, 83-92.

Gombert, P. (1998) "Synthèse sur la Géologie et l'Hydrogéologie de la Série Sédimentaire du Sud-Ouest du Burkina Faso, Programme RESO, Sous-Programme 'Ressources en Eau'." IWACO, Ouagadougou.

Gyau-Boakye, P.; Tumbulto, J> W> (2000) "The Volta Lake and Declining Rainfall and Streamflows in the Volta Basin." Environment, Development and Sustainability, 2, 1-10.

Gyau-Boakye, P. (2001) "Sources of Rural Water Supply in Ghana." Water International, 26, 1, 96-104.

IGP (1999) "Promotion of District Capitals 1 (Ejura, Kintampo & Nkoranza) – Hydrogeological Investigations, Technical Report." Ministry of Local Government and Rural Development, Accra.

Inter-African Committee for Hydraulic Studies (1986) "Explanatory Notice and Recommended Usage of the Map of Potential Groundwater Resources in Western and Central Africa 1: 5,000,000." ICHS, BRGM, Orleans.

Kortatsi, B.K. (1984) "Groundwater Utilization in Ghana." In: Future Groundwater Resources at Risk (Proceedings of the Helsinki Conference), IAHS Publication No. 222, IAHS Press, Wallingford.

Kwei, C. A. (1997) "Evaluation of Groundwater Potential in the Northern Region of Ghana." CIDA, Quebec.

Leduc, C.; Desconnets, J. C. (1994) "Variability of Groundwater Recharge in the Sahel: Piezometric Survey of the Continental Terminal Aquifer (Niger)." In: Future Groundwater Resources at Risk (Proceedings of the Helsinki Conference), IAHS Publication No. 222, IAHS Press, Wallingford.

Milville, F. (1991) "Etude Hydrodynamique et Quantification de la Recharge des Aquifères en Climat Soudano-Sahélien: Application a un Bassin Expérimental au Burkina Faso." In: Soil Water Balance in the Sudano-Sahelian Zone, IAHS Publication No. 199, Proceedings of an International Workshop, Niamey, 18-23.02.1991, IAHS Press, Wallingford.

Ministry of Works and Housing (1998) "Water Resources Management Study, Information 'Building Block' Study. Part II, Volta Basin System, Groundwater Resources." Ministry of Works and Housing, Accra.

Ministry of Works and Housing (2001) "2000 Population & Housing Census – Summary Report on Final Results." Ministry of Works and Housing, Accra.

Minor, T. B., Carter, J. A., Chesley, M. M., Knowles, R. B. (1994) "An Integrated Approach to Groundwater Exploration in Developing Countries Using GIS and Remote Sensing", ASPRS/ACSM, <http://www.odyssey.maine.edu/gisweb/spatdb/acsm/ac94048.html>.

New, M., Hulme, M. and Jones, P. D. (1999) "Representing Twentieth Century Space-Time Climate Variability. Part 1: Development of a 1961-90 Mean Monthly Terrestrial Climatology." *J. Climate* 12, 829-856 (dataset available through the IPCC DDC).

ONEA (1999) « Rapport Technique d'Exploitation, Exercice 1998. » Office National de l'Eau et de l'Assainissement, Ouagadougou.

Taylor, K. C., Minor, T. B., Chesley, M. M., and Matanawi, K. (1999) "Cost effectiveness of well site selection methods in a fractured aquifer." *Ground Water* 37/2, 271-274.

Thierry, D. (1990) "Analysis of Long Duration Piezometric Records from Burkina Faso to Determine Aquifer Recharge." In: D. N. Lerner, A. S. Issar, I. Simmers; *Groundwater Recharge – A Guide to Understanding and Estimating Natural Recharge; International Contributions to Hydrogeology, &, International Association of Hydrogeologists*. Heise, Hannover.

Wardrop & Associates Ltd. (1977) "CIDA Ghana Upper Region Water Supply Project, Phase 1, Hydrogeological Report Based on 1460 Well Supplies." CIDA, Quebec.

Wardrop Engineering (1987) "Report on Shorting of Wells in the Upper Regions of Ghana." Unpublished report.

WRI-Water Resources Research Institute / DANIDA (1993) "Rural Drinking Water Supply and Sanitation Project in the Volta Region – Final Report on the Inventory and Assessment of Potential for Hand Dug Wells in the Volta Region, Vol I – Main Report." WRI, Accra.

N'Djim, H and Doumbia, B. 1998. Case study: Mali Population and water issues. In: Alex B. Sherbinin and Victoria Dompka (Eds.). *Water and Population Dynamics: Case studies and policy implications*. <http://www.aaas.org/international/ehn/waterpop/mali.htm>

Telmo, A. 2002. A water Supply and Sanitation Study of the Village of Gouansolo in Mali, West Africa. MSc Report. Michigan Technological University, USA. http://www.cee.mtu.edu/peacecorps/documents_november_02/andrea_telmo.pdf

CIA (Central Intelligence Agency), 2004. *The World Factbook—Mali*. <http://www.cia.gov/cia/publications/factbook/geos/ml.html>

Bracken, P. and Mang, H.P. 2002. Sanitary systems in Ouagadougou, Burkina Faso: Current practices and future potential. <http://www.gtz.de/ecosan/download/PatrickBracken-traineereport.pdf>

Ministry of Works and Housing. 1998. Ghana's Water Resources: Management Challenges and Opportunities. Water Resources Management Study.

Gyau-Boakye, P and Dapaah-Siakwan, S. 1999. Groundwater: solution to Ghana's rural water supply industry? Water Resources Research Institute

Kankam-Yeboah, K. 1987. Status Report on the pilot scheme in Groundwater Utilization in the Accra plains. WRRI-CSIR Accra, Ghana.

Kortatsi, B.K. 1994. Groundwater utilization in Ghana, in: Future Groundwater Resources at Risk (Proceedings of the Helsinki Conference), IAHS Publication no. 222, 1994.

Cornish, G.A. and Aidoo, J.B. 2000. Informal Irrigation in the Peri-urban zone of Kumasi, Ghana: Findings from an initial questionnaire survey. Report OD/TN 97. HR Wallingford, UK.

Andah, A.I. 1993. Groundwater Utilization for small-scale Irrigation in the Accra Plains: Kponkpo and Ablekuma pilot schemes. Water Resources Research Institute (C.S.I.R). Accra, Ghana.

Agodzo, S.K. 2003. Shallow well irrigation for shallot farming in the Keta Lagoon Area of Ghana. Journal of the Ghana Institution of Engineers. Volume 1, Number 1:pp1-6

Nathan Consortium, 1970. Framework for River Basin Planning. Ghana Sector Studies. Interim Report for Ministry of Finance and Economic Planning.

Nicholson, S.E., 1983. Sub-Saharan rainfall in the years 1976 – 80: Evidence of continued drought. Monthly Weather Review. III. 1646 – 1654.

ACOPS. WGPI (Working Group on Programme on Interventions): WGPI-I (2001); WGPI-II (2002)

ACOPS. WGPIA (Working Group on Integrated Problem Analysis): WGPIA-I (2000); WGPIA-II (2001); WGPIA-III (2001).

Andreini M., Vlek, P. D. Nick van de Giesen, 2002. Water sharing in the Volta basin. FRIEND 2002 – Regional Hydrology: Bridging the Gap between Research and practice. IAHS Publ. No. 274.

Andreini M., van de Geisen, N., van Edig, A., Fosu, M. and W. Andah, 2000. Volta Basin Water Balance. ZEF Discussion Papers on Development and Policy No. 21.

Armah, A. K. and D. S. Amlao. 1998. Coastal Zone Profile of Ghana. Gulf of Guinea Large marine Ecosystem Project. Ministry of Environment, Science and Technology. Accra, Ghana.

CONAGESE 10 BP 6486 Ouagadougou 10 – National Action Plan for the Fight Against Desertification, Burkina Faso – 1999.

Cote d'Ivoire, 2001. Plan of Installation of the National Park of Comoe and of Development of its Periphery, 2002-2001.

Gyau-Boakye and P. Tumbulto, 2000. The Volta Lake and declining rainfall and streamflow in the Volta River basin. *Environment, Development, and Sustainability*, 2, 1 – 10.

GCLME. UNIDO/UNDP/NOAA/UNEP Guinea Current Large Marine Ecosystem, Review of Existing Information and Recommendations on Transboundary Priority Issues: Regional Report for Stocktaking Meeting, May 2001.

GCLME. INTEGRATED ENVIRONMENTAL AND LIVING RESOURCES MANAGEMENT IN THE GULF OF GUINEA – The Large Marine Ecosystem Approach. Proceedings of the first Regional Symposium on the Gulf of Guinea Large Marine Ecosystem. GOG (1998).

GCLME. PERSPECTIVES IN INTEGRATED COASTAL AREAS MANAGEMENT IN THE GULF OF GUINEA. Gulf of Guinea Large Marine Ecosystem Project. GOG (1998).

GCLME. STATE OF THE COASTAL AND MARINE ENVIRONMENT OF THE GULF OF GUINEA. Gulf of Guinea Large Marine Ecosystem Project. GOG (1998)

GCLME. TOWARDS INTEGRATED COASTAL ZONE MANAGEMENT IN THE GULF OF GUINEA. Gulf of Guinea Large Marine Ecosystem Project. GOG (1998).

GCLME. GEF: REGIONAL REPORT FROM THE STOCKTAKING WORKSHOP (May 2001) – PDF – B “Development of a Strategic Action Program (SAP) for the Guinea Current Large Marine Ecosystem (GCLME)”

Mali, 1983. Inventory Project of Terrestrial Resources.

Minister of Environment and Water (BURKINA FASO). – National Monograph on the Biological Diversity of Burkina Faso – Ouagadougou, February 1999.

Moniod, F., B. Pouyaud, and P. Sechet, 1977. Le Bassin du Fleuve Volta. Monographies Hydrologiques Orstom, No. 5. Paris.

Nathan Consortium, 1970. Framework for River Basin Planning. Ghana Sector Studies. Interim Report for Ministry of Finance and Economic Planning.

Nicholson, S. E., 1983. Sub-Saharan rainfall in the years 1976 – 80: Evidence of continued drought. *Monthly Weather Review*. III. 1646 – 1654.

Opoku-Ankomah, Y., 2000. Impacts of Potential Climate Change on River Discharge in *Climate Change Vulnerability and Adaptation Assessment on Water Resources of Ghana*. Water Research Institute (CSIR), Accra. Ghana.

Opoku-Ankomah, Y., 1998. Volta Basin System Surface Water Resources in Water

Resources Management Study. Information Building Block. Part II, Vol. 2. Ministry of Works and Housing. Accra. Ghana.

UNEP, 2001. Ghana Country Report for the Integrated Management of the Volta River Basin Project, Accra.

UNEP, 2001. Mali Country Report for the Integrated Management of the Volta River Basin Project, Accra.

UNEP, 2001. Togo Country Report for the Integrated Management of the Volta River Basin Project, Accra.

UNEP, 2001. Benin Country Report for the Integrated Management of the Volta River Basin Project, Accra.

UNEP, 2001. Burkina Faso Country Report for the Integrated Management of the Volta River Basin Project, Accra.

UNEP, 2001. Cote d'Ivoire Country Report for the Integrated Management of the Volta River Basin Project, Accra.

UNEP, 2001. Report of the first Regional Meeting of the National Coordinators for the Preparatory Process of a full GEF Project Proposal, Accra, Ghana, June 5 – 8.

UNEP, 2001. Some Technical Information and country Reports, Integrated Management of the Volta River Basin, PDF B. Accra, Ghana, June, 5 – 8.

UNEP, 2001. Presentations on other Initiatives. First Regional Meeting of the National Coordinators. Integrated Management of the Volta River Basin, PDF B. Accra, Ghana, June 5 – 8.

UNEP, 2001. Proceedings of the first Steering Committee Meeting. Integrated Management of the Volta River Basin. Accra, Ghana, June 8, 2001.

UNEP, 2001. Report on the Second Regional Meeting of the National Coordinators. Integrated management of the Volta River Basin, PDF B. Accra, Ghana, November 27 – 30.

UNEP, 2001. Some technical Information. Second Regional Meeting of the National Coordinators. Integrated Management of the Volta River Basin, PDF B. Accra, Ghana November 27 – 30.

UNEP, 2001. Proceedings of the Second Steering Committee Meeting. Integrated Management of the Volta river Basin. Accra, Ghana, November 30.

UNEP, 1999. Regional Seas Reports and Studies No. 171, "Overview of Land-based

Sources and Activities Affecting the Marine, Coastal and Associated Freshwater Environment in the West and Central African Region.”

UNEP/GEF/VOLTA/SCM.1/1, 2001. Proceedings of the First Steering Committee meeting. Accra.

UNEP/GEF/VOLTA/SCM.2/1, 2001. Proceedings of the Second Steering Committee meeting. Accra.

UNEP/GEF/VOLTA/SCM.1/1, 2001. Report of the First Regional Meeting of the National Coordinators for the preparatory process of a full GEF project proposal. Accra.

UNEP/GEF/VOLTA/SCM.2/1, 2001. Report of the Second Regional Meeting of the National Coordinators. Accra.

Water Resources Management Study (WARM), 1997. Information Building Block, Part III, Vol. 1.

World Bank. World Development Report 2000/2001.